



- Bedrock
- Buried bedrock ridge
- Fan delta (a) splay (b) channels
- Blocky (a) coarse (b) talus
- Distributary mouth bar
- Channel
- Scarps
- Slump scarp
- Fiord floor terrace
- Lineations
- Flow-transverse steps and ridges
- Pressure ridges
- Prominent ridge crest
- a) flute (b) depression
- Spillover lobe
- Steep slope
- Hummocky debris flow
- Shallow linear gullies
- High-relief dendritic gullies

SEAFLOOR MORPHOLOGY – Bute Inlet, British Columbia

Brian D. Bornhold
Pacific Geoscience Centre
Geological Survey of Canada

and

David B. Prior
Coastal Studies Institute and
Department of Geology & Geophysics
Louisiana State University

These maps depict the seafloor morphology related to the principal Holocene sediment transport systems in Bute Inlet. The interpretations are based on sidescan sonar and high-resolution seismic surveys conducted between 1984 and 1986.

References: Prior, D.B., Bornhold, B.D. and Johns, M.W., 1986. Active sand transport along a fjord-bottom channel, Bute Inlet, British Columbia: *Geology*, v. 14, p. 581-584.
Prior, D.B., Bornhold, B.D., Wiseman, W.J. and Lowe, D.R., 1987. Turbidity current activity in a British Columbia fjord: *Science*, v. 237, p. 1330-1334.

Prior, D.B. and Bornhold, B.D., 1989. Submarine morphology and processes of fjord fan deltas and related high-gradient systems: modern examples from British Columbia: in Nemec, W. and Steel, R.J. (Eds.), *Fan Deltas: Sedimentology and Tectonics Settings*, Blackie and Sons, p. 125-143.

GSC OPEN FILE 1989
Sheet 2 of 2