

This document was produced  
by scanning the original publication.

Ce document est le produit d'une  
numérisation par balayage  
de la publication originale.

COMMISSION GÉOLOGIQUE DU CANADA



GEOLOGICAL SURVEY OF CANADA

**OVERVIEW, LEGEND AND MINERAL DEPOSIT TABULATIONS FOR:  
GEOLOGICAL SURVEY OF CANADA**

**O.F. 2188**

**GEOLOGY OF SW KLUANE LAKE MAP AREA  
(115G & F[E½]), YUKON TERRITORY**

**O.F. 2189**

**GEOLOGY OF MOUNT ST. ELIAS MAP AREA  
(115B & C[E½]), YUKON TERRITORY**

**O.F. 2190**

**GEOLOGY OF SW DEZADEASH MAP AREA  
(115A), YUKON TERRITORY**

**O.F. 2191**

**GEOLOGY OF NE YAKUTAT (1140) AND TATSHENSHINI RIVER (114P)  
MAP AREAS, BRITISH COLUMBIA**

**C.J. Dodds and R.B. Campbell  
1992**



**Énergie, Mines et  
Ressources Canada**

**Energy, Mines and  
Resources Canada**

**Canada**

## CONTENTS

	Page #
--	--------

### OVERVIEW AND LEGEND

Introduction	1
Acknowledgments	3
Geology overview	4
Geological setting	4
Economic geology	11
History	12
Placer deposits	13
Lode deposits	15
Nonmetallic sedimentary deposits	18
Stratified rock map unit descriptions	20
Intrusive rock map unit descriptions	32
Map symbols	38
Notes on map unit descriptions	39
Mapping sources	41
References	72

### Figures

1. Map area index	2
2. Terranes, major faults, cover rocks, plutonic rocks, and morphogeological belts (inset map), Saint Elias Mountains and adjoining areas and legend	5
3. National topographic system (NTS) grid, Saint Elias Mountains and adjoining areas	40

### Tables

1. Occurrence of map units in terranes and cover rocks	7
2. Mineral deposits SW Kluane Lake map area (115G & F(E1/2))	43
3. Mineral deposits Mount St. Elias map area (115B & C(E1/2))	54
4. Mineral deposits SW Dezadeash map area (115A)	57
5. Mineral deposits of Tatshenshini River (114P) and NE Yakutat (114O) map areas	62

### MAPS

- O.F. 2188: Geology of SW Kluane Lake map area (115G & F(E1/2)), Yukon Territory;  
1 sheet at 1:250 000 scale
- O.F. 2189: Geology of Mount St. Elias map area (115B & C(E1/2)), Yukon Territory;  
1 sheet at 1:250 000 scale
- O.F. 2190: Geology of SW Dezadeash map area (115A), Yukon Territory; 1 sheet at  
1:250 000 scale
- O.F. 2191: Geology of NE Yakutat (114O) and Tatshenshini River (114P) map areas,  
British Columbia; 2 sheets at 1:250 000 scale

**GEOLOGY OF SW KLUANE LAKE (115G & F(E1/2) (O.F. 2188),  
MOUNT ST. ELIAS (115B & C(E1/2) (O.F. 2189), SW DEZADEASH (115A) (O.F. 2190),  
& NE YAKUTAT (114O) AND TATSHENSHINI RIVER (114P) (O.F. 2191) MAP AREAS,  
SOUTHWESTERN YUKON AND NORTHWESTERN BRITISH COLUMBIA**

## INTRODUCTION


These open files are the result of geological investigations during Operation Saint Elias, a project undertaken by the Geological Survey of Canada intermittently during the 1970s and 1980s to complete geological mapping of the Saint Elias Mountains of Canada. They update and revise previously issued open files on the area (Campbell and Dodds, 1982a,b,c, and 1983a,b), and include 1:250 000 scale geological maps with fossil and numerical age data locations accompanied by a single comprehensive legend common to all sheets, and mineral deposit locations and data tabulations.

The map areas are located in southwesternmost Yukon and northwesternmost British Columbia (Fig. 1) and encompass the entire Saint Elias Mountains within Canada. The mountains are the highest and among the youngest in North America, and contain the world's largest nonpolar icefields. Much of the region is extremely rugged with 20 peaks cresting 4200 m. (14 000 ft.) which include Canada's highest Mount Logan (5951 m.; 19 525 ft.). Most of Saint Elias Mountains in Yukon are within Kluane National Park Reserve and Kluane Game Sanctuary and the area as a whole affords spectacularly beautiful mountain scenery and a wide variety of fauna and flora (Theberge, 1980). The Alaska Highway and Haines Road bound the map areas to the northeast, however there are no roads in the region and access is by aircraft, boat or foot only.

Previous geological mapping in Saint Elias Mountains was undertaken mainly in the more accessible, easternmost parts of the region. Earlier exploratory mapping was carried out in various parts of SW Kluane Lake by Brooks (1900), McConnell (1905), Cairnes (1915a), Sharp (1943) and Bostock (1952), followed by the first regional geological synthesis of about half of the area by Muller (1967). The major past contribution to geological mapping in Mount St. Elias was that in the northeasternmost corner of the sheet by Wheeler (1963). Initial geological explorations were undertaken in SW Dezadeash by Cockfield (1928) and Mandy (1933), however the first regional geological map and synthesis of the area was that of Kindle (1953). Earlier geological mapping in Tatshenshini River sheet were conducted by Mandy (1933) and Watson (1948) in Squaw Creek and Rainy Hollow areas, whereas no previous mapping has been recorded from NE Yakutat.

Prior to Operation Saint Elias more than seventy percent of the region was essentially unstudied. The objective of the project was to update previous work and complete geological mapping of Saint Elias Mountains in Canada. Helicopter supported field work was undertaken approximately during 2 weeks in 1973, 3 months in 1974 and 1977, 2 months in 1978, 1 month in 1979, 2 weeks in 1981 and 1983, and 3 weeks in 1986 and 1987. Due to the extremely rugged, and extensively ice-covered and remote terrane, most of the Icefield and Alsek ranges were reconnaissance-mapped by R.B. Campbell from helicopter fly-by and landing-site observations. Detailed ground coverage was

## MAP AREA INDEX



**Figure 1.** O.F. 2188 SW Kluane Lake (115G & F(E1/2)); O.F. 2189 Mount St. Elias (115B & C(E1/2)); O.F. 2190 SW Dezadeash (115A); and O.F. 2191 NE Yakutat (114O) and Tatshenshini River (114P) map areas.

restricted mainly to the relatively accessible eastern perimeter of the region (for major contributors see Mapping Sources). Much of the data involved in the compilation of these open files is from the many participants in Operation Saint Elias: geological mapping and special studies (Campbell and Dodds, 1975, 1978, 1979, 1982a,b,c, and 1983a,b; Campbell and Eisbacher, 1974; Dodds, 1988; Dodds and Campbell, 1988; Eisbacher, 1975, 1976; Eisbacher and Hopkins, 1977; Read, 1976; Read and Monger, 1975, 1976; Souther, 1974; and Souther and Stancui, 1975), and thesis studies (Campbell, 1981; Downey, 1978; Downey et al., 1980; Jacobson, 1979; Jacobson et al., 1980; Lowey, 1980; Sturrock, 1975, Sturrock et al., 1980). Only a few mineral deposits were visited by the authors. Data on them was mostly from literature surveys and where possible from discussions with government, university and mining industry geologists.

### Acknowledgments

The authors wish to thank the many individuals involved in the success of Operation Saint Elias. They include both those who worked directly on the project, and those who indirectly contributed generously their exchange of ideas, data and services.

Very capable field assistance was given by the following: F. Taxboc, J. Mitchell, L. Werner, and T. Booth in 1974; J. Dudley, M. Downey, C. Roots, J. Gaspard, and K. Pivnick in 1977; G. Lowey, B. Jacobson, D. Wood, D. Stone, and W. Walker in 1978; M. Eckervogt in 1979; P. Benham in 1986; and S. Dennison in 1987. Very safe and efficient helicopter support was provided by the following companies and (pilots): Transwest Helicopters Ltd. (R. Dunbar, T. Okusako, and L. Smailles) in 1974 and 1977; Shirley Helicopters Ltd. (B. Cleveland) in 1978; Yukon Helicopters Ltd. (M. Duff) in 1979; Pacific Helicopters Ltd. (J. Cadham) in 1983; Tundra Helicopters Ltd. (D. Cassidy) in 1986 and 1987; and Airlift Corp. (J. Clancy) in 1987. Cooks who served excellent fare were Janet Green and Gail Dodds in 1974; Marg Yandel in 1977; and Nancy Hoyle in 1978.

The authors particularly wish to thank D.A. Brew, H.C. Berg, E.M. MacKevett Jr., and G. Plafker of the U.S. Geological Survey (Alaska Branch, Menlo Park), who gave freely of their data and ideas during field trips to the map areas, to McCarthy and Yakutat Quadrangles, on two extensive field trips throughout SE Alaska, and in many other written and verbal discussions.

Macro fossil identifications were by E.W. Bamber, P. Harker, J.A. Jeletzky, D.J. McLaren, B.S. Norford, A.W. Norris, A.E.H. Pedder, T.P. Poulton, H.W. Tipper, E.T. Tozer (all of Geological Survey of Canada); and also by G.A. Cooper (National Museum of Natural History - Smithsonian Institution), R.H. Flower, and B.D.E. Chatterton (University of Alberta, Edmonton). Conodont identifications were mostly by M.J. Orchard; some were also by B.E.B. Cameron, T.T. Uyeno, and S.M.L. Pohler (all of Geological Survey of Canada). Age determinations were made mostly by R.K. Wanless, R.D. Stevens and other staff of the Geological Survey of Canada; a few were also by D.L. Sturrock, M.E. Downey, and B. Jacobson as part of B.Sc. thesis studies at University of British Columbia, Vancouver.

The authors wish to thank T. Bremner and other staff at Indian and Northern Affairs Canada, Whitehorse for mineral deposit data and editorial assistance; and also D.G. MacIntyre and T.G. Schroeter of British Columbia Ministry of Energy, Mines, and Petroleum Resources for sharing geological information and ideas on the British Columbia part of the map areas. They are also indebted to Falconbridge Ltd. (J.J. McDougall, J.B. Gammon, and T. Chandler), Geddes Resources Ltd. (D. Little, R. Beckett, Mary P. Webster, and J. Ricker), Stryker Freeport Resources Ltd. (D. Perkins, A. Wilkinson, and G. della Val) for some logistical support, and exchange of ideas and data; and also Archer, Cathro, and Associates Ltd. (R. Carne) for discussions on mineral deposits of the region.

## GEOLOGICAL OVERVIEW

### Geological Setting

The Saint Elias Mountains in Canada comprise the Icefield, Donjek, Kluane, Fairweather, Alsek, Squaw, Datlasaka and Kusawak ranges. Large valley glaciers including the Klutlan, Donjek, Logan, Seward, Hubbard, Kaskawulsh, Lowell, Fisher, and Tweedsmuir, flow from the vast ice-cover of the Icefield Ranges. Major, heavily-laden, milky, extensively braided rivers such as the White, Donjek, Alsek and Tatshenshini, transect the region. The mountains lie along the southwestern edge of the Coast Belt and are within the northern extension of the Insular Belt in southwestern Yukon and northwestern British Columbia and adjacent Alaska (Fig. 2, inset). For the most part they are southwest of Denali Fault System. The Saint Elias Mountains are divisible into six northwest-trending, major-fault bounded geological terranes with differing cover rock sequences, all of which extend into adjoining Alaska (Fig. 2 and Table 1). These comprise the Alexander Terrane, three segments of Wrangellia (W1, W2, and W3), the Chugach Terrane and a small sliver of the Yakutat Terrane. The Saint Elias Mountains are dominated by the Alexander Terrane. Wrangellia segments (W2 and W3) may have been moved northeast of Alexander Terrane by large dextral displacements along Duke River and Denali faults (Campbell and Dodds, 1983c). The terranes are characterized by distinctive plutons, metamorphic rocks, structural styles and their accretion is believed to have occurred mostly in Mesozoic to Cenozoic time (Berg et al., 1972, 1978; Coney et al., 1980; Campbell and Dodds, 1983c; Monger and Berg, 1984; Wheeler et al., 1988; Plafker et al., 1989). Large scale, both syn- and post-accretion, mainly dextral transcurrent faulting, has further complicated the geology of the region.

The Saint Elias Mountains are predominantly underlain by the Alexander Terrane, which is also extensively exposed throughout adjacent southeastern Alaska (Gehrels and Berg, 1984; Gehrels and Saleeby, 1987). It consists of thick sequences of mainly Paleozoic strata, but which range in age from Late Cambrian and earlier (?) to latest Triassic. Cambrian to lowermost Ordovician and older (?) rocks comprise abundant mainly marine basic volcanics, volcaniclastics and minor carbonate. Those of Ordovician-Silurian age include widespread laminated and thicker bedded carbonates, calcareous clastics and rare basic volcanics. Devonian rocks consist of shallow-marine carbonate, fine clastics and local basic volcanics. Undifferentiated mostly marine Devonian to latest Triassic age strata include fine clastics, basic volcanics, carbonates, and rare evaporites.

TERRANES, MAJOR FAULTS, COVER ROCKS, PLUTONIC ROCKS, AND  
MORPHOGEOLOGICAL BELTS (INSET MAP), SAINT ELIAS MOUNTAINS AND ADJOINING AREAS  
(modified from Wheeler and McFeely, 1987)




Figure 2.

## LEGEND

## COVER ROCKS

Tertiary

Tvs

nonmarine volcanics and sediments  
(mostly Wrangell Lava).

Cretaceous

Ks

shallow marine sediments.

Upper Jurassic-Lower Cretaceous

JKs

flysch, local volcanics (Gravina-  
Nutzotin).

JKm

metasediments ("Kluane schist", and  
older rocks).

## TERRANES

YA

YAKUTAT (local Prince William).

CG

CHUGACH.

W1,W2

WRANGELLIA (W1, W2).

W3,TU

WRANGELLIA (W3), TAKU.

W1?

WRANGELLIA (W1)?

WM

WINDY-MCKINLEY.

NS

NISLING (local undivided rocks).

AX

ALEXANDER.

## PLUTONIC ROCKS

Tertiary

Mg

Wrangell suite (6-16 Ma):  
subvolcanics and granitoids.

Og

Tkope suite & others (23-33 Ma):  
granitoids and subvolcanics.

Eg

Seward suite (41-52 Ma).

Cretaceous-Tertiary

KTg

Coast Plutonic Complex (undivided).

KTb

gabbro-diabase plutons.

Late Early Cretaceous

Kg

Kluane Ranges suite (106-121 Ma),  
& Alaskan-type mafic-ultramafics.

Late Jurassic-earliest Cretaceous.

JKg

Saint Elias suite (130-160 Ma).

Late Pennsylvanian-Early Permian

PPg

Icefield Ranges suite (270-290 Ma).

## MAJOR FAULTS

T.F.S.	-	Transition Fault System.
C.F.S.	-	Contact Fault System.
C.S.F.S.	-	Chugach-St. Elias fault system.
F.F.	-	Fairweather Fault.
B.R.F.S.	-	Border Ranges Fault System.
C.F.	-	Chitina Fault.
H.F.	-	Hubbard Fault.
D.R.F.	-	Duke River Fault.
T.F.	-	Totschunda Fault.
D.F.S.	-	Denali Fault System.
C.S.F.	-	Chatham Strait Fault.

## OCCURRENCE OF MAP UNITS IN TERRANES AND COVER ROCKS

TERRANES	STRATIFIED UNITS	INTRUSIVE UNITS		STRATIFIED AND INTRUSIVE UNITS IN VARIOUS COVER ROCKS			
		GRANITOID	MAFIC/ULTRAMAFIC	JURA-CRETACEOUS COVER ROCKS	TERTIARY COVER ROCKS	STRATIFIED UNITS	INTRUSIVE UNITS
YAKUTAT (YA)	KY	Eg, Oggb				EK	
CHUGACH (CG)	KV, KVm, KVgn, KVv, KVvm	Eg	Oggb				
WRANGELLIA (WI?)	PMm, PMsv	PPy, JKg, JKgr, JKgd, Eg					
WRANGELLIA (WI)	PPs, Pc, uRN, uRC, uRM	PPy?, JKg		Ks			
ALEXANDER (AX)	IPgn, IPvm, IPvs, IPcs, Psa, Pvb, Ec, Ec, Evs, Eoc, EOwb, EOls, EOw, EOwv, ODcs, OSwp, SDc, Dp, Dc, Psp, uPs, uPv, uPvs, uPc, uPpc, uPp, uPcs, uPcp, uPe	PPgd, PPgdi, PPydi, PPy, PPdi, PPg, JKg, JKgd, JKgr, Eg, Ogb, Oqd, Ogdp, Ogd, Ogr, Ogp, Og, Of, Mg, Mf	EOb uPbv?, PRub?, PRb?, Mb, KTb,	Ks		Pvs, Ps, Os, Nv, NW, NWI, NW <sub>2</sub> , NW <sub>3</sub> , NW <sub>4</sub> , NW <sub>L</sub> , NWpy, NWcg, NWm, NWu	Mf
WRANGELLIA (W2)	Pv, Pvx, Psv, Pc, Ps, PMvs, PRv, mRv, uRN, uRe, uRC, uRM, RKp?	Kg, Kgd, Kqd, Kdi, Kqm, Of, Mdi, Mf	uPb, uPbv?, PRub, PRb, KTb	JKD, RKp	Kg	Os, Nv, NW, NW <sub>2</sub> , NW <sub>L</sub> , NWpy, NWcg, NWm, NWu, Pt	Mf
WRANGELLIA (W3, TAKU)	PMvs			JKD	IKub, IKb, Kgd, Of,	Os	
"COAST BELT" (?TERRANES(S))	PMgn, Msm (in part), Mc	KTg		Msm (in part)	KTg	Pv	

Table 1.

Several unconformities, such as documented in southeastern Alaska (Gehrels and Saleeby, 1987), are suspected but unproven within the successions. Two distinct episodes of plutonism occurred in the Alexander Terrane of Saint Elias Mountains during Late Pennsylvanian to Early Permian (270-290 Ma) and Late Jurassic to earliest Cretaceous (130-160 Ma) time. Mafic plutonism also accompanied volcanism during the Cambrian-Ordovician, mid to late Paleozoic and Permo-Triassic (Dodds and Campbell, 1988) periods. Strata mostly are regionally metamorphosed to sub- or low-greenschist facies, but in the western parts the grade locally reaches upper greenschist to mid-amphibolite facies. The Alexander Terrane has been complexly deformed and rocks generally exhibit a pervasive cleavage or foliation. The dominant structural and metamorphic grain is northwest-southeast and along the eastern perimeter large northwest-trending, northeast-verging synclinoria and anticlinoria have been outlined. In the northern and central parts of the terrane, large folds, some of which are recumbent and southeast-verging, trend northeasterly. The widespread metamorphism and deformation seems to be post-Late Triassic and pre-earliest Cretaceous. Upright and overturned west-northwest-trending folds, locally superimposed on the major structures, are probably related to major, mostly Cenozoic strike-slip faulting. The Alexander Terrane is bounded by the Hubbard and Duke River faults; the former possibly a pre-mid-Pennsylvanian terrane boundary or suture of unknown displacement (Gardner et al., 1988), the latter a post-Triassic dextral transcurrent fault with large displacements offsetting terrane boundaries.

Segments of Wrangellia with their Mesozoic cover sequences (Berg et al., 1972; Monger and Berg, 1984) flank the Alexander Terrane on the west and southwest (W1) and north and northeast (W2, W3). W1 is confined by the Border Ranges and Hubbard faults and in Saint Elias Mountains only locally includes the weakly metamorphosed late Paleozoic and Triassic rocks typical of Wrangellia in the McCarthy area Alaska (MacKevett, 1978). In Canada, W1 largely comprises pervasively deformed argillite, schist, greenstone, amphibolite and marble, which have been interpreted as a deeper, more metamorphosed equivalent of Wrangellia (Plafker et al., 1989). W1 is intruded by syenitic dykes and small plutons (270-290 Ma) and large elongate batholithic complexes (130-160 Ma) (Dodds and Campbell, 1988). The Jurassic-Cretaceous cover rocks to W1 are shallow marine sediments (MacKevett, 1978), but these have only locally been recognized in Yukon. Wrangellia W2 is bounded by the Duke River and Denali faults. It consists of Pennsylvanian oceanic arc volcanics and sediments, Lower Permian shallow marine sediments, and mid to Upper Triassic basic rift volcanics and variable amounts of shallow marine evaporite, carbonate, and argillite (Muller, 1967; Read and Monger, 1976). The strata of W2 are generally in the sub-greenschist grade and have been deformed into upright to northeast verging folds. W2 is intruded by upper Paleozoic (?) and Upper Triassic mafic and ultramafics probably related to the mafic volcanism, and by large, elongate, syn- or post-tectonic, granitic plutons (106-117 Ma; Dodds and Campbell, 1988). In the narrow belt between the Duke River and Denali faults, W2 is cut by anastomosing networks of faults, which bound lenses of folded rocks with coherent stratigraphy in a fault melange. Wrangellia W3, which lies northeast of Denali Fault System along the western margin of the Coast Plutonic Complex, comprises little known pervasively deformed greenschist to amphibolite grade metasediments, metavolcanics and local mafic-ultramafic intrusions. In adjacent southeast Alaska W3 abuts the northern part of the Taku Terrane, which has been correlated with rocks of Wrangellia in the

eastern Alaska Range (Plafker and Hudson, 1980; Davis and Plafker, 1985; Plafker et al., 1989). The Jurassic-Cretaceous cover rocks to W2 and W3 consist of weakly metamorphosed and deformed deep marine flysch and local basic volcanics (Eisbacher, 1976; Richter, 1976). The cover is intruded by an Alaskan-type concentric ultramafic complex (Sturrock, 1975; Sturrock et al., 1980) and granitic plutons of Late Early Cretaceous age (Richter et al., 1975; Dodds and Campbell, 1988). Northeast trending folds of probable Cretaceous age in the cover rocks were refolded, presumably the result of dextral-strike-slip displacements along Denali Fault System which during latest Mesozoic to latest Cenozoic time amounted to some 300 km. (Eisbacher, 1976). Wrangellia W3 and its metamorphosed Jurassic-Cretaceous cover rocks are intruded along their east flank by the Coast Plutonic Complex (Kindle, 1953; Wheeler, 1963; Muller, 1967; Tempelman-Kluit, 1974; Eisbacher, 1976; Erdmer, 1989, 1990), which locally includes part of the "Tonalite Sills" (of Brew and Ford, 1978, 1981, 1985).

The Chugach Terrane is bounded on the northeast by the Border Ranges Fault System (MacKevett and Plafker, 1974; Plafker et al., 1976; Plafker and Campbell, 1979; Plafker et al., 1985) and on the south and southwest by Chugach-Saint Elias and Fairweather faults. It is an accretionary prism consisting of variably metamorphosed and deformed Cretaceous and older (?) deep-marine flysch and oceanic crustal rocks (Plafker et al., 1977b; Hudson and Plafker, 1982; Plafker et al., 1989). In the Seward Glacier area and adjacent Alaska the terrane is partitioned by the Columbus Fault into two north-dipping thrust sheets in which metamorphic grade and deformation increases downward. These comprise a northern high temperature-low pressure assemblage of subgreenschist to amphibolite-grade metasediments, granitoid gneisses and pegmatite, and a southern assemblage of low grade basic volcanics and foliated crystalline amphibolite. The metasedimentary and metavolcanic division of the terrane is also recognizable in the Fairweather Ranges (G. Plafker, pers. comm., 1979). The Chugach Terrane is intruded by Eocene (41-52 Ma) and Oligocene granitic plutons (Hudson et al., 1977; Hudson, 1983; Dodds and Campbell, 1988), and by mid Tertiary layered gabbros (Loney and Himmelberg, 1983). Accretion and deformation of the terrane commenced in the Early Jurassic and continued into the latest Cretaceous and/or earliest Tertiary (Plafker et al., 1989).

The Yakutat Terrane is mostly bounded on the north and northeast by the Chugach-St. Elias and Fairweather faults, and on southwest by the Transition Fault System. It is a composite terrane and comprises an eastern upper Mesozoic flysch and melange assemblage and a western sequence of Eocene oceanic crustal rocks, which are basement to an Eocene to Quaternary dominantly marine clastic cover (Plafker et al., 1989). The terrane is believed to have been transported to its current position from a site some 600 km to the southeast, by dextral strike slip along the Fairweather-Queen Charlotte transform fault system during Neogene to present time (Plafker, 1987). Only a very narrow sliver of the Yakutat terrane occurs in Saint Elias Mountains of Canada adjacent to Newton and Seward glaciers. There the terrane is represented by north-dipping thrust sheets of low grade metamorphosed and folded late Mesozoic flysch, melange, and minor early Tertiary clastics within the lower plate of the St. Elias fault. It is intruded by small granitic plutons of Eocene and Oligocene age (Hudson et al., 1977).

Tertiary nonmarine clastic sediments and volcanics unconformably overlie the deformed rocks of the Alexander Terrane and segments of Wrangellia. The sedimentary sequences consist of Eocene (?) and Oligocene mostly fluvial deposits with local landslide, debris flow, lacustrine and pull-apart basin deposits, and rarer volcanics (Eisbacher and Hopkins, 1977; Dodds, 1987; Ridgway et al., 1989). These clastic rocks were apparently deposited on a gently undulating low elevation surface, and once were distributed extensively over the eastern Saint Elias Mountains (Eisbacher and Hopkins, 1977). They now mostly remain only in small faulted, erosional remnants. Overlying and overlapping the sedimentary strata are locally very thick Miocene-Pliocene subaerial calc-alkaline volcanics of the Wrangell Lava, which occur in widely scattered deposits on the Alexander Terrane and segments of Wrangellia (Souther and Stanciu, 1975; Souther, 1977; MacKevett, 1978). Where mapped in detail, the Wrangell Lava has been subdivided into the Canyon Mountain, St. Clare and Alsek provinces (Souther and Stanciu, 1975). The clastic strata are only locally openly to tightly folded close to faults (Eisbacher and Hopkins, 1977; Dodds, 1988; Ridgway et al., 1989). The Wrangell Lava is mostly fairly flat lying or shallow dipping. However, in the northern Saint Elias Mountains they have been openly folded and locally overturned adjacent to a group of reverse faults (Muller, 1967; Souther and Stanciu, 1975). Oligocene (24-31 Ma) plutonism present in southeasternmost Saint Elias Mountains and adjoining southeastern Alaska (MacKevett et al., 1974; Brew et al., 1978; Jacobson, 1979; Jacobson et al., 1980; Brew and Morrell, 1983; Brew and Ford, 1985; Dodds and Campbell, 1988) may be coeval with volcanics similar to those in the O'Connor River area. This granitoid plutonism and also felsite, latite and crowded feldspar porphyry subvolcanic intrusion widespread in Kluane and Kusawak ranges are both probably strike-slip fault related. Miocene (6-16 Ma) basic to acidic subvolcanics and local deeper-level granitoid plutons (Downey, 1978; Downey et al., 1980; Dodds and Campbell, 1988) within or close by Wrangell Lava are considered comagmatic to the volcanics, and related to the northward oblique subduction of the Pacific Plate beneath the continent (MacKevett, 1978). Pliocene and younger (?) diamictite and fluvial clastics occur in the Bull Creek area and also along the White River in Alaska. They have been variously interpreted as subglacial tillite deposits (Denton and Armstrong, 1969), lahars derived from glaciated volcanic cones (Plafker et al., 1977c), or subaerial slope and lacustrine fan-delta deposits in a strike-slip setting (Eyles and Eyles, 1989). These clastics have been locally tilted and faulted. Latest Pliocene and Pleistocene (?) volcanics are speculated to be present within parts of the Wrangell Lava sequence and also in the Blanchard River area on the westernmost flanks of the Coast Belt. Large-scale strike-slip faults, such as the Denali (Eisbacher, 1976; Eisbacher and Hopkins, 1977; Clague, 1979a), Fairweather (Plafker et al., 1978; Carlson et al., 1985; and others) and Totschunda (Richter and Matson, 1971; Plafker et al., 1977a) faults, were active during mid- to latest Cenozoic time in Saint Elias Mountains and closely adjacent parts of Alaska. Dextral displacements along Denali Fault System may have been transferred to Totschunda Fault during latest Cenozoic time, by folds and reverse faults displayed in the Wrangell Lava of St. Clare Province. Linear ponds and mounds along sections of Denali Fault System suggest that some movements may be Pleistocene, but evidence for post latest Pleistocene strike-slip displacements is lacking (Clague, 1979a). However, Holocene dip-slip fault scarps occur (Eisbacher and Hopkins, 1977; Clague, 1979a) and a zone of microseismicity has been outlined (Boucher and Fitch, 1969; Horner, 1983) along the Denali-Totschunda fault linkage. Late Cenozoic faults and related deformation are suspected elsewhere in the Alexander Terrane of the region.

These include a major reverse fault (southwest side up) between Klutlan and Kaskawush glaciers which produced the Wood-Steele-Walsh mountain front, and strike-slip-fault related brittle deformation in O'Connor River and Tats Creek areas (Dodds, 1988). Dextral strike-slip displacements along Fairweather Fault coupled with underthrusting on St. Elias and adjacent faults during latest Cenozoic time resulted in the spectacular relief of the Mt. St. Elias-Mt. Augusta and Mt. Logan fronts. The rise of the Saint Elias Mountains was probably initiated during the Miocene with most of the tectonism concentrated in the Pliocene and Pleistocene, and has resulted largely from the oblique subduction of the Yakutat Terrane and the Pacific Plate.

Regional studies of surficial geology and geomorphology have been undertaken in the Saint Elias Mountains (Rampton, 1979a,b,c,d, 1981, 1982a,b,c). More detailed investigations have been made in the mountains of surge-type glaciers (Stanley, 1969; Clarke et al., 1986), ice damned lakes and related glacial outburst flood hazards (Clague, 1979b; Clarke and Mathews, 1981; Clarke, 1982; Clague and Rampton, 1982), and landslides (Clague, 1981). The White River ash eruptions of about 1200 and 1500 to 1800 years age close to the Alaska-Yukon boundary (Lerbekmo and Campbell, 1969; Hughes et al., 1972; Lerbekmo et al., 1975; MacKevett, 1978; and others), represent the last known major volcanic activity in the region. Large historical earthquakes have occurred in Saint Elias Mountains and immediately adjacent parts of Alaska, with several major events concentrated along the Fairweather and related faults. These include the 1899-1900 sequence of great earthquakes of the Yakutat Bay area estimated at magnitudes (M) of 8.5, 7.8, 8.4 and 8.1 during a thirteen month interval (Tarr and Martin, 1912; Thatcher and Plafker, 1977); the 1958 Lituya Bay earthquake magnitude (M) 7.9 (Stauder, 1960; Tocher, 1960; Tobin and Sykes, 1968; Plafker et al., 1978), which resulted in the landslide induced giant tidal wave at Lituya Bay (Miller, 1960); and the 1979 St. Elias earthquake (M) magnitude 7.5 (Lahr et al., 1979, 1980; Stephens et al., 1980; Stover et al., 1980), which triggered landslides in the Seward, Tweedsmuir, Towagh and Jarvis glaciers areas. Recent seismic monitoring (Horner, 1983, 1989; and others) attests to the continuing tectonic instability and uplift in the Saint Elias Mountains.

## Economic Geology

Following white man's first visit in 1891, the recorded history of mineral exploration and development in Saint Elias Mountains as elsewhere has been cyclical. Mineral occurrences discovered to date include a variety of deposit types, however many seem to be minor showings. Placers are probably all fluvial deposits. Lode types include assorted skarn, magmatic, volcanic red bed, volcanogenic massive sulphide, porphyry, and vein and/or replacement deposits. Nonmetallic sedimentary types include lignite coal, gypsum-anhydrite evaporite, and bentonite clay deposits. Until now placers (gold ± minor platinum, silver, copper) have been the dominant producers with Burwash Creek recording much of the yield. To date production from lode deposits amounts to relatively small tonnages of high-grade copper, lead, zinc, nickel (± precious metals), while no production has been recorded from nonmetallic sedimentary deposits. Currently placer gold operations in the Burwash Creek area and Squaw Creek account for all mineral production from the region. By far the most important known lode occurrences are Windy Craggy volcanogenic massive sulphide (Cu,Co,Au,Ag) and Wellgreen magmatic

(Ni,Cu,Co,PGE,Au) deposits. The more significant nonmetallic mineral prospects outlined are O'Connor (gypsum, anhydrite) and Datlasaka (bentonite) deposits. Only a few of the mineral deposits in Saint Elias Mountains have been visited by the authors, and tabulation by map sheet of information on them (Tables 2, 3, 4, and 5) is largely from literature surveys.

### **History**

Placer copper was known and utilized by local native Indians, prior to white man's arrival in 1891 (Hayes, 1892). During 1896-1900 pack-horse-access on the Dalton Trail opened up the area resulting in the first recorded discoveries of placer gold in Dezadeash map area (Brooks, 1900; Tyrrell, 1901), vein copper on Kletsan Creek (Brooks, 1900) and skarn copper at Rainy Hollow. In 1903 placer gold finds near Silver City, precipitated the initial discoveries of placers and lignite coal in Kluane Ranges between Donjek and Dezadeash rivers (McConnell, 1905). Between 1905-1908 the earliest staking took place of vein copper showings in White River and Burwash Creek areas (Cairnes, 1915a; Muller, 1967). From 1908 to 1911 other copper finds were made in the Rainy Hollow area and small tonnages of high-grade ore shipped. The Chisana Gold Rush in 1913-14 kindled placer prospecting in the White River area (Cairnes, 1915a). In 1915 lode gold was found at Gold Cord west of Pleasant Camp (Fraser, 1915). During 1920-22 development work was done at Maid of Erin with some shipment of high-grade ore (Watson, 1948). In 1927 coarse gold was discovered on Squaw (Dollis) Creek (Cockfield, 1928; Mandy, 1933, 1934). Except for small scale placer operations little mining activity took place in the area between 1930-44. Public access to Alaska Highway-Haines Road in 1944 and World War II technology, significantly enhanced placer production and lode exploration in the region. Fixed-wing and later helicopter supported mineral exploration and Government geological mapping in the 1950s led to discoveries of significant deposits of lode copper at Bornite Creek in 1950 (Kindle, 1953) and at Johobo in 1959 (Read and Monger, 1976), magmatic nickel-copper-platinum group elements at Wellgreen in 1952 and at Canalask (Micro) in 1953 (Muller, 1967; Read and Monger, 1976)), and massive copper sulphides at Windy Craggy in 1957-58 (MacIntyre, 1983). Exploration immediately following these finds, resulted in the location of many of the smaller lode copper and nickel showings now known in the eastern parts of the region. During 1959-61 some high-grade ore was shipped from Johobo. Between the 1960s and mid 1970s improved geophysical and geochemical survey techniques helped to better outline many of the known deposits and to target other bodies or areas for assessment. During this period development work was done on Wellgreen, Canalask, White River, Airways, Lynda, Cork, Glen, Dickson, Kloo, Jackpot and other deposits. Between 1968-1971 Kane, Lunar and Hum Bird lead-zinc-precious metal vein, and Lep, Taylor, Sanpete and Bornite skarn deposits among others, were staked and worked on. Around 1970 some evaluation was made of Niamodlaoc Mtn. and other lignite coal deposits (Cameron and Birmingham, 1970). In 1973 underground mining problems resulted in closure of Wellgreen Mine; stockpiled concentrates to that date amounted to 33 924 tonnes grading 7.4% nickel and 6.6% copper. In 1974 authorization of Kluane National Park reserve resulted in some 22 015 km<sup>2</sup> (8500 miles<sup>2</sup>) of Saint Elias Mountains in Yukon Territory being set aside as a National Heritage Site, and restriction of mining activities to areas outside park-limits. Mid- to late 1970s exploration in Alsek Ranges by Swiss Aluminum Mining Company of Canada led to discoveries of small massive sulphide deposits

including Mus (X) and a subglacial magnetic anomaly near Windy Craggy, and to Alu skarn deposit above Tarr Inlet. Between 1980-83 an intensive drilling, sampling and airborne geophysical program was undertaken at Windy Craggy by Falconbridge Ltd. and Geddes Resources Ltd., which outlined for the first time the potential size of the massive sulphide deposits (MacIntyre, 1983, 1984; Gammon and Chandler, 1986). In 1983 St. Joe Canada Inc. acquired lapsed Swiss Aluminium claims in East Arm glacier area. They revealed promising precious metal values in volcanogenic copper-zinc-cobalt mineralization at Rime showings (MacIntyre, 1984), substantiated the subglacial anomaly and postulated a hidden subglacial source for abundant massive sulphide glacial erratics in Henshi Creek valley (Day, 1985; Day et al., 1987). In 1987 on-ice drilling of the anomaly intersected massive sulphides just beneath the glacier. Mid-1980s exploration by Stryker-Freeport Resources Ltd. in Tsirku-Jarvis glaciers area resulted in discoveries of volcanogenic copper ± barite deposits and to significant float boulderados on the flanks of Mount Henry Clay (Still, 1984; MacIntyre and Schroeter, 1985). Also during that time regional programs by Noranda Exploration Co. Ltd. led to among others small finds of copper skarns and copper-precious metal veins at Red Mountain, and in the upper Klehini and Kelsall rivers areas. Between 1984-87 Haines Gypsum Inc. and Queenstake Resources Ltd. did extensive work on O'Connor gypsum deposits (White, 1986). During 1986-88 Kluane Joint Venture (All-North Resources Ltd. and Chevron Minerals Ltd.) optioned and reassessed Wellgreen and Canalask properties particularly for platinum group elements (PGE). Kluane Joint Venture restaked other known magmatic nickel-copper prospects including Onion, Cats and Dogs, Airways, Lynda, Duke South, Frohberg, and C and E North, and under option agreements with a variety of companies, also explored them for PGE potential (INAC, 1988; R. Carne, pers. comm.). During 1985 to late 1989 Geddes Resources Ltd., operating from an all-weather camp and a DC3 airstrip in Tats valley flats, undertook major underground exploration and first phase mine feasibility studies at Windy Craggy. Early in 1987 work began on a 1852-meter adit system accessible from Tats Camp by a road constructed on Tats Glacier. Intensive year-round work was carried out from 1987 to late 1989, by which time the adit system and an extensive underground drilling and sampling program had been completed. By early 1990 two major ore bodies had been outlined and Geddes Resources Ltd., now with Northgate Ltd. and Cominco Ltd. as large shareholders, released ore reserve estimates of 165 million tonnes grading 1.9% copper, 0.08% cobalt, 0.2 g/tonne gold and 3.9 g/tonne silver (Geddes Res. Ltd., 1990).

### ***Placer deposits***

Auriferous placer deposits in Saint Elias Mountains of Canada have been worked since about 1898 (Brooks, 1900; Tyrrell, 1901; Kindle, 1953). They are probably all fluvially derived and have accounted for almost the entire metal production from the region. Most are in unconsolidated Quaternary sediments, but one Tertiary paleoplacer has been reported (Muller, 1967). They occur mainly along the eastern parts of the region in Nutzotin, Donjek, Kluane, northern Alsek and Squaw ranges. Areas placers are known from and have been worked include: upper White River; upper Koidern River; tributaries of Donjek River; Reed Creek and closeby creeks; Burwash and adjacent creeks; Bullion and neighbouring creeks; Jarvis River and tributaries; many creeks throughout SW Dezadeash map sheet; and Squaw and nearby creeks. Gold is recorded from all past or present producers and most showings. Native copper has been obtained from some

operations, platinum and/or silver from a few deposits, and tin has been reported from at least one creek.

Most of the placer deposits are located on or nearby the complexly deformed and intruded, relatively narrow belt of Wrangellia (segments W2, W3) volcanic and sedimentary rocks and their Jura-Cretaceous flyschoid and Tertiary volcanic and nonmarine clastic covers. These bedrock sequences have been intermittently and extensively intruded by a spectrum of Mesozoic and Tertiary ultramafic, mafic, intermediate, and acid plutons and subvolcanic intrusions. The belt is crudely contained and deformed by two major strike-slip faults; the Duke River Fault to the southwest and the Denali and possibly related faults abutting the Coast Belt to the northeast. Combined large-scale displacements along these faults have produced the present, fairly narrow, complex melange zone of moderately to highly deformed and altered diverse rocks types intricately bounded by anastomosing fault networks and gouges. An assortment of lode deposit types also occur throughout this fault melange belt, and most carry copper ± precious metals. These include: Late Triassic mafic-ultramafic magmatic nickel-copper-platinum ± gold; Late Triassic red bed copper ± native copper and silver; Late Triassic skarn ± precious metal; Late Early Cretaceous Alaskan concentric ultramafic chrome, platinum; Late Triassic and/or older volcanogenic copper ± precious metal; Mesozoic and Tertiary age fault- or fold-related quartz ± carbonate vein and/or replacement ± copper, gold, silver; and Oligocene and Miocene intermediate to acid subvolcanic porphyry copper, molybdenum ± gold, silver deposits. Erosion, transportation and sorting of these diverse rock types and lode deposits, and possibly also reworking of Tertiary fluvial and ubiquitous Pleistocene surficial clastics, by juvenile pre, syn and post glacial river systems, have almost certainly contributed metals to the placer deposits of the belt.

Operations undertaken on Burwash, Bullion, Squaw (Dollis), Shorty, Squirrel creeks and Reed Creek tributary account for most of the recorded placer gold, platinum, silver and native copper production from the region to date. Smaller-scale workings on Sugden, Sheep, Arch, Kimberley, Quill, Tatamagouche, Swede Johnson, Porcupine and Beloud creeks yielded much of the remaining reported production. Most of the recently past to current production and activity has come from Burwash, Squirrel, Reed (tributary), Kimberley, Swede Johnson, Porcupine, and Squaw creeks and Donjek River. The majority of other placer occurrences listed in the report, constitute minor or "panned-colours" only showings. The following are some incomplete recorded yields (in crude oz.) of gold from the more significant placer producing creeks in Saint Elias Mountains: Burwash Creek (24 322 oz., 1904-1970; 2013 oz., 1978-88); Bullion Creek (9487 oz., 1904-70); Squaw (Dollis) Creek (5077 oz., 1927-70); Shorty Creek (1863 oz., 1945-46; 2436 oz., 1947-59); Squirrel Creek (1279 oz., 1978-88); Reed Creek tributary (1082 oz., 1978-88); Sugden Creek (682 oz., 1947-59); Sheep Creek (676 oz., 1904-70); Arch Creek (540 oz., 1904-1973; 102 oz., 1978-88); Kimberley Creek (560 oz., 1978-88); Quill Creek (108 oz., 1978-88); Tatamagouche Creek (100 oz., 1904-70); Swede Johnson Creek (65 oz., 1978-88); and Porcupine Creek (35 oz., 1978-88). Placer production figures quoted are from the following sources: 1904-1973 and 1927-70 (Read and Monger, 1976); 1945-46 (Kindle, 1953); 1947-59 (Debicki, 1982); and 1978-88 (LeBarge and Morison, 1990).

## *Lode deposits*

A fairly wide variety of lode deposit types occur in Saint Elias Mountains of Canada. However, many appear to be relatively minor showings. They include (classifications mostly after Eckstrand, 1984): lead-zinc-silver, tungsten, and iron skarn ( $\text{Cu} \pm \text{Au}, \text{Bi}, \text{Mo}$ ); magmatic (mafic/ultramafic-hosted) nickel, copper ( $\pm \text{PGE}, \text{Au}, \text{Zn}$ ); magmatic (mafic/ultramafic-hosted) chromite (+ Pt, Fe, Ti); volcanic red bed copper ( $\pm$  native Cu, Ag); volcanogenic massive sulphide ( $\text{Cu} \pm \text{Co}, \text{Zn}, \text{Au}, \text{Ag}, \text{Ba}$ ); porphyry copper, molybdenum ( $\pm \text{Au}, \text{Ag}$ ); vein and/or replacement  $\pm$  ( $\text{Cu}, \text{Au}, \text{Ag}$ ) and lead-zinc ( $\pm \text{Cu}, \text{Au}, \text{Ag}$ ); and ultramafic-hosted asbestos deposits. Classifications of many of the deposits are crude or tentative due to varying degrees of alteration (hydrothermal and metamorphism) and deformation, and to lack of detailed studies. Most of the deposits are located in the less rugged and ice-covered more easily accessible eastern parts of the region. Production to date amounts to only small tonnages of high-grade copper, lead, zinc, and nickel ( $\pm$  precious metals). The most significant lode occurrences in the area are Windy Craggy massive sulphide ( $\text{Cu}, \text{Co}, \text{Au}, \text{Ag}$ ), Wellgreen magmatic ( $\text{Ni}, \text{Cu}, \text{Co}, \text{PGE}, \text{Au}$ ) and Canalask (Micro) magmatic ( $\text{Ni}, \text{Cu}, \text{Co}, \text{Zn}, \text{PGE}$ ) deposits, all of Triassic age.

An assortment of lead-zinc-silver, tungsten, and iron (magnetite) type skarns ( $\text{Cu} \pm \text{Au}, \text{Bi}, \text{Mo}$ ) are undoubtedly present in the region. However, subdivision of most occurrences is not possible due to alteration, deformation and insufficient data. The majority of known copper skarn deposits are in the Rainy Hollow or adjoining areas, but there are a few others adjacent to White River. A good quality wollastonite skarn is also reported from the White River area (Archer, Cathro and Associates Ltd., pers. comm., 1990). Many of the deposits could be hosted by carbonates of Late Triassic age, but Alu copper-molybdenum-tungsten skarn deposit among others is in older rocks. Almost all occurrences seem to be minor, however some in the Rainy Hollow area have produced small tonnages of high-grade ore. Notable among these is Maid of Erin, which between 1911 and 1922 shipped a total of 157 tons of sorted ore to Tacoma yielding 77 658 lb. of copper, 5849 oz. of silver, and 6 oz. of gold (Watson, 1948).

Magmatic (mafic/ultramafic-hosted) nickel-copper ( $\pm \text{PGE}, \text{Au}, \text{Zn}$ ) deposits are present along the easternmost part of the region. They occur in Triassic and possibly older mafic-ultramafic intrusions and contact host rocks, exclusively within Wrangellia segments (W2, W3). Many of the deposits have undergone some degree of deformation and alteration, due to late Mesozoic and Cenozoic folding and faulting. The most significant occurrences of this type are Wellgreen ( $\text{Ni}, \text{Cu}, \text{Co}, \text{PGE}, \text{Au}$ ) and Canalask (Micro) ( $\text{Ni}, \text{Cu}, \text{Co}, \text{Zn}, \text{PGE}$ ) deposits. Since its discovery in 1952, considerable development work has been done on the Wellgreen property (Campbell, 1956, 1960; Muller, 1967; Campbell, 1981; Carriere et al., 1981; Hulbert et al., 1988; INAC, 1989). Hudson Yukon Mining Co. Ltd. produced 33 924 tonnes of concentrates grading 7.4% nickel and 6.6% copper, but underground problems resulted in mine closure in 1973. Intermittent exploration work has also been carried out on Canalask following its discovery in 1953, but no ore has been produced. During 1986-88 Kluane Joint Venture (All-North Resources Ltd. and Chevron Mineral Ltd.) optioned and reassessed Wellgreen and Canalask properties particularly for PGE content. In 1989 All-North Resources Ltd. released estimated ore reserves at Wellgreen of 50 million tonnes grading 0.36% nickel, 0.35% copper, 0.51 g/tonne platinum, 0.34 g/tonne Pd with significant values of other PGE, cobalt and gold. In 1990 All-North Resources

Ltd. published a potential ore reserve at Canalask (Micro) of 1.8 million tonnes grading 0.86% nickel in a footwall replacement deposit and also reported widespread disseminated nickel-copper-PGE mineralization in a nearby mafic-ultramafic sill. Other apparently much smaller deposits of this type occurring mostly in Kluane Range include Onion, Cats and Dogs, Airways, Lynda, Wash, Glen (Tatamagouche), Destruction, Duke South, Dickson, Frohberg, and C and E North. Many of these prospects were also initially staked in the 1950s and developed for nickel-copper. In 1986-88 Kluane Joint Venture and other companies restaked most of these and other prospects and evaluated them mainly for PGE content, but there has been no production from any of them to date.

A single small magmatic (mafic/ultramafic-hosted) chromite (+ Pt,Fe,Ti) occurrence (Stride) is reported (Kindle, 1953) from the Late Early Cretaceous Pyroxenite Creek Alaskan-type concentric ultramafic complex intruding Jura-Cretaceous flysch in Kluane Ranges just west of Haines Junction (Sturrock, 1975; Sturrock et al., 1980). Pan tests of gravels on one of the creeks dissecting the intrusion revealed fine platinum (Kindle, 1953). Little work has been recorded from the Stride prospect, which now is in Kluane National Park.

Volcanic red bed copper ( $\pm$  native Cu,Ag) deposits are present in various parts of Wrangellia W2 of the region. They all appear to be hosted by subaerial rift volcanics of the Upper Triassic Nikolai Greenstone, and most seem to be fairly minor showings. The better documented examples occur in the White River area at Kletsan (K-CU) (Hayes, 1892; Brooks, 1900; Moffit and Knopf, 1910; Cairnes, 1915a) and at Discovery (White River) (Cairnes, 1915a; Muller, 1967; Sinclair et al., 1979; Carriere et al., 1981; Eckstrand, 1984). Native copper occurring as placers near these deposits was used and traded by local native Indians before the arrival of white man. Several very large native copper nuggets including one weighing 1175 kg (2590 lb) were found close to the adit at Discovery (Cairnes, 1915a; Muller, 1967). Other small deposits of this type also occur in the Tatamagouche Creek area (Jacquot, Mary and Teddy, Quill (Ram), Quill Creek (Fossil and Hudson Bay), Burwash, and possibly Maple Creek (Vug)), and in the Mush Lake area (Husky, Wren, and Kel). Native copper found in placers close by, almost certainly were derived from deposits such as these. The Johobo (Yukon Star) and Bornite Creek (Kindle, 1953; Warnock, 1962; Read and Monger, 1976; Carriere et al., 1981) deposits, which are both hosted by Nikolai Greenstone, are most probably originally of this type. Deformation arising from movements along the nearby Denali Fault, however has probably resulted in remobilization and redeposition of the copper sulphides in gash-vein fills. Hand-sorted ore shipped from the deposits at Johobo included 2585 tons grading 23% copper and 2 oz./ton silver to Japan in 1959-61, and 1062 tons averaging 20.2% copper and 1 oz./ton silver to Tacoma in 1962. About 150 tons of ore was high-graded from Bornite Creek. These deposits, however, now lie within Kluane National Park. Other small volcanic deposits of uncertain genesis also occur in older strata of Wrangellia in the region. These include Pickhandle (M), Sandy (Cave), Shaft, Snow (Mush) and Jackpot (Pirate Creek) (Read and Monger, 1976; Carriere et al., 1981; INAC, (1989)), which are hosted by oceanic arc volcanics of the mostly Pennsylvanian Station Creek Formation in Wrangellia W2. Minor deposits of questionable volcanic origin also occur in Wrangellia W3, and include Kloo (Ellen) (INAC Overview, (1990)) and possibly Kelsal 32 (BCMEMPR Minfile, 114P (1988)) prospects.

Other volcanogenic ( $\text{Cu} \pm \text{Co}, \text{Zn}, \text{Au}, \text{Ag}, \text{Ba}$ ) deposits are also present in the Alexander Terrane of Saint Elias Mountains of Canada. Most of these occurrences are in Tatshenshini River map area (BCMEMPR Minfile, 114P (1988); NBCMI, 114P (1989)). The most significant by far is the Windy Craggy massive sulphide ( $\text{Cu}, \text{Co}, \text{Au}, \text{Ag}$ ) deposit (MacIntyre, 1983, 1984, 1986; Gammon and Chandler, 1986; Peter, 1989; Geddes Res. Ltd., 1990). It is hosted by a sequence of low greenschist grade, folded and faulted, Upper Triassic (Early Norian) submarine basaltic to andesitic arc volcanics and fine calcareous clastics, and is of exhalative type. Considerable underground development work by Geddes Resources Ltd. has revealed at least two major deposits containing ore reserve estimates of 165 million tonnes grading 1.9% copper, 0.08% cobalt, 0.2 g/tonne gold and 3.9 g/tonne silver with 69% of this amount classified as proven and probable and the remainder as being possible (Geddes Res. Ltd., 1990). First phase mine feasibility studies have been undertaken at Windy Craggy and are under review, but as yet there has been no production. On-ice drilling in 1987 of a sizeable magnetic anomaly beneath East Arm Glacier by St. Joe Canada Inc. in joint venture with Newmont Mines Ltd., revealed massive sulphide mineralization and the potential of another significant probably Upper Triassic exhalative copper deposit just beneath the glacier. Massive sulphide boulders are plentiful at the toe of East Arm Glacier and in Hensi Creek valley (Day, 1985; Day et al., 1987). Other apparently minor Upper Triassic volcanogenic deposits of probable exhalative type in the Windy Craggy area include among others Tats, Rime (X, Mus), Alsek (Tats Glacier), Barbican Mount, Aeolian Steeple and Pampero Ridge (MacIntyre, 1983, 1984; BCMEMPR, Minfile, 114P (1988)). Volcanic-hosted polymetallic sulphide  $\pm$  barite deposits occur adjacent to Tsirku-Jarvis glaciers in Tatshenshini River map area (MacIntyre and Schroeter, 1985; Schroeter and MacIntyre, 1986; MacIntyre, 1986; BCMEMPR Minfile, 114P (1988)). These deposits include Low Herbert, Low Jarvis (Jumar), High Herbert North, and Herbert (Mouth) East and West and were explored by Stryker-Freeport Resources Ltd. through the mid-1980s. Significant massive sulphide float boulderados were also discovered on the flanks of Mount Henry Clay on both sides of the international border. Most of the deposits appear to be relatively minor, and are of uncertain age and origin due largely to limited study. Although Upper Triassic occurrences are strongly suspected, all are not necessarily of that age. Some of the mineralization examined by Noranda Exploration Co. Ltd. at Red Mountain (Fair) may be in Upper Triassic volcanic and sedimentary strata (Schroeter and MacIntyre, 1986). Mineralized gossans at Valia and Mount Bigger, however, are hosted by volcanics of possibly Paleozoic age. A minor copper showing of uncertain origin is also reported in Upper Cambrian volcanics at Fenton (INAC, (1989)) adjacent to Bates River.

Porphyry copper, molybdenum ( $\pm \text{Au}, \text{Ag}$ ) deposits are locally present in Saint Elias Mountains both in Wrangellia and the Alexander Terrane. Most appear to be related to Oligocene or Miocene subvolcanic intrusions, and almost all seem to be minor showings. Development work has been done at Cork prospect on Burwash Creek found in the mid-1960s, and which occurs in Oligocene latite porphyry subvolcanic intrusions hosted by Lower Permian sediments of Hasen Creek Formation (Christopher et al., 1972; Read and Monger, 1976; Carriere et al., 1981). Some mineralization at Glen (Burwash) on Burwash Creek (Christopher et al., 1972; Read and Monger, 1976; INAC, (1987)) may also be of porphyry type and related to Oligocene latite intrusions in mafic-ultramafic and volcanic rocks of Wrangellia W2. Small deposits possibly related to Oligocene intrusions

elsewhere in Wrangellia W2 include Garlic, Tatshenshini, Dalton and some of the mineralization at Mohawk (Kane). Minor copper-gold bearing quartz stockwork and veins are also reported at Canyon Mountain (Can) in Wrangell Lava overlying Wrangell W2. Very minor insitu and float porphyry copper-molybdenum occurrences are recorded from the Alexander Terrane. These include Mineral Ridge (Sharp) and float in Steele Glacier area found by the Wood Yukon Expedition in 1941 (Sharp, 1943), and also St. Elias, Galloping and other mineralized float boulders on moraines of Steele, Spring and Donjek glaciers located by others (Muller, 1967). Elsewhere in Alexander Terrane minor copper-molybdenum mineralization with precious metal values is reported from a breccia pipe in Wrangell Lava of the Alsek Province associated with a Miocene acid subvolcanic intrusion (Souther and Stancui, 1975). All the above copper-molybdenum occurrences in the Alexander Terrane are now within Kluane National Park.

An assortment of quartz ± carbonate vein and/or replacement deposits with ± copper, gold, or silver, and lead-zinc ± copper, gold or silver are present in Wrangellia and the Alexander Terrane of the region. Many are likely fault- or fold-related, and almost all seem to be minor occurrences. In Wrangellia segments W2, W3 veins ± Cu,Au,Ag occur at Rabbit Creek (CC), Kelli (Reed Creek), Colton 1, and Watson; and veins with Pb-Zn ± Cu,Au,Ag are present at Liberty, Jackpot (Pirate Creek), Mohawk (Kane), Mount Mansfield (Bear), and Kel. Of these more exploration work has been carried out on Jackpot (Pirate Creek) (Read and Monger, 1976; Carriere et al., 1981; INAC, (1985)) and Mohawk (Kane) (Read and Monger, 1976; INAC, (1986)) prospects. In the Alexander Terrane veins ± Cu,Au,Ag occur at Jennifer 2 (Kincora), Burger King, Kud, Saddle (3/4), Klehini River (KR1, KR4, KR7), Gold Cord (Stampede), and Grizzly Heights; and veins with Pb-Zn ± Cu,Au,Ag are present at Bates (Iron Creek), Hum Bird (Discovery, Creek, South, Camp, Dome), Fault Creek, and Parton. Of these more development work has been done on Gold Cord (Stampede) (Watson, 1948; Schroeter and MacIntyre, 1986; BCMEMPR, 114P (1988)), Grizzly Heights (MacIntyre and Schroeter, 1985; BCMEMPR, 114P (1988)), and HUM BIRD (Schroeter, 1978; BCMEMPR, 114P (1988)) prospects.

Ultramafic-hosted asbestos is present in various parts of the region, but all occurrences are very minor and of low quality. Veinlets of asbestos are reported in Late Triassic ultramafics in Wrangellia W2 at Duke and Ptarmigan-Windgap, and in ultramafic rocks of unknown age in Wrangellia W3 at Decoeli. Picrolite asbestos veinlets up to one inch wide are also recorded from peridotite of unknown age in the Alexander Terrane, both on an island in Bates Lake (Kindle, 1953) and at Nadahini Mountain (Watson, 1948).

### ***Nonmetallic Sedimentary Deposits***

Sedimentary deposits of nonmetallic type in Saint Elias Mountains of Canada include lignite coal, gypsum and anhydrite, and bentonite clay. Many of the deposits are small or uneconomic, and to date there has been no recorded production from any of them. The most development work done has been on the O'Connor gypsum deposit.

Low-grade coal occurs exclusively in Tertiary sedimentary rocks in widely scattered parts of the region. Seams up to three metres thick have been reported from the Oligocene and older Amphitheatre Formation. However, the coal is mostly lignite and where tested has been ranked sub-bituminous C or lower grade and appears uneconomic

(Cameron and Birmingham, 1970). Notable seams of lignite coal have been recorded from the following areas: north of White River at McLellan (Coal Creek) and Memoir (Cairnes, 1915a); at Cement Creek, and adjacent to Duke River at Amphitheatre, Granite Creek (Hoge), and Niamodloac Mtn. (Muller, 1967; Cameron and Birmingham, 1970; Read and Monger, 1976); and nearby Slims River at Stove (1,2), and at Kimberley Creek (McConnell, 1905; Muller, 1967; Cameron and Birmingham, 1970; Read and Monger, 1976). Very minor occurrences of lignite coal have been also documented in Tatshenshini River map area on Coal Creek, Squaw Creek, Talbot Creek, in the Kusawak Range, and in several localities on Tats Creek. Thin beds of very low-grade coal and coaly siltstone also occur locally within the Wrangell Lava of St. Clare Province (Souther, 1974; Souther and Stancui, 1975).

Gypsum-anhydrite deposits of highly varying sizes are present mainly in the eastern parts of Saint Elias Mountains. They are more numerous in Wrangellia W2, but also occur in the Alexander Terrane. Most have undergone some degree of alteration and remobilization. Within Wrangellia W2 gypsum-anhydrite deposits of evaporite (possibly sabkha) type are locally known in the basal part of the Upper Triassic Chitistone Limestone. Although this relationship is not often seen, probably due to fault- and fold-related remobilization and redeposition, most deposits occur in close proximity to volcanic and carbonate strata of Late Triassic age and are believed to be of that age and of evaporite type. The largest gypsum and anhydrite deposits in Wrangellia W2 in the region occur adjacent to Bullion Creek in Kluane National Park. Variably sized smaller deposits are also present elsewhere in Wrangellia W2 in the following localities: on Wade and Burwash creeks; adjacent to Bock's Brook; along the east flanks of Kluane Ranges between Slims and Jarvis rivers; and reportedly in the lower parts of Nadahini Creek. Several gypsum-anhydrite deposits are known within the Alexander Terrane, and most of these are probably of evaporite type. They are of uncertain age, but some may be Late Triassic. The most significant of these are the O'Connor River (Snow) deposits.

Exploration work done on the O'Connor River deposits during 1984-1987 by Haines Gypsum Inc. and Queenstake Resources, outlined reserves of 8.85 million tonnes of gypsum with up to 8.0% anhydrite (White, 1986; BCMEMPR Minfile, 114P (1988)). Other smaller deposits occur in Kluane Ranges between Slims and Jarvis rivers, in the northern part of Squaw Range, and in Alsek Ranges just northeast of the Windy Craggy deposit. There has been no recorded production from any of the gypsum-anhydrite occurrences.

A bentonite clay deposit is reported from Datlasaka Range in Tatshenshini River map area. The setting of the Datlasaka bentonite deposit is not well known, but it is at least 15 m thick and estimated at 135 000 cubic metres (NBCMI, 114P (1988)).

# STRATIFIED ROCK MAP UNIT DESCRIPTIONS

## QUATERNARY

### PLEISTOCENE AND RECENT

**Qs**

undivided surficial deposits: includes glacial deposits, alluvium, colluvium; local landslide, lacustrine, eolian and ash deposits.

## TERTIARY AND (?) YOUNGER

### NEOGENE AND (?) YOUNGER PLIOCENE AND (?) PLEISTOCENE

**Pv**

"BLANCHARD RIVER VOLCANICS": reddish-brown basaltic flows; minor interbedded basaltic tuff, agglomerate, and conglomerate (predominantly basaltic clasts); local basaltic neck: (114P/15,16).

**Pt**

"BULL CREEK DIAMICTITE": diamictite, fluvial gravel, and sand: may locally include uppermost Miocene: ? equivalent to White River ("tillite") diamictite: (115F/8,9).

## MIOCENE TO PLIOCENE AND (?) YOUNGER

### WRANGELL LAVA: (Nw<sub>1</sub> to Nw inclusive)

(subaerial, calc-alkaline basaltic andesite and andesitic lava and pyroclastics; local rhyolite, dacite, olivine basalt and nonmarine volcaniclastics: arc volcanics: 300 to 2000 m).

**Nw**

undivided Wrangell Lava: rusty red-brown basaltic to andesitic flows; minor creamy-white to yellow acid pyroclastics and local acid flows: may locally include Qs and/or intra-Wrangell sediments: (115G/2-6; 115F/1,7,10,15; 115B/8-10,15,16; 115A/2,4-6,12).

### ST. CLARE PROVINCE: (Nw<sub>L</sub> to Nw<sub>U</sub> inclusive)

**Nw<sub>U</sub>**

basalt, trachybasalt, andesite flows and pyroclastic rocks; minor volcanic conglomerate: (115G/5; 115F/8,10).

**Nw<sub>m</sub>**

porphyritic and nonporphyritic basaltic andesite flows (minor pillow lava), interbedded with abundant felsic ash flows, acid tuff, coaly tuff, volcanic sandstone and conglomerate: acid pyroclastics may be related to intra-Wrangell (IMf) intrusions: (115G/5,12; 115F/7-10).

**Nw<sub>og</sub>**

volcanic conglomerate - intra-Wrangell; clasts all Wrangell derived: (115G/5,12; 115F/7-9).

**Nw<sub>py</sub>** acid tuff or ash flows - intra-Wrangell: (?) related to IMf subvolcanic intrusions: (115G/5; 115F/8).

**Nw<sub>L</sub>** thick, blocky, mainly nonporphyritic basaltic andesite flows, locally interbedded with white or light-grey clay and coaly siltstone; overlain by thin, closely stacked basaltic andesite flows: (115G/5,12; 115F/7-10).

**ALSEK PROVINCE: (Nw<sub>1</sub> to Nw<sub>4</sub> inclusive)**

**Nw<sub>4</sub>** thin, uniform basaltic andesite and andesite flows: (115B/9; 115A/5,12).

**Nw<sub>3</sub>** felsic pyroclastics; locally includes acid breccia, ash flows and air fall pumice; possibly related to porphyritic rhyolite intrusion (IMf) adjacent to Dusty River: (115B/9; 115A/5,12).

**Nw<sub>2</sub>** basalt and basaltic andesite flows and pyroclastics: related to central conduit (breccia pipe): (115B/8,9; 115A/5,12).

**Nw<sub>1</sub>** locally includes rhyolite and/or dacite lava, thin to thick ash flows and pumice: (115B/8,9; 115A/5,12).

## **TERTIARY**

### **NEOGENE AND (?) OLDER**

#### **MIocene TO PLIOCENE AND (?) OLDER**

**Nv** basaltic flows; minor rhyolitic flows and tuffs: probably equivalent to Nw, but may locally include Pvs: (114P/11,15).

## **PALEOGENE**

### **OLIGOCENE AND, OLDER AND (?) YOUNGER**

**Os** **AMPHITHEATRE FORMATION:** yellow-buff to grey-buff sandstone, pebbly sandstone, polymictic conglomerate, siltstone, mudstone; minor brown-grey carbonaceous shale and thin lignitic coal beds: (nonmarine clastics; predominantly fluvial deposits; local landslide, debris-flow, and lacustrine deposits: fault-trough clastic wedge: 0 to 1100 m): may locally include older, or intra-Wrangell: (115G/2,3,5,6; 115F/15; 115B/16; 115A/2-6,12; 114P/15).

### (?) EOCENE AND OLIGOCENE

- |     |   |
|-----|---|
| Ps  | sandstone, pebble conglomerate, polymictic conglomerate; minor carbonaceous shale; rarer thin coaly beds: may in part be equivalent to Os: (114P/6,9-11,14).  |
| Pvs | "O'CONNOR RIVER VOLCANICS": volcanic breccia, tuff; locally interbedded predominantly basaltic and andesitic flows: C. meta. locally by Og: (114P/2,7,10,11). |

### EOCENE AND PALEOCENE

- |    |  |
|----|--|
| Ek | <u>KULTHIETH FORMATION</u> : light grey partly arkosic sandstone, interbedded with dark grey siltstone, pebble conglomerate and thin coal beds: (shallow marine and nonmarine delta-beach bar deposits): (115C/7). |
|----|--|

## CRETACEOUS

### LOWER CRETACEOUS AND (?) YOUNGER

- |    |   |
|----|---|
| Ks | sandstone, conglomerate, shale, siliceous tuff: (shallow-marine shelf deposits): (115C/15). |
|----|---|

### CRETACEOUS AND (?) OLDER

#### YAKUTAT GROUP:

(deep marine; flysch, melange; unseparated on map: accretionary prism: 115C/1,2,7,8).

- |    |   |
|----|---|
| Ky | <p><u>"melange facies"</u>: melange blocks (to several kilometres) of greenstone, oolitic limestone, marble, granitic plutonic rocks, chert, greywacke, set in pervasively sheared matrix of black cherty and tuffaceous pelite: may be U. Jurassic to L. Cretaceous: R. meta. commonly to (2b), locally to (4).</p> <p><u>"flysch facies"</u> (local): grey to brown lithofeldspathic greywacke and conglomerate, in thick channel deposits or interbedded with grey to black siltstone, argillite, or slate: may be U. Cretaceous: R. meta. (1) to (2).</p> |
|----|---|

#### VALDEZ GROUP: (Kv<sub>vm</sub> to Kv inclusive)

(deep-marine flysch, submarine basic volcanics, and metamorphic equivalents: accretionary prism: 115C/7-10; 115B/5; 114O/8; 114P/3-6).

- |    |  |
|----|--|
| Kv | dark grey argillite, greywacke: may locally include Ks: R. meta. (2) to low (3): (115B/5; 115C/9). |
|----|--|

Kv <sub>m</sub>	metamorphic equivalents of Kv: includes granitic dykes, sills, and locally plugs grades to brown schists and granitoid gneisses (Kv <sub>gn</sub> ): in 114O,P includes Kv and (?) Kv <sub>gn</sub> : R. meta. high (3) or low (4).
Kv <sub>gn</sub>	brown granitoid gneisses, minor dark brownish-grey schists; includes granitoid dykes, sills, and locally plugs.
Kv <sub>v</sub>	green pillow lava, breccia, tuff: R. meta. (1) to (2): (115C/7).
Kv <sub>vm</sub>	very dark green amphibolite: (?) metamorphic equivalent of Kv <sub>v</sub> : R. meta. to high (3) or (4).

## JURASSIC AND CRETACEOUS

### UPPER JURASSIC TO LOWER CRETACEOUS

JKD	<u>DEZADEASH GROUP</u> : interbedded light to dark buff-grey lithic greywacke, sandstone, siltstone, thin dark grey shale, argillite and conglomerate; mass-flow conglomerate common in middle part; rare light grey tuff: (flysch; deep-marine fan deposits: ~3000 m): R. meta. (1) to (2a): (115F/15; 115B/16; 115A/2,3,5,6,11-13; 114P/15).
-----	--

## TRIASSIC TO CRETACEOUS

### UPPER TRIASSIC TO LOWER CRETACEOUS

TRKp	dark grey phyllite; minor greywacke and conglomerate: includes JKD and upper noncalcareous member of McCarthy Formation; may locally include uRM: (115G/2,3,5-7; 115B/15,16).
------	---

## TRIASSIC

### UPPER TRIASSIC

uRM	<u>McCARTHY FORMATION</u> (lower calcareous member): thin interbedded light to dark grey argillaceous limestone and dark grey argillite: (marine: <u>0 to 400 m</u> ): (115G/2,3,5-7; 115F/15; 115B/16; 115C/15; 114P/14,15).
uRC	<u>CHITISTONE and NIZINA LIMESTONES</u> : massive light grey limestone, limestone breccia and darker grey, well-bedded limestone: (shallow marine; lowest part intertidal-supratidal, local sabkha: <u>0 to 700 m</u> ): may locally include uRM: (115G/2,3,5-7,12; 115F/15; 115B/15,16; 115C/15,16; 114P/14,15).

**uRe**

white to creamy-white gypsum and anhydrite: (115G/2; 115B/15,16).

**uRN**

**NIKOLAI GREENSTONE:** dark green and maroon amygdaloidal basaltic and andesitic flows, locally interbedded with maroon tuff and breccia, and rarer maroon and green shales and thin-bedded bioclastic limestone; grey-green volcanic breccia, pillow lava and conglomerate at base: (subaerial, locally submarine: rift volcanics: 0 to 1000 m): may locally include uRC, uRM, and rare uRe: R. meta. commonly to (2b): (115G/1-3,5-7,12; 115F/10,15; 115B/15,16; 115C/16; 115A/3,5,6; 114P/14).

### MIDDLE TRIASSIC

**mTp**

dark grey phyllite (locally limy); minor thin grey limestone: (marine: 0 to 200 m): (115G/3,5,6; 115F/10; 115B/16).

### MESOZOIC AND (?) OLDER

**Msm**

biotite schist; minor amphibolite, gneiss: in part "Kluane Schist" (? metamorphosed JKd); includes older rocks: R. meta. to (4): (114P/9,15,16).

**Mc**

marbles in Msm: may be U. Triassic and/or L. Permian, or older: R. meta. to (4): (114P/9,16).

### PALEOZOIC AND (?) MESOZOIC

**PTv**

basic volcanics (locally pillowed), greenstone; rare limestone, sandstone, shale: may include Pv, uRN, and PTb, and locally (?) older rocks: R. meta. to low (3): (115A/3; 114P/9,14-16).

**PMm**

schist, amphibolite, dark argillite, marble, conglomerate; may locally include basic intrusions: (?) equivalent to Strelna Metamorphics; may locally include metamorphosed PPs, and younger rocks: R. meta. to low (4): (115C/10,15).

**PMsv**

dominantly greenstone; includes argillite, metachert; local marble, diorite, gabbro; rarer lenses serpentinite: may include PMm, PPs, and locally U. Triassic, Valdez Group, or (?) older rocks: rocks are moderately to highly deformed; mainly or all melange: R. meta. generally to low (3): (115B/2,3; 114O/8,9,14; 114P/3,5,6).

**PMvs**

- (i) Kluane Range, Donjek to White rivers area (115G/12; 115F/9,15,16): basic metavolcanics, schist, marble; minor granitic rocks and granitoid gneiss: may be U. Paleozoic and U. Triassic; may locally include (?) older and (?) younger rocks.
- (ii) area NW and SE Mt. Decoeli (115B/16, 115A/13): greenstone, greenschist; minor metagreywacke-argillite: may be equivalent to rocks of areas (i) and (iii); may locally include JKD, and (?) PRub, PRb, IKub, IKb, or KTb.
- (iii) area NW and SE Kelsall Lake (114P/9,15,16): amphibolite, greenstone; minor greenschist; rare metagreywacke-argillite: may be equivalent to rocks of areas (i) and (ii); may include metamorphosed PRv, and locally metamorphosed JKD: R. meta. to (4).

## PALEOZOIC AND/OR MESOZOIC, OR (?) OLDER

**PMgn**

gneiss complex; minor schist: R. meta. (4): (114P/9).

## CARBONIFEROUS TO PERMIAN

### PENNSYLVANIAN TO LOWER PERMIAN

#### SKOLAI GROUP: (Pvx to PPs inclusive)

**PPs**

undivided Skolai Group: volcaniclastic sediments, argillite; local limestone, marble, chert: includes felsite dyke swarms, minor syenite: R. meta. (2) to low (3): (115C/9,15,16).

### LOWER PERMIAN

#### HASEN CREEK FORMATION: (Pc, Ps)

**Ps**

thin-bedded siliceous argillite, siltstone; minor greywacke and conglomerate; local thin basaltic flows (some pillowled), breccia and tuff: (marine: 0 to 800 m): may locally include Pv, Pc, PRub, PRb; in 114P/15 may locally include U. Triassic and JKD: R. meta. to (2): (115G/2,3,5-7,12; 115F/7,15,16; 115B/15,16; 115A/3,5,6,12; 114P/15).

**Pc**

buff bioclastic limestone, calcarenite; local conglomerate at base: (marine: 0 to 100 m): equivalent to Golden Horn Limestone: R. meta. to (2): (115G/2,3,5,6,12; 115F/10,15; 115B/16; 115C/16; 115A/3,5,6,12,13).

### PENNSYLVANIAN TO (?) LOWERMOST PERMIAN

#### STATION CREEK FORMATION: (Pvx, Pv, Psv)

(oceanic arc volcanics and sediments)

**Psv**

tuff, breccia, siliceous argillite, rare andesitic flows: may locally include Ps: R. meta. to (2): (115A/3,5,6,12; 114P/14,15).

**Pv**

area NW of lower Duke River (115G/5,6,12; 115F/7,9,10,15,16): predominantly pyroclastics; includes grey-green massive volcanic breccia and agglomerate (clasts basalt and plagioclase porphyry), grading into lithic-vitric tuff, and well-bedded pale green and white siliceous tuff; rare argillite and basic flows.

Duke to Slims rivers area (115G/2; 115B/16): increasing amounts of flows, lesser amounts of pyroclastics.

area NW and SE of Mush Lake (115A/3,5,6; 114P/15): predominantly flows; includes dark green massive porphyritic (augite) basaltic to andesitic flows (locally pillowied); minor volcanic breccia, rare argillite.

locally Pv unit may include PRub, PRb, uRN: R. meta. commonly to (2b).

**Pvx**

pillowed basic flows in unit Pv: (115A/6).

## UPPER PALEOZOIC AND (?) OLDER AND YOUNGER DEVONIAN TO UPPER TRIASSIC AND (?) OLDER

Adjacent to Duke River Fault (SW side) and locally in Icefield and Alsek ranges:

(uPs to uPe inclusive): (units of this grouping not necessarily in order of age: many rocks are lithologically similar to those of late Paleozoic and early Mesozoic ages closely juxtaposing, but NE of Duke River Fault: mostly, rocks are complexly deformed: R. meta. generally high (2) to mid (3)).

**uPe**

white to creamy-white gypsum and anhydrite; locally size of lenses are exaggerated to show on map: (?) mostly uRe: (115B/16; 114P/10,14).

**uPcp**

thin-bedded, dark bluish-grey limestone or marble; local light grey, thicker bedded limestone; minor dark grey calcareous argillite: U. Triassic and (?) older: (115G/4; 115B/6-11,13,14; 115A/5).

**uPcs**

laminated dull ochre yellow calcareous siltstone-sandstone, silty limestone, grey phyllite; minor intercalated thin-bedded dark blue-grey limestone; may locally include uPcp: (?) Carboniferous or Permian; may include older, and locally Triassic rocks: (115G/3; 115B/8-10,13-15).

**uPp**

- (i) Slims to Jarvis rivers area (115B/16): black siliceous argillite, argillite (locally calcareous); includes interbedded argillite, dark siltstone; rarer buff limestone or marble, black ultramafic "pods" and dykes, and local gypsum-anhydrite: (?) equivalent to mRp, Ps, or Dp; with uPc, PRub, uRe.
- (ii) Squaw and Datlasaka ranges area (115A/3; 114P/10,14,15): similar rocks to (i): (?) mostly U. Paleozoic; may locally include older and U. Triassic.

(iii) Donjek to Disappointment glaciers area (115B/9,10,13-15): black graphitic argillite, commonly siliceous, and dark quartz-rich argillaceous siltstone; minor sandstone and impure quartzite; local dull ochre calcareous sandstone-siltstone, dark calcareous argillite and dark grey limestone or marble: (?) U. Paleozoic, and (?) older; may locally include U. Triassic.

**uPpc**

- (i) area adjacent to Duke River Fault (SW side), Steele Glacier to Marble Creek (115G/2-6; 115F/1,8; 115B/9,14-16; 115A/5): dark argillite (mainly calcareous, but locally siliceous) and interbedded dark siltstone-argillite (commonly calcareous); minor thin-bedded dark blue-grey limestone and phyllite: local "lenses" of more massive light grey limestone and greenstone (? massive volcanics and/or basic intrusions); rare gypsum-anhydrite: U. Paleozoic to U. Triassic; may locally include older.
- (ii) "Tats Glacier" and Carmine Mountain areas (114P/12 and 114P/11): similar rocks to (i) and (iii): in part or all U. Triassic; may include older.
- (iii) Squaw-Datlasaka Ranges area (115A/3; 114P/10,14,15): similar rocks to (i) and (ii): mostly U. Paleozoic to U. Triassic; may include older.

**uPc**

massive light grey and thin-bedded dark blue-grey limestone or marble; local buff-grey crinoidal limestone: mostly U. Paleozoic to U. Triassic; includes Devonian and (?) older, Pennsylvanian, (?) L. Permian, and U. Triassic: (115G/2-5; 115B/15,16; 115A/3; 114P/8-10,14,15).

**uPvs**

green and purple volcanic breccia, tuff, agglomerate: rarer cherty sediments, black argillite, grey phyllite: (?) equivalent to uPv, Pv, Psv, uRN, or older: (115B/16).

**uPv**

Slims to Jarvis rivers area (115B/15,16): dark green (locally purple), porphyritic (augite and/or plagioclase), or nonporphyritic basaltic to andesitic flows; local volcaniclastic sediments, grey limestone or marble, and basic intrusions: Devonian (? Middle or Upper) or (?) younger.

Alsek River area (115A/5): green and purple basaltic to andesitic flows: (?) U. Triassic, but may be older.

"Tats Glacier" area (114P/12): dark greenish-grey basaltic to andesitic pillow lava and massive flows; includes interbedded limy argillite-siltstone and basic intrusions; local limestone, tuff and agglomerate; rare gypsum: (marine: ? arc volcanics): U. Triassic and (?) younger; may locally include older.

Carmine Mountain area (114P/11): basaltic to andesitic pillow lava and massive to fragmental dacite; locally interbedded calcareous and cherty argillite: (marine: ? bimodal rift volcanics): (?) U. Triassic, but may be older.

**uPs**

undivided mostly units, OSwp, Dc, uPpc, uPp, uPc, Mb: (114P/8-10).

## UPPER PALEOZOIC AND OLDER AND YOUNGER DEVONIAN TO UPPER TRIASSIC AND OLDER

Psp

**"ICEFIELD RANGES PELITIC ASSEMBLAGE"**: black graphitic argillite, phyllite, and schist (commonly siliceous; locally dominantly calcareous), and dark quartz-rich argillaceous siltstone; lesser thin-bedded mica quartzite, mica quartz-rich sandstone-siltstone; rarer dark blue-grey platy limestone or marble, and thin-bedded grey limestone or marble; local thicker bedded grey limestone or marble, basic volcanics or intrusions, and gypsum-anhydrite: (diverse grouping mostly of dark grey, fine- to very fine-grained clastics; thickness unknown): may be mostly U. Paleozoic to U. Triassic, but include older: equivalent to Dp, uPpc, uPp, uPcs, uPcp; locally includes SDc, Dc, uPc, uPv, uPe; may include finer clastic (distal) equivalent of Psa (and Psa unit), COwb, COw, COwv, OSwp, and ODcs (locally ODcs unit): rocks are complexly deformed: R. meta. commonly to low (3), locally to mid-(3): (115G/4; 115B/1,2,6-14; 115A/5; 114O/15,16; 114P/2,6-8,10-14).

## DEVONIAN

### MIDDLE DEVONIAN AND, OLDER AND YOUNGER

Dc

**"BULLION CREEK LIMESTONE"**: massive to well-bedded light grey limestone or marble, thin-bedded dark grey limestone or marble; minor dark blue-grey calcareous argillite (locally interbedded, and at base): (marine: bank/reef carbonates: 0 to ? 300 m): mostly Devonian (Middle Devonian, but includes older and younger): R. meta. (2) to low (3): (115G/2,3; 115B/9,15,16; 115A/3,5,12,13; 114P/10,14,15).

## DEVONIAN AND (?) OLDER

### LOWER DEVONIAN AND (?) OLDER

Dp

dark blue-grey argillite, phyllite, and minor greywacke siltstone-sandstone; upper part more calcareous, lower part more greywacke; locally may include massive limestone (? Dc), and greenstone (uPv or uPbv): (marine): R. meta. (2) to low (3): 115G/2,3; 115B/9,15,16; 115A/12,13).

## SILURIAN AND DEVONIAN, AND (?) OLDER

SDc

massive to thick-bedded light grey limestone or marble, and well-bedded dark blue-grey limestone or marble; minor calcareous argillite or phyllite: (marine: up to ~2000 m): in part Devonian, probably includes Silurian, and locally (?) Ordovician; equivalent in part to ODcs: R. meta. (2) to (3), locally to (4): (115G/4,5; 115F/1,2,7; 115B/2,3,5-8,10-14; 115C/9,15,16; 114O/16; 114P/2,3,6,7).

## (?) ORDOVICIAN TO SILURIAN, AND (?) YOUNGER

OSwp

area adjacent to Duke River Fault (SW side), Steele Glacier to Marble Creek (115G/2-5; 115B/9,15,16; 115A/5,12): dull rusty-buff or green-grey greywacke siltstone-sandstone, and argillite or phyllite; minor grit; rarer limestone, pebble conglomerate, conglomerate: may locally include argillite (Dp): (marine: ? 0 to ~1000 m): R. meta. high (2) to mid-(3).

Squaw-Datlasaka ranges area (114P/10,14,15): interbedded rusty, buff to darker buff-grey greywacke siltstone-sandstone, dark grey argillite (commonly quartz-rich); locally includes quartzite and rarer conglomerate: may be partly equivalent to Psa: R. meta. low to mid (3).

## ORDOVICIAN AND (?) OLDER AND YOUNGER

### LOWER ORDOVICIAN TO DEVONIAN AND (?) OLDER

#### "GOATHERD MOUNTAIN CARBONATE ASSEMBLAGE":

ODcs

(a) laminated carbonate-calcareous mudstone-siltstone facies: dominantly interbedded yellow to ochre-buff calcareous mudstone-siltstone, grey silty limestone and platy to thicker bedded blue-grey, cryptocrystalline limestone; occurs generally in lower to middle parts of unit: includes Ordovician; (?) facies equivalent in part to Psa.

(b) thicker bedded to massive carbonate facies: thick to massive-bedded, light to medium grey limestone; local well-bedded, mid- to dark blue-grey limestone: occurs generally in middle to upper parts of unit: probably equivalent to SDc, but locally includes Ordovician.

Abrupt facies changes, diachronous and unconformable relationships suspected within this unit: locally facies (a) rapidly grades upward and laterally into (b): locally (a) grades downward into greywacke-argillite (C Owb) (Range Creek area): this unit is believed to grade rapidly into part of Psa (Marble Creek area): (marine: ? to or >1000 m): rocks are moderately to complexly deformed: R. meta. high (2) to mid (3): 115B/1,8; 115A/3-5; 114O/9,15,16; 114P/2,6-8,10-14).

## CAMBRIAN TO ORDOVICIAN AND (?) YOUNGER

#### "DONJEK RANGE GREYWACKE-GREENSTONE ASSEMBLAGE": (COs, COw, COwv)

(marine: volcaniclastics; minor basic volcanics and intrusions: ? arc volcanic clastics: up to ~2000 m): rocks pervasively deformed: R. meta. high (2) to mid (3): 115G/2,3; 115B/9,14-16; 115A/5,12).

COwv

massive to well-bedded, buff-green, coarse- to medium-grained greywacke and/or volcanic sandstone (clinopyroxene and plagioclase clasts abundant); minor greywacke siltstone-sandstone, argillite, phyllite or schist, and basic intrusions; rare conglomerate, basic flows and (?) pyroclastics, and thin limestone or marble.

COw

green to maroon-buff greywacke siltstone-sandstone, and argillite, phyllite, or schist; minor coarser greywacke and/or volcanic sandstone (with augite/plagioclase clasts); rare conglomerate, and limestone or marble.

**€ Os**

undivided mainly units €Ow, €Ovw; may locally include OSwp.

**"NORTHERN ALSEK RANGES GREYWACKE-GABBRO ASSEMBLAGE": (€Oc, €Owb)**

(marine: volcaniclastics; basic volcanics and intrusions: (?) arc volcanic clastics: >1000 m: (?) equivalent to €Ovw, €Ow, €Os; (?) in part facies equivalent to, or younger than €v, €c, €vs: deformation only locally intense: R. meta. high (2) to low (3); C. meta. by €Ob: 115B/1; 115A/3-5; 114P/13).

**€Owb**

greenish-grey greywacke sandstone-siltstone (volcanic derived) and dark argillite; minor laminated buff-grey silty limestone and limy siltstone, and basaltic to andesitic flows (some pillowied); rarer, more massively bedded blue-grey limestone (€Oc): grades upwards into ODcs: includes abundant gabbro-diabase sills/dykes (€Ob).

**€ Oc**

light grey laminated silty limestone and limy siltstone, and well-bedded limestone or marble: (distinct carbonate units in €Owb).

## CAMBRIAN AND (?) YOUNGER

**"FIELD CREEK VOLCANICS": (€v, €c, €vs)**

(marine and nonmarine: basaltic to andesitic flows; minor carbonate and volcaniclastics: (?) arc volcanics: ? >2000 m, base unobserved): deformation only locally intense: R. meta. high (2) to low (3): 115A/3,4).

### UPPER CAMBRIAN AND (?) YOUNGER

**€ vs**

calcareous tuff, volcanic breccia; dark green and rare purple, porphyritic andesitic flows; locally intercalated with thin-bedded limestone, and "lenses" of purple and green tuffaceous sandstone and pebble conglomerate.

**€ c**

light grey massive limestone and thin-bedded silty limestone, or marble: ("main" limestone 0 to ? ~200 m, with several, much thinner, discontinuous beds).

### UPPER CAMBRIAN AND (?) OLDER

**€ v**

massive dark green (rare purple) porphyritic (plagioclase) basaltic to andesitic flows (locally pillowied), volcanic breccia; minor tuff and tuffaceous sandstone; rare thin limestone or marble: (base not observed).

## PALEOZOIC AND YOUNGER

Pvb

massive and pillowd basaltic to andesitic flows, volcaniclastics, and gabbro-diabase intrusions; minor pyroclastics, argillite-siltstone, and limestone; rare acid volcanics: (?) mostly Paleozoic; includes Cambro-Ordovician, Devonian, and locally U. Triassic; includes units (?) Cv, (?) Cvs, COwb, COb, uPv, (?) Mb, (?) KTb, and locally (?) Cc, COc, OSwp, Dc; may locally include intrusions related to Pvs, Og, or Nv: R. meta. low to mid (3): (114P/7,8,10,11,14).

## LOWER PALEOZOIC AND (?) YOUNGER

(?) ORDOVICIAN TO (?) SILURIAN AND (?) YOUNGER

Psa

**"NORTHEASTERN ICEFIELD RANGES CLASTIC ASSEMBLAGE"**: thin to medium-bedded, fine- to medium-grained, khaki buff, quartz-rich, micaceous, variably calcareous siltstone and sandstone, mica quartzite, psammitic schist, intercalated with minor thin grey phyllite, argillite and schist; includes minor volcanic wacke and rarer thin buff ochre-orange limestone or marble, and silty limestone: (monotonous sequence of quartz-rich, calcareous, fine- to medium-grained clastics: marine: thickness unknown): may be partly equivalent to OSwp, and partly facies equivalent to ODcs; may include distal equivalents of COw, COwv: rocks complexly deformed: R. meta. low to mid (3): (115G/3,4; 115B/8-10,13-15; 115A/5).

## (?) LOWER PALEOZOIC AND (?) YOUNGER

IPcs

rusty brown, thin-bedded limestone, marble; minor greenstone: R. meta. variable, low (3) to (4): (115F/2; 115C/15,16).

## (?) CAMBRIAN TO (?) ORDOVICIAN AND (?) YOUNGER

IPvs

greenstone, amphibolite; minor schist, argillite, greywacke, mica quartzite; locally includes marble: R. meta. to low (4): (115F/1,2,7; 115B/5,6,12; 115C/8,15,16; 114O/15).

## (?) CAMBRIAN AND (?) YOUNGER

IPvm

massive dark green to black amphibolite: (? basic metavolcanics): R. meta. to (4): (115B/2,3,7; 114O/9,10,15,16).

## (?) LOWER PALEOZOIC AND (?) OLDER

IPgn

**"VERN RITCHIE GLACIER GNEISSIC ASSEMBLAGE"**: grey gneiss and granitoid gneiss; includes amphibolite, amphibole schist, quartz mica schist, metaquartzite and marble: may be partly equivalent to IPvm, IPvs: locally complexly deformed: R. meta. to (4): (114O/9,10,15; 114P/6,11,12).

# INTRUSIVE ROCK MAP UNIT DESCRIPTIONS

## TERTIARY

### NEOGENE

#### MID TO LATE MIOCENE

##### WRANGELL PLUTONIC SUITE: (6-16 Ma): (IMdi, IMg, IMf)

(calc-alkaline: subvolcanics; basalt, andesite, rhyolite: high-level plutons; granodiorite, diorite, gabbro: comagmatic with Wrangell Lava).

IMf

subvolcanics: (6-16 Ma): white to creamy-white hornblende and/or biotite rhyolite, rhyodacite, dacite, and trachyte; felsite: suite includes unmapped basalt, andesite, gabbro: (comprise sills, dykes, domes, small plutons): in part or all intra-Wrangell Lava: may include Of: (115G/3,5; 115F/7,8,15; 115B/9; 115A/3-5,12).

##### high-level plutons: (IMdi, IMg)

IMg

Mt. Steele pluton: (8.4 Ma): fine medium grained, creamy-grey hornblende and/or biotite granodiorite, and porphyritic (K-feldspar) hornblende granodiorite; includes creamy-grey, biotite-quartz-plagioclase leucogranite porphyry (equivalent to IMf): extent of phases and pluton uncertain, due to ice cover: probably includes JKg: (115F/1,2,7; 115C/16).

IMdi

Bock's Brook stock: (12.8-13.3 Ma): includes narrow border phase of dark grey, medium grained, pyroxene gabbro, cored and intruded by light buff-grey, medium grained, uniform biotite diorite: both phases cut by younger Wrangell dykes: (rounded, discordant, concentrically zoned pluton: contacts sharp; C. meta. of gabbro by diorite): (115G/2).

## PALEOGENE

### OLIGOCENE

Of

subvolcanics: (26-32 Ma): white to creamy-white felsite (? rhyolite, rhyodacite); biotite and/or hornblende quartz latite porphyry; locally crowded hornblende plagioclase porphyry: (mainly sills, dykes; some small plutons): may include IMf: (115G/2,6; 115B/7,15,16; 115A/3,5,6,11,13; 114P/2,6,7,9,10,14,15).

##### TKOPE PLUTONIC SUITE: (24-31 Ma): (Ogb to Og inclusive)

(calc-alkaline: epizonal: elongate, discordant, composite batholithic complexes and plutons: granite; lesser granodiorite, gabbro; rarer quartz diorite and granophyre: cut by diabase, quartz feldspar porphyry dykes: contacts sharp; thermal aureoles; C. meta. (1) to (2)).

Og

undivided Tkope plutonic suite: includes light pinkish-grey, medium- to coarse-grained, homogeneous, biotite and/or hornblende granite (locally miarolitic); lesser light creamy-grey biotite hornblende granodiorite, dark grey biotite hornblende quartz diorite and gabbro-diorite: locally includes older and (?) younger plutons: (114P/7-11).

Øgb-  
Øgp

Tkope River batholith: (24.7-31.1 Ma): (Øgb to Øgp inclusive)

- Øgp: fresh, light pink, fine grained, homogeneous, miarolitic, biotite granophyre: (114P/11).
- Øgr: fresh, pink, medium grained, homogeneous, hornblende biotite granite: dominant phase: (114P/7,10,11).
- Øgd: altered, dark grey, massive, fine- to fine medium-grained, heterogeneous, porphyritic (plagioclase), biotite hornblende (rarer augite) granodiorite: (114P/7,10).
- Øgdp: altered, grey, massive, medium grained, biotite granodiorite porphyry (phenocrysts; plagioclase, quartz): (114P/7,10).
- Øqd: altered, white to light grey, massive, medium grained, biotite hornblende quartz diorite: (114P/11).
- Øgb: altered, dark grey to black, biotite augite hypersthene microgabbro to gabbro: (114P/7,10,11).

Other high-level plutons:

Øggb

Mt. Newton pluton: (27.4 Ma): foliated, dark greenish-grey, fine coarse grained, homogeneous, hornblende gabbro-diorite: (115C/7).

Mt. Owen pluton: light grey, fine- to medium-grained, homogeneous, biotite hornblende granite; rhyolite porphyry: composite body; may include Eg and/or younger: (115C/1,8).

EOCENE

Eg

SEWARD PLUTONIC SUITE: (41-52 Ma): nonfoliated to weakly foliated, light to mid-brownish-grey, medium grained, homogeneous, biotite and hornblende biotite - tonalite, granodiorite; rarer granite and quartz diorite: (calc-alkaline: epizonal: elongate, discordant and concordant, single or multiphase plutons: contacts sharp; distinct thermal aureoles): may locally include younger plutons: (115B/4,5; 115C/1,8-10; 114P/3-6).

## CRETACEOUS TO TERTIARY

### LATE CRETACEOUS TO EARLY TERTIARY

KTg

COAST PLUTONIC COMPLEX (Northern part): (55.5-72.4 Ma): biotite hornblende granodiorite, biotite hornblende quartz diorite; lesser biotite granite: (multiphase plutonic complex): (115B/16; 115A/2; 114P/9,15,16).

KTb

medium- to coarse-grained, dark grey hornblende pyroxene gabbro; minor hornblende quartz diorite, diabase: (?) equivalent to 1Kb or Tertiary: (115A/2; 114P/7,10,15).

## CRETACEOUS

### LATE EARLY CRETACEOUS

**KLUANE RANGES PLUTONIC SUITE:** (106-121 Ma): (Kqm to Kg inclusive) (calc-alkaline: epizonal to (?) mesozonal: elongate, discordant and concordant, batholithic complexes and single or multiphase plutons: granodiorite, quartz diorite, diorite; rarer quartz monzonite: contacts sharp; thermal aureoles; C. meta. (1) to (2): may include older and younger plutons).

#### undivided Kluane Ranges plutonic suite:

Kqm-  
Kg

Kg: multiphase batholithic complexes and small plutons: (115G/2,5,6,12; 115F/9,10,15,16; 115B/15,16; 115A/2,3; 114P/14,15).

#### subdivided Kluane Ranges plutonic suite: (Kqm to Kgd inclusive)

Kgd: buff to mid-grey, medium grained, biotite hornblende granodiorite: (115G/2; 115A/6).

Kqd: mid-grey, coarse medium grained, biotite hornblende quartz diorite: (115A/3,6).

Kdi: mid- to dark greenish-grey, coarse medium grained, hornblende diorite: (115B/16; 115A/3,5,6).

Kqm: buff to mid-grey, coarse medium grained, biotite hornblende quartz monzonite: (115A/5).

#### **Pyroxenite Creek Ultramafic Complex:** (113-124 Ma): (IKb, IKb)

(Alaskan-type complex: epizonal: includes three crudely concentric clinopyroxenite phases: cut by mafic and ultramafic dykes, and small mafic plutons: contacts sharp; thermal aureole; C. meta. to (3): 115A/12)).

IKb

medium grained hornblende pyroxene gabbro, and biotite hornblende diorite: small plutons intruding ultramafite.

IKub

two central phases: (a) coarse grained olivine clinopyroxenite (minor biotite); (b) fine grained magnetite clinopyroxenite (minor hornblende); narrow border phase of hornblende clinopyroxenite.

## JURASSIC TO CRETACEOUS

### LATE JURASSIC TO EARLIEST CRETACEOUS

SAINT ELIAS PLUTONIC SUITE: (130-160 Ma): (JKgr, JKgd, JKg)

(calc-alkaline: crudely comprise (a) narrow western belt (mainly U. Jurassic); (?) mesozonal to epizonal; elongate, concordant/discordant batholithic complexes; foliated tonalite, quartz diorite, diorite, lesser granodiorite, quartz monzodiorite, rarer granite: (b) broad eastern belt (mainly L. Cretaceous); epizonal; elongate to rounded, discordant, single or multiphase plutons; granodiorite, lesser tonalite, rarer granite, quartz monzodiorite, quartz diorite, diorite: contacts sharp, thermal aureoles, C. meta. (1) to (2): may include older and younger plutons).

#### undivided Saint Elias plutonic suite:

JKgr-  
JKg

JKg: mainly batholithic complexes and multiphase plutons: (115G/4; 115F/1,2; 115B/1,2,5-9,12-15; 115C/9,10,15,16; 115A/4,5; 114O/8,9,15,16; 114P/2,3,5-7,11-14).

#### subdivided Saint Elias plutonic suite: (JKgr, JKgd)

JKgd: mostly nonporphyritic and porphyritic (K-feldspar), biotite hornblende granodiorite; lesser nonporphyritic biotite and/or hornblende tonalite; locally includes biotite hornblende quartz monzodiorite, biotite hornblende quartz diorite, dark green-grey diorite and hornblendite: phases mostly light to mid-grey, coarse medium grained, nonfoliated and homogeneous: (115G/3,4; 115F/1; 115B/1,8,10,11,13-15; 115C/15,16; 115A/4; 114O/16; 114P/13).

JKgr: light creamy-grey, medium grained, homogeneous, nonfoliated hornblende biotite granite, quartz monzonite: (115F/2; 115C/10,15; 115A/5).

## MESOZOIC AND/OR OLDER

### LATE PALEOZOIC AND/OR MESOZOIC

Mb

"SQUAW-DATLASAKA RANGES GABBRO-DIABASE SILLS": medium to dark rusty grey-green, massive to schistose greenstone: predominantly metagabbro or metadiabase sills; may locally include basic volcanics and minor sediments: (?) equivalent to PRb: may include uPv and (?) older rocks: R. meta. high (2) to low (3): (115A/3; 114P/10,14,15).

## (?) PERMIAN AND TRIASSIC

### LATE TRIASSIC AND (?) OLDER

"KLUANE-TYPE MAFIC-ULTRAMAFICS": (PRub, PRb)

PRb

medium grey-green, massive, medium grained, pyroxene gabbro sills: may locally include IKb: R. meta. (2) to low (3): C. meta. by larger sills (1) to (2): (115G/2,3,5,6; 115B/15,16; 114P/15).

PRub

sheeny black peridotite; rare dunite: sills, "pods": may locally include IKub: R. meta. (2) to low (3); C. meta. by larger intrusions to (3): (115G/2,3,5,6; 115F/15,16; 115B/16; 115A/3,5,6; 114P/14,15).

## LATE PALEOZOIC AND (?) YOUNGER (?) DEVONIAN AND/OR YOUNGER

**uPbv**

"**MT. CAIRNES GABBRO-GREENSTONE COMPLEX**": complex of massive coarse medium- to coarse-grained, mid- to dark rusty grey-green hornblende pyroxene gabbro, minor medium grained gabbro-diabase and gabbro-pegmatite intrusions, and rare small black peridotite "pods"; with screens of green or lesser purple flows, pyroclastics, and volcaniclastics, minor dark siliceous argillite, and rare chert: may include PRb, PRub, Mb, IKb, KTb, and (?) IMdi; with screens of uPv, uPvs; similar to uPb: rocks are variably deformed: R. meta. high (2) to mid-(3): (115G/2; 115B/9,15,16; 115A/12).

## LATE PALEOZOIC OR (?) OLDER

**uPb**

"**STEELE CREEK - MT. CONSTANTINE GABBRO COMPLEXES**": (364 Ma): complexes of massive, fractured, dark rusty grey-green hornblende pyroxene gabbro, gabbro-pegmatite and diabase: (115G/5; 115F/7).

## CARBONIFEROUS TO PERMIAN

### LATE PENNSYLVANIAN TO EARLY PERMIAN

**ICEFIELD RANGES PLUTONIC SUITE**: (270-290 Ma): (PPgd to PPg inclusive) (alkaline to calc-alkaline: (?) mesozonal to epizonal: elongate to rounded, concordant/discordant, agmatite or multiphase batholithic complexes and simple-phase plutons: commonly quartz monzodiorite to diorite phases are veined and intruded by leucocratic quartz syenite to granodiorite: several plutons are uniform granodiorite or syenite: contacts both complex and sharp; some thermal aureoles; C. meta. (1) to (2): may locally include JKg and (?) younger intrusions).

#### undivided Icefield Ranges plutonic suite:

PPg: multiphase batholithic complex and pluton: (115G/3-6; 115B/13).

#### subdivided Icefield Ranges plutonic suite: (PPgd to PPdi inclusive)

PPdi: mid- to dark greenish-grey, medium grained, nonfoliated and foliated, biotite hornblende quartz monzodiorite-quartz diorite-diorite, veined and intruded by leucocratic granodiorite: (agmatite complexes): (115B/2,3,6,7,12,13).

PPy: salmon pink, uniform, medium grained, hornblende syenite: (small simple-phase plutons): (115C/9,16).

PPydi: mid- to dark greenish-grey, medium grained, nonfoliated and foliated biotite and/or hornblende quartz monzodiorite-monzodiorite-diorite, veined and intruded by pink to creamy-grey, leucocratic quartz syenite to granodiorite: (syenodiorite complexes): (115B/6,11-14; 115C/15,16).

PPqdi: Art Lewis Glacier pluton: complex dominantly composed of dark greenish-grey, altered, medium grained, foliated biotite hornblende quartz diorite-diorite; cut by numerous fine grained leucocratic dykes and granitic pegmatite: (114O/15).

**PPgd-  
PPg**

PPgd: light buffish-grey, coarse medium grained, fairly homogeneous, nonporphyritic and porphyritic (K-feldspar) hornblende and/or biotite granodiorite: includes Steele Glacier and Fisher plutons: (115G/4; 115F/1,8; 115B/2).

## CAMBRIAN TO ORDOVICIAN, AND (?) YOUNGER

€ Ob


### "NORTHERN ALSEK RANGES GABBRO-GREENSTONE COMPLEX":

variably sized, massive, dark rusty grey green, gabbro or diabase sills and dykes (>80%), with screens of rusty € Ow: includes basic volcanics (? Cv, ? € vs, and € Ow); may include younger intrusions: R. meta. high (2) to low (3); C. meta. by larger sills (1) to (2): (115B/1; 115A/3-5; 114P/13,14).

## MAP SYMBOLS


### Geological boundary

defined, approximate, assumed  
gradational  
outcrop limit/drift boundary  
limit of geological mapping



### High-angle Fault

defined, approximate, assumed  
relative displacement:  
strike-slip; downthrow (barb)



### Thrust Fault (teeth upthrust side)

defined, approximate, assumed




### Holocene Fault Scarp

defined, assumed



### Fold

anticline : overturned anticline  
syncline : overturned syncline  
homocline : plunge of fold  
fold axial trace  
defined, approximate, assumed



### Bedding: strike/dip

tops known:  
horiz., inclin., vert., overturned  
tops unknown:  
inclined, vertical



### Foliation, cleavage: strike/dip horizontal, inclined, vertical



### Lineation: minor fold axes



Eruptive centre



Breccia pipe

"b.p."

Hydrothermal alteration zone

"h.a.z."

## Mineral deposits

Locality: map to legend reference number  $\otimes^4$

## Macro and micro fossils

Locality: macro, conodont, macro and conodont      ◎, ●, ▲

## Numerical age determinations

(Dodds and Campbell, 1988; Farrar et al., 1988)

Numerical age determination locality

**coding:** material, method, age  
(multiple determinations grouped) hK267 (e.g.)

method: potassium-argon, rubidium-strontium,  
fission-track K, R, F

age: in millions of years 267 (e.g.)

## **NOTES ON MAP UNIT DESCRIPTIONS**

- I colours in rock descriptions, refer to weathered surfaces.

## II (R. meta.; C. meta.)

### **Regional Metamorphism (R. meta.):**


- (1) unmetamorphosed.
  - (2) subgreenschist facies.
  - (2a) laumontite-prehnite-quartz facies.
  - (2b) pumpellyite-prehnite-quartz facies.
  - (3) greenschist facies.
  - (4) amphibolite facies.

### Contact Metamorphism (C. meta.):

- (1) albite-epidote - hornfels facies.  
 (2) hornblende - hornfels facies.  
 (3) pyroxene - hornfels facies.

三

National Topographic System (NTS) grid, Saint Elias Mountains and adjoining areas. Location of map units are indicated (in legend) by this system: for example (115F/1); or (115F/7-10, inclusive).



**Figure 3-**

## MAPPING SOURCES

**Geology mostly from Operation Saint Elias: 1973-74, 1977-79, 1981, 1983, 1986-87**

### Project leaders

R.B. Campbell: (1973-1983)  
 R.B. Campbell & C.J. Dodds: (1983-1988)  
 C.J. Dodds: (1988-1990)

### Major contributors, Operation Saint Elias:

R.B. Campbell (1973-74, 1977-79, 1981, 1983)	Paleozoic and younger rocks in Icefield, Fairweather, Alsek, Kusawak and Boundary ranges.
C.J. Dodds (1974, 1977-79, 1981, 1983, 1986-87)	Paleozoic and younger rocks SW of Duke River Fault in northeastern Icefield, Donjek, Kluane, northern Alsek, Squaw and Datlasaka ranges.
G.H. Eisbacher (1973-74)	Dezadeash Group and Amphitheatre Formation rocks in Kluane Ranges.
J.W.H. Monger (1974)	Pre-Cenozoic rocks in Kluane Ranges (between Denali and Duke River faults).
P.B. Read (1974-75, 1977)	Pre-Cenozoic rocks in Kluane Ranges (between Denali and Duke River faults) and in northernmost Alsek Ranges.
J.G. Souther (1973-74)	Wrangell lava rocks of St. Clare, Alsek and Canyon Mountain Provinces.

### Major previous mapping sources:

E.D. Kindle (1953): parts of SW Dezadeash map area.  
 J.E. Muller (1967): parts of SW Kluane Lake map area.  
 K.deP. Watson (1948): parts of Tatshenshini River map area.  
 J.O. Wheeler (1963): parts of Mount St. Elias map area.

Geological compilation and cartography by C.J. Dodds

**Topographic bases:** Surveys & Mapping Branch, Energy, Mines and Resources Canada.

Kluane Lake, 115G & F(E1/2) (edition 1, 1961); Mount St. Elias, 115B & C(E1/2) (edition 2, 1967); Dezadeash, 115A (edition 1, 1951); Yakutat, 114O (edition 1, 1966); Tatshenshini River, 114P (edition 2, 1975).

N.B. Kluane National Park Reserve and Kluane Game Sanctuary boundaries added to topographic bases from information courtesy D.E. Culham, Regional Surveyor Yukon Territory, Energy, Mines and Resources Canada (pers. comm., 1990).

Table 2. MINERAL DEPOSITS SW KLUANE LAKE MAP AREA (115G AND F [E1/2])

OF.	INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrain	Mgeo Belt	NTS (115)	Status References
1	(1)	SUGDEN (METALLINE)	Cu	vn	hyd	Pv?	W2	IB	G/02E (7)op	1,p118;9;58,p35;59
2	(42)	DISCOVERY (WHITE RIVER)	Cu,Ag	dis,vn	vlc,hyd,rpl	uTN	W2	IB	F/15W (7)op	2,p53;3,p139;5,p108; 6,p85;10,p46;11,p247; 17;18,p138;22,p51; 27,p68;28,p40;29,p138; 30,p130;31;40,p204; 58,p35;59
3	( )	SPY	Cu,Ni;Zn,Pb,Fe	mas,dis,vn	mgn;hyd,rpl	PRub;Ps	W2	IB	G/02W (7)op	8,p66;9,p56;10,p56;17; 58,p35;59
4	(70)	CONGDON (I)	Cu	dis	mgn	PRub	W2	IB	G/02W (7)op	9;14,p247;58,p35;59;62
5	( )	DC	Ni,Cu	dis,vn	mgn;hyd,rpl	PRub;Ps	W2	IB	G/02W (6)op	5,p111;9,p55;10,p47; 17;58,p35;59
6	(5)	DICKSON	Cu,Ni,PGE,Co	dis,vn	mgn;hyd,rpl	PRub,b;Ps	W2	IB	G/02W (6)op	5,p111;9,p55;10,p47; 15,p165;17;58,p35;59; 62
7	(6)	DESTRUCTION (RAM)	Ni,Cu,Co,Pt	mas,dis,vn	mgn;hyd,rpl	PRub,b;Ps	W2	IB	G/02W (6)op <sup>1</sup>	14,p235;15,p165;17; 58,p35;59
8	(11)	WADE	Cu,Ag,Au	unk	unk	uTN?	W2	IB	G/06W (7)op	9,p63;10,p59;13,p314; 16;17;58,p35;59
9	(12)	CORK	Cu,Mo	stw,bx,dis	pfp	Og;Ps	Op;W2	IB	G/06W (5)op	7,p101;9,p64;10,p49; 13,p315;16;17;27,p73; 32,p918;58,p35;59; 60,p1
10	(13)	GLEN (TATAMAGOUCHE)	Ni,Cu,PGE;Pb,Zn	mas,dis,vn	mgn;hyd,rpl	PRub;Ps	W2	IB	G/06W (6)op <sup>1</sup>	9,p56;10,p49;11,p246; 12,p178;13,p315; 15,p165;16;17;27,p72; 58,p35;59
11	(13)	GLEN (BURWASH)	Cu,Au,Mo	dis,vn	pfp?	Og;Pv	Op;W2	IB	G/06W (6)op <sup>1</sup>	9,p56;10,p49;11,p246; 12,p178;13,p315; 15,p165;16;17;58,p35; 59;60,p1

OF. No.	INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (115)	Status References
12 ( )	GLEN (MARY)	Cu,Ni,Co,Au,Ag	dis,sig	mgm	PRub,Ps	W2	IB	G/06W (6)op <sup>1</sup>	8,p65;9,p56;10,p56;17; 58,p35;59	
13 (15)	JACQUOT	Cu,Ag	stb;yn,dis	vlc;hyd,rpl	uTN	W2	IB	G/06W (7)op <sup>1</sup>	4,p30;5,p109;6,p85; 7,p103;9,p60;10,p49; 17;29,p65;58,p35;59	
14 (15)	MARY AND TEDDY	Cu	stb;yn,dis	vlc;hyd,rpl	uTN	W2	IB	G/06W (7)op <sup>1</sup>	5,p109;6,p85;7,p103; 9,p60;10,p49;17; 29,p65;58,p35;59	
15 (14)	BURWASH	Cu	stg	vlc;rpl	uTN	W2	IB	G/06E (7)op <sup>1</sup>	4,p31;15,p164;17; 58,p35;59	
16 (16)	QUILL (RAM)	Cu,Ag	stb;yn,dis	vlc;hyd,rpl	uTN	W2	IB	G/06W (5)op	5,p109;9,p61;10,p47; 15,p166;17;22,p53; 27,p70;58,p35;59	
17 (17)	LINDA (LOWER)	Ni,Cu,PGE	mas,dis,yn	mgm;hyd,rpl	PRub,b,Pv	W2	IB	G/06W (6)op	5,p109;6,p85;9,p61; 10,p48;14,p237; 15,p166;17;22,p53; 27,p70;34,p195;35; 36,p662;37;58,p35;59; 62	
18 (17)	LINDA (UPPER)	Ni,Cu,PGE	mas,dis	mgm	PRub,b,Pv	W2	IB	G/06W (7)op	5,p109;6,p85;9,p61; 14,p237;15,p166; 22,p53;27,p70;58,p35; 59;62	
19 (18)	WELLGREEN (EAST ZONE)	Ni,Cu,Co,PGE,Au	mas,dis	mgm	PRub,b,Pv	W2	IB	G/05E (3)op	5,p110;7,p100;8,p64; 9,p57;10,p48;13,p317; 14,p252;15,p166;17; 19,p36;22,p52;28,p43; 34,p184;35;36,p662;37; 38,p41;39,p55;44; 45,p96;58,p35;59;62	
20 (18)	WELLGREEN (WEST ZONE)	Ni,Cu,Co,PGE,Au	mas,dis	mgm	PRub,b,Pv	W2	IB	G/05E (3)op	5,p110;9;8,p64;9,p57; 10,p48;17;34,p184; 37,p11;39,p55;58,p35; 59;62	

OF. No.	INAC (No)	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (115)	Status References
21	(18)	WELLGREEN (NORTH ZONE)	Ni,Cu,PGE,Au	mas,dis	mgm	P Rub,b,Pv	W2	IB	G/05E	(6)op 9;10,p48;17;34,p184; 37,p11;39,p55;58,p35; 59;62
22	( )	NICKEL CREEK (MULLER)	Ni,Cu	dis	mgm	P Rub,b,Pv	W2	IB	G/05E	(7)op 5,p110,map?;10,p48; 17;34,p184;37,p11; 39,p55;58,p35;59
23	(19)	AIRWAYS (ARCH)	Cu,Ni,PGE,Co	mas,dis	mgm	P Rub,b	W2	IB	G/05E	(6)op 5,p110;9,p54;10,p47; 11,p245;12,p179; 13,p317;14,p238; 15,p167;34,p192;35; 37,p7;58,p35;59;62
24	(19)	AIRWAYS (FW)	Cu,Ni,PGE,Co	mas,dis	mgm	P Rub,b	W2	IB	G/05E	(6)op 5,p110;9,p54;10,p47; 11,p245;12,p179; 14,p238;15,p167;17; 34,p192;35;37,p7; 58,p35;59;62
25	(20)	MUSKETEER	Cu	dis	hyd,pl	Pv	W2	IB	G/12E	(7)op 9;14,p240;17;58,p35; 59;62
26	(22)	ST. ELIAS	Mo	vn,dis	hyd,pl	uPpc	AX	IB	G/05W	(7)?ip 15,p164;17;19,p36; 58,p35;59
27	(23)	MINERAL RIDGE (SHARP)	Mo,Cu	vn,dis	prp	PPgd?;Or? Mt?	Mg? Og? Pg,AX	IB	F/01E	(7)?ip 5,p112;15,p164; 10,p46;17;19,p36; 41,p636;42,p41;58,p35; 59;60,p1
28	(24)	GALLOPING	Mo	dis	prp?	?	?AX	IB	F/01E	(7)?ip 5,map;15,p164; 17;19,p36;58,p35;59; 60,p1
29	(26)	GARLIC	Cu,Au,Mo,Fe	vn	prp?	PMvs?;?	?W2	IB	F/09E	(7)op 11,p245;16;17;58;59; 60,p1
30	(27)	LIBERTY	Au,Cu,Pb,Zn	vn	hyd	Pv	W2	IB	F/16E	(7)op 15,p167;58,p35;59
31	(29)	CATS AND DOGS	Cu,Ni	dis	mgm	P Rub	W2	IB	F/16W	(7)op 11,p245;14,p241;34;37; 58,p35;59;62
32	(30)	MEXICO	Cu,Fe	stg	skn	Pc?	W2	IB	F/16W	(7)op 15,p164;17;58,p35;59; 60,p1

OF.	INAC No.	Deposit(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgso Belt (115)	Status References
33	(57)	PICKHANDLE LAKE (M)	Cu	stb;vn,dis	vc;hyd,tpl	Pv?	W2	IB	F/16W (7)op 6,p85;10,p47;11,p246; 17;30,p130;34,p180; 43,p165;58,p35;59;62
34	(44)	KLETSAN (K-CU)	Cu,Ag	dis,vn	vc;hyd	uRN	W2	IB	F/10W (7)op 2,p51;3,p133;5,p108; 10,p46;11,p246;17; 28,p42;57,p379;58,p35; 59
35	(33)	CANALASK (MICRO)	Ni,Cu,Co,Zn,PGE,Fe	dis,mas,vn,stg	mgm;hyd,tpl	Pv,PRub,b	W2	IB	F/15E (2)op 5,p111;8,p60;10,p46; 14,p242;15,p169;17; 28,p39;34,p197;35; 36,p662;37,p8;44,p81; 58,p35;59;62
36	(34)	CANALASK (EPIC)	Cu,Mo	vn	hyd	Pv	W2	IB	F/15E (7)op 10,p46;11,p245;17;58; 59;62
37	(35)	TAYLOR (ARN)	Au,Cu,Mo,Fe	unk	skn	uRN	Kg7:W2	IB	F/15E (7)op 11,p245;58,p35;59; 60,p1;62
38	(36)	RAY (SANPETE)	Cu,Fe	dis	skn	uRC	Kg7:W2	IB	F/15W (7)op 10,p47;11,p245;17; 38,p37;58,p35;59;60,p1
39	(40)	RABBIT CREEK (CC)	Cu,Ag	vn,stg	hdv,tpl	Pv	W2	IB	F/15W (7)op 3,p123;10,p47;11,p245; 17;58,p35;59
40	(41)	LEP	Cu,Zn,Pb,Fe,Ag	mas,pod	skn	Pc?	Kg:W2	IB	F/15E (7)op 10,p47;11,p245;17; 28,p73;33,p38,p38; 58,p35;59;60,p1
41	( )	WHITE RIVER (MULLER)	Cu,Ag	stb;vn,dis	vc;hyd,tpl	uRN	W2	IB	F/15W (6)op 5,p108,map;58,p35;59
42	( )	MUS	Ni,Cu	dis	mgm	PRub	W2	IB	G/05E (7)op 9;58,p35;59;62
43	( )	SHARE	Cu,Fe	dis	unk	Pv	W2	IB	F/15E (6)op 58,p35;59;62
44	(56)	NICK	Ni,Cu	unk	mgm?	?	AX	IB	G/05W (7)ip 15,p165;58,p35;59
45	(25)	MT. STEELE (ICEFIELD)	Cu,Mo,Fe	vn,dis	prp?	Mg	Mg AX	IB	F/01W (7)ip 5,map;58,p35;59;60,p1
46	(58)	CANYON MOUNTAIN (CAN)	Cu,Ag	stw,vn,dis	hyd,tpl,prp?	Nw,IMf	Mg:Vs/W2	IB	F/15W (7)op 11,p246;58,p35;59; 60,p1

OF. No.	INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS Belt	Status References
47	(69)	ONION	Ni,Cu,PGE	mas,dis,vn	mgm;hyd,rpl	P <sup>+</sup> Rub,b,Pv	W2	IB	F/15E (6)op	13,p318;14,p245; 15,p169;58,p35;59;62
48	(80)	DUKE SOUTH	Cu,Ni,PGE	pod,mas,dis,vn	mgm;hyd,rpl	P <sup>+</sup> Rub,b,Ps	W2	IB	G/02W (7)op	9;14,p249;58,p35;59;62
49	(89)	WASH	Cu,Ni,PGE	dis	mgm	P <sup>+</sup> Rub,b	W2	IB	G/06W (7)op	9;14,p250;58,p35;59;62
50	( )	TECK	Au,Ag	dis,vn	hyd,rpl	P <sup>+</sup> Rub,Pv	W2	IB	G/06E (7)op <sup>1</sup>	9,p68;58,p35;59
51	( )	PAT	Cu	dis	mgm?;hyd?,rpl?	P <sup>+</sup> Rub,b, u <sup>+</sup> RN?	W2	IB	G/06W (7)ip	9,p63;10,p59;17; 58,p35;59
52	( )	PTARMIGAN- WINDGAP	Ab	vn	hyd,rpl	P <sup>+</sup> Rub	W2	IB	G/06E (7)op <sup>1</sup>	9,p68;17;58,p35;59
53	(8)	DUKE	Ab	vn	hyd,rpl	P <sup>+</sup> Rub	W2	IB	G/06E (7)op <sup>1</sup>	9;17;58,p35;59
54	( )	MAPLE CREEK (VUG)	Cu	vn	hyd,rpl	u <sup>+</sup> RN	W2	IB	G/05E (7)op	9;10,p49;17;58,p35;59
55	(68)	KELLI (REED CREEK)	Au,Cu,As	vn	hyd	Pv,Of	Φg?;W2	IB	G/12E (7)op	9;13,p318;15,p169; 58,p35;59;60,p1;61,p9
56	(17)	VERSLUCE (LINDA)	Cu,Ag	dis,vn	unk	Pv	W2	IB	G/06W (7)op	9,p61;14,p237;15,p166; 58,p35;59
57	(64)	SWEDE JOHNSON (SJ)	Au,Cu	dis;flt	unk	?	W2	IB	G/12E (7)op	9;11,p247;13,p317; 14,p248;58,p35;59
58	(74)	SWEDE	Au,PGE	unk	unk	?	W2	IB	G/12E (7)op	9;14,p248;15,p170; 58,p35;59
59	(66)	WADE CREEK	Au	unk	unk	?	Tvs;W2?;W2?	IB	G/05E ( )op	9;12,p179;58,p35;59
60	( )	SEXSMITH	Cu	dis	mgm	P <sup>+</sup> Rub	W2	IB	G/12W (6)op <sup>1</sup>	58,p35;59;62
61	( )	QUILL CREEK (FOSSIL #1)	Cu	dis,vn	vlc;hyd,rpl	u <sup>+</sup> RN	W2	IB	G/06W (7)op	9;10,p48;17;28,p70; 58,p35;59
62	( )	QUILL CREEK (FOSSIL #2)	Cu	dis,vn	vlc;hyd,rpl	u <sup>+</sup> RN	W2	IB	G/06W (7)op	9;10,p48;17;28,p70; 58,p35;59
63	( )	QUILL CREEK (HUDSON BAY)	Cu	dis,vn	vlc;hyd,rpl	u <sup>+</sup> RN	W2	IB	G/06W (7)op	9;10,p48;17;22,p53; 58,p35;59
64	(37)	HUMP	Au,As	flt,(vn,dis)	(hyd)	?	NW2	IB	F/15W (8)op	46,p193;58,p35;59
65	(31)	PICKHANDLE	Cu,Ni	flt,(unk)	(mgm)	P <sup>+</sup> Rub	NW2	IB	F/16W (8)op	15,p164;58,p35;59

OF. No.	INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrain	Mgeo Belt (115)	Status References
66 ( )	STEELE GLACIER (SHARP)	Mo,Cu,Fe	fit,(vn,dis)	(prp?)	Mf?	/AX	IB	G/05W	(8)ip 5,p112;19,p36;41,p636; 42,p41;58,p35;59;60,p1
67 ( )	NIAMODLAOC MOUNTAIN 1	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/06E	(7)op <sup>1</sup> 3,p14;14,p32;5,p113; 7,p15;9,p71;17; 25,p20;26,p18;47,p19; 48,p319;58,p35;59
68 ( )	NIAMODLAOC MOUNTAIN 2	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/03E	(7)ip <sup>1</sup> 3,p14;14,p32;5,p113; 7,p15;9,p71;17; 25,p20;26,p18;47,p19; 48,p319;58,p35;59
69 (9)	GRANITE CREEK (HOGE)	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/06W	(7)ip <sup>1</sup> 5,p113;9,p69;11,p245; 17;25,p20;26,p18; 48,p319;47,p19; 49,pA50;58,p35;59
70 (10)	AMPHITHEATRE	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/06W	(7)op <sup>1</sup> 4,p32;5,p113;7,p9; 9,p69;11,p245;17; 25,p20;26,p18;42,p42; 48,p319;49,pA50; 58,p35;59
71 (21)	CEMENT	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/05W	(7)op <sup>1</sup> 5,p113;7,p9;9,p69; 11,p245;17;25,p20; 47,p21;26,p18;47,p19; 48,p319;49,pA50; 58,p35;59
72 (39)	MCLELLAN (COAL CREEK)	coal	cnc,stf	sed	0s	Tvs/W2	IB	F/15W	(7)op <sup>1</sup> 3,p14;11,p245;17; 58,p35;59
73 (38)	MEMOIR	coal	cnc,stf	sed	0s	Tvs/W2	IB	F/15W	(7)op <sup>1</sup> 3,p14;11,p245;17; 58,p35;59
74 (2)	STOVE 1	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/02E	(7)ip 1,p18;4,p32;5,p113; 9,p71;11,p245;17; 26,p17;42,p42;48,p319; 58,p35;59
75 ( )	STOVE 2	coal	cnc,stf	sed	0s	Tvs/W2	IB	G/02E	(7)ip 1,p18;4,p32; 5,p113,map;9;26,p17; 42,p42;48,p319;58,p35; 59

OF.	INAC No.	Deposit(s) Name(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (115)	Status References
76	( )	WADE CREEK (MAPLE)	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/05E	(7)op 5,p115,map;9,p73; 58,p35;59
77	( )	BURWASH CREEK GYPSUM	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/06W	(7)ip 5,p115;9;58,p35;59
78	( )	"T"	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/02W	(7)op 5,p115,map;9;58,p35; 59
79	( )	TONY	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/02W	(7)op 5,p115,map;9;58,p35; 59
80	( )	METALLINE (BULLION CREEK)	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/02E	(7)ip 5,p115,map;9,p73;17; 58,p35;59
81	( )	BOCK'S BROOK	Gy,An	dsc,mas	sed,evp;hyd	uRe	w2	IB	G/02W	(7)op 5,p115,map;9,p73; 58,p35;59
82	( )	WOLLI	Wo	mas	skn	uRc	Kg:W2	IB	F/15W	(7)op 58,p35;59;62
83	( )	BULLION CREEK 1	Au,Pt,Cu	flv,unc	plc	Qs	/Ax	IB	G/02E	(4)ip 1,p13;4,p14,p19; 5,p105;7,p152;8,p141; 9,p47;10,p46;17; 19,p17;20,p21;21,p62;
										22,p87;27,p113;28,p65; 38,p139;42,p40;47,p23; 50,p4;51,p6;52,p5; 53,p9;54,p17;58,p35; 59;64
84	( )	TATAMAGOUCHE CREEK	Au,Pt,Cu	flv,unc	plc	Qs	W2	IB	G/06W	(4)op <sup>1</sup> 4,p22;8,p105;9,p52; 18,p184;27,p112; 29,p120;47,p23;58,p35; 59;64;65,p87
85	( )	ARCH CREEK	Au	flv,unc	plc	Qs	W2	IB	G/05E	(1)op 4,p10;5,p107;8,p143; 9,p46;17;19,p18; 20,p17;47,p23;54,p18; 55,p360;58,p35;59;64; 65,p87;66,p69;67,p42
86	( )	DUKE RIVER	Au	flv,unc	plc	Qs	/JK&W2;/W2	IB	G/06E	(1)op <sup>1</sup> 8,p14;1;9,p49;23,p81; 58,p35;59;63

OF.	INAC No.	Deposit(s) Name(s) (No)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (115)	Status References
87	( )	BURWASH CREEK	Au, Pt, Ag, Cu, Sn, Fe	flv, unc	plc	Qs	NW2; Tvs/W2	IB	G/06W (1)op <sup>1</sup> 1,p15;4,p14;5,p105; 7,p152;8,p143;9,p48; 10,p49;17;18,p184; 19,p16;20,p20;21,p63; 22,p86;23,p80;24,p82; 29,p120;27,p112; 28,p65;38,p138;42,p41; 47,p23;51,p6;58,p35; 59,p63;64;65,p87; 66,p66;67,p40
88	( )	QUILL CREEK	Au	flv, unc	plc	Qs	NW2	IB	G/06W (4)op 9;58,p35;59;65,p87; 66,p70;67,p44
89	( )	PORCUPINE CREEK	Au	flv, unc	plc	Qs	NW2	IB	G/05E (4)op 9;58,p35;59;66,p69;67
90	( )	PTARMIGAN CREEK	Au	flv, unc	plc	Qs	NW2; Tvs/W2?	IB	G/06E (7)op <sup>1</sup> 9;58,p35;59;65,p90; 66,p65
91	( )	GRANITE CREEK	Au	flv, unc	plc	Qs	Kg;W2	IB	G/06W (7)op <sup>1</sup> 9;58,p35;59;60,p1; 65,p90;66,p65
92	( )	KLETSAN CREEK	Cu	flv, unc	plc	Qs	NW2; Tvs/W2?	IB	F/10W (7)op 2,p51;3,p133;5,p108; 10,p46;11,p246;17; 28,p42;47,p25;56,p143; 57,p379;58,p35;59
93	( )	SHEEP CREEK 1	Au	flv, unc	plc	Qs	NW2; Tvs/W2	IB	G/02E (4)ip 1,p15;4,p21;5,p106; 9,p50;17;19,p18; 42,p41;58,p35;59;64
94	( )	WHITE RIVER	Cu	flv, unc	plc	Qs	NW2	IB	F/15W (7)op 2,p53;3,p139;5,p109; 10,p46;17;58,p35;59
95	( )	PAN CREEK 1	Au	flv, unc	plc	Qs	NW2; JKs;W2	IB	F/15W (7)op 3,p124;5,p107;58,p35; 59;64
96	( )	PAN CREEK 2	Au	flv, unc	plc	Qs	NW2; JKs;W2	IB	F/15W (7)op 3,p124;5,p107;58,p35; 59;64
97	( )	BOWEN CREEK	Au	flv, unc	plc	Qs	Kg;W2;W2	IB	F/15W (7)op 3,p124;5,p107;58,p35; 59;60,p1
98	( )	HIDDEN CREEK	Au	flv, unc	plc	Qs	Kg;W2;W2	IB	F/15W (7)op 3,p124;5,p107;58,p35; 59;60,p1

OF.	INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt	NTS Belt (115)	Status References
99	( )	EDITH CREEK (KOIDERN RIVER)	Au	flv,unc	plc	Qs	/W2;Kg:W2	IB	F/09E (7)op	3,p124;5,p107;42,p40; 58,p35;59;60,p1
100	( )	HOGE CREEK 1	Au,Pt	flv,unc	plc	Qs	/W2;AX	IB	G/05E (7)ip	8,p141;9,p49;17; 58,p35;59;67,p40
101	( )	HOGE CREEK 2	Au	flv	ppl	0s	Tws/W2	IB	G/05E (7)ip	5,p108;8,p141;9,p49; 17;58,p35;59
102	( )	SQUIRREL CREEK	Au	flv,unc	plc	Qs	/JKs/W2;W2	IB	G/06E (4)op <sup>1</sup>	9;58,p35;59;65,p89; 66,p66;67
103	( )	REED CREEK	Au	flv,unc	plc	Qs	/W2	IB	G/12E (1)op	9;58,p35;59;61,p9;63; 65,p86;66,p70;67,p43
104	( )	BADLANDS CREEK	Au	flv,unc	plc	Qs	/W2	IB	G/06E (7)op <sup>1</sup>	9;58,p35;59;65,p90
105	( )	SWEDIE JOHNSON CREEK	Au,Pt	flv,unc	plc	Qs	/W2	IB	G/12E (1)op	9;58,p35;59;63;65,p86; 66,p65;67
106	( )	JODY CREEK	Au	flv,unc	plc	Qs	/W2	IB	G/12E (1)op	9;58,p35;59;63;65,p86; 66,p65
107	( )	WADE CREEK	Au	flv,unc	plc	Qs	/W2;Tvs/W2	IB	G/05E (4)op	9;58,p35;59;66,p65; 67,p43
108	( )	QUILL, NICKEL CREEKS	Au	flv,unc	plc	Qs	/W2	IB	G/06W (7)op	9;58,p35;59;66,p65; 67,p40
109	( )	MAPLE CREEK	Au	flv,unc	plc	Qs	/W2	IB	G/05E (7)op	9;58,p35;59;66,p65
110	( )	WOLVERINE CREEK	Au	flv,unc	plc	Qs	/W2;Tvs/W2	IB	G/12W (7)op	58,p35;59;64;65,p86
111	( )	LYNX CREEK	Au	flv,unc	plc	Qs	/W2	IB	G/12W (7)op	58,p35;59;65,p86; 66,p65
112	( )	HALFBREED CREEK	Au	flv,unc	plc	Qs	/JKs/W2;W2	IB	G/06E (7)op <sup>1</sup>	9;58,p35;59;65,p86
113	( )	LEWIS CREEK	Au	flv,unc	plc	Qs	/W2;JKs/W2	IB	G/02W (7)op <sup>1</sup>	9;58,p35;59;65,p86

### Explanation of Mineral Deposit Information for SW Kluane Lake (115G / F (E1/2)) map area

- ⊗ 4 mineral deposit(s) locality and map reference number (on map); location accuracy may vary.  
 N.B. locations of placers (plc) mostly are very approximate and are "spot placed" only to indicate creeks or sections of creeks where placers have been mined or occur.

OF. No. map reference number.

INAC (No) INAC (Indian and Northern Affairs Canada) mineral occurrence file number.  
 Deposit(s) Name(s) name (alternate name) of deposit(s). N.B. for placers (plc) only; name is creek or river where deposit(s) occur.

**Commodity(s) Abbreviations** elements present are listed in decreasing order of abundance; elements present in trace amounts are included, but not specified.

Ab	asbestos	Au	gold	Gy	gypsum	Pt	platinum
Ag	silver	Co	cobalt	Mo	molybdenum	PGE	platinum gp. elements
An	anhydrite	Cu	copper	Ni	nickel	Sn	tin
As	arsenic	Fe	iron	Pb	lead	Wo	wollastonite

**Deposit(s) Form Abbreviations**

cnc	concordant	stf	stratiform	mas	massive	bx	breccia
disc	discordant	stg	stringer	dis	disseminated	flv	fluvial deposit
sbt	stratabound	stw	stockwork	vn	vein	unc	unconsolidated

**Deposit(s) Type Abbreviations**

hyd	hydrothermal	skn	skarn	ppp	porphyry	sed	sedimentary
rpl	replacement	vlc	volcanic	evp	evaporite	ppl	paleoplacer
mgm	magmatic					plc	placer

Host Unit(s) 1:250 000 scale map unit(s) host to mineral deposit(s).

Plut.:Cov./Terrane

(Fig. 2 map unit(s) host to mineral deposit(s). Plut. = Plutonic Rocks; : = Intruded into: Cov. = Cover Rocks; / = deposited on: Terrane = Terrane.  
 e.g. Mg? Ogr? PPg AX = Mg? or Ogr? = intruded into PPg; = intruded into AX.

N.B. for placer deposits and float only; e.g. W2/Kg/W2 = Qs (not depicted in Fig. 2) / = deposited on W2 and on Kg : = intruded into W2.

Mg = Miocene granitoids and or subvolcanics: Ogr = Oligocene subvolcanics: PPg = Late Early Cretaceous granitoids: PPg = Late Pennsylvanian-Early Permian granitoids.  
 Tvs = Tertiary volcanics and or subvolcanics: JKs = Jura-Cretaceous flysch (Gravina-Nutzotin Belt); W2 = Wrangellia Terrane, segment W2: AX = Alexander Terrane.

Mgeo Belt Morphogeological Belts (Fig. 2 inset) in which mineral deposit(s) occur (IB = Insular Belt).

NTS National Topographic System map sheet reference.  
 (115) = 1:1 000 000 scale; G / F = 1:250 000 scale; 01 to 16 = 1:50 000 scale; E = East, or W = West half.

#### Status Abbreviations

- (1) Being produced: commodity(s) is (are) being extracted for sale.
- (2) Never produced: reserves (grade + 3 dimens.) or resources known.
- (3) Past producer : reserves (grade + 3 dimens.) or resources known.
- (4) Past producer : reserves or resources exhausted, or not known.
- (5) Grade, 2 dimensions (length and width) known: reserves unknown.
- (6) Grade, 1 dimension (length or width) known : by drill or chip sample.
- (7) Outcrop showing; or commodity(s) known, but unpublished: ± grab sample assays.
- (8) Float; provenance known or unknown: ± grab sample assays.

ip = inside, or op = outside of Kluane National Park Reserve: op<sup>1</sup> = op, but on lands withdrawn from staking due to Native Land Claims.

**References** (e.g. 9,p50 = Read and Monger (1976) page 50; NB. p = initial page of reference cited; (1976) = year of publication.

N.B. references listed are main sources only; for references such as INAC publications (not given), individual assessment report numbers and summaries, and claims information, the reader should see INAC (1989) or literature cited there.

- 1 McConnell (1905).
- 2 Moffit and Knopf (1910).
- 3 Cairnes (1915a).
- 4 Cairnes (1915b).
- 5 Muller (1967).
- 6 KirKham (1971).
- 7 Craig and Laporte (1972).
- 8 Sinclair and Gilbert (1975).
- 9 Read and Monger (1976).
- 10 Carrier et al. (1981).
- 11 INAC, (1985).
- 12 INAC, (1986).
- 13 INAC, (1987).
- 14 INAC, (1988).
- 15 INAC, (1989).
- 16 Morin (1989).
- 17 Canmindex, 115G/F (1989).
- 18 Sinclair et al. (1975).
- 19 Skinner (1961).
- 20 Skinner (1962).
- 21 Green and Goodwin (1963).
- 22 Findlay (1967).
- 23 Green (1965).
- 24 Green and Goodwin (1964).
- 25 Cameron & Birmingham (1970).
- 26 Campbell, J.D. (1967).
- 27 Findlay (1968a).
- 28 Findlay (1969b).
- 29 Green (1966).
- 30 Sinclair et al. (1976).
- 31 Sinclair et al. (1979).
- 32 Christopher et al. (1972).
- 33 Mähnyk (1972).
- 34 Campbell, S.W. (1981).
- 35 Campbell, F.A. (1956).
- 36 Campbell, F.A. (1960).
- 37 Campbell, S.W. (1976).
- 38 Craig and Milner (1975).
- 39 Campbell, S.W. (1977).
- 40 MacLean (1914).
- 41 Sharp (1943).
- 42 Bostock (1952).
- 43 Morin et al. (1977).
- 44 Evenchick et al. (1987).
- 45 Hulbert et al. (1988).
- 46 Johnston (1915).
- 47 McConnell (1906).
- 48 Eisbacher & Hopkins (1977).
- 49 Ridgway et al. (1989).
- 50 Bostock (1935).
- 51 Bostock (1936).
- 52 Bostock (1937).
- 53 Bostock (1938).
- 54 Bostock (1941).
- 55 Bostock (1957).
- 56 Hayes (1892).
- 57 Brooks (1900).
- 58 Campbell and Dodds (1978).
- 59 Campbell and Dodds (1982a).
- 60 Dodds and Campbell (1988).
- 61 INAC Overview, (1990).
- 62 Archer Cathro L. P. com. (1990).
- 63 T. Bremner, P. com. (1990).
- 64 Debicki (1982).
- 65 Debicki (1983).
- 66 Debicki and Gilbert (1986).
- 67 LeBarge and Morison (1990).

Table 3. MINERAL DEPOSITS MOUNT ST. ELIAS MAP AREA (115B AND C [E1/2])

OF. INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt	NTS (115)	Status References
1 (2)	KASKAWULSH	Cu,Ag	unk	unk	uPbv?	W2?;AX?	IB	B/09E (7)ip	11,p162;17,p35;18
2 (11)	JENNIFER 1 (KINNORA)	Zn	dis?	skn	uPvs?	Og?: (W2?;AX?)	IB	B/16W (7)op	9,p310;17,p35;18; 19,p122,p10
3 (11)	JENNIFER 2 (KINNORA)	Cu,Ag	stw,vn	hyd,rpl	Dc?	AX	IB	B/16W (7)op	9,p310;17,p35;18; 22,p10
4 (6)	GIBBONS (VULCAN)	Ni,Cu	unk	mgn	Pfub,b	W2?;AX?	IB	B/15E (7)ip	4;7,p55;11,p162;12; 17,p35;18
5 (7)	FROHBERG (ULTRA)	Cu,Ni,PGE,Zn	stw,vn,dis	mgn;hyd,rpl	Pfub,ub, uPp	W2?	IB	B/16W (6)op	10,p230;13;17,p35;18; 23
6 (7)	TELLURIDE	Zn,Cu,Pb,Ag	stb,mas,dis	vlc,exh?	uPbv	W2?;AX?	IB	B/16W (7)op	10,p229;11,p162;12;13; 17,p35;18;23
7 (7)	CUB (TELLURIDE)	Zn,Cu,Pb,Ag	ftt,(stb)	(vlc,exh?)	uPbv?	/W2	IB	B/16E (8)op	4;7,p67;8,p61;10,p229; 12;13;17,p35;18;23
8 (1)	PLUG	Cu,Ag	ftt,(unk)	(unk)	?	/AX	IB	B/01E (8)ip	11,p162;17,p35;18;23
9 (5)	DULUTH	Ni,Cu	ftt,(unk)	(mgn?)	uPbv?	/W2?;AX?	IB	B/15E (8)ip	4;11,p162;17,p35;18
10 (8)	BULLION	Gy,An	dsc,mas	sed,evp;hyd	uRe	W2	IB	B/15E (7)ip	5,p115;7,p73;11,p162; 17,p35;18
11 (3)	KIMBERLEY CREEK COAL	coal	cnc,stff	sed	Os	Tvs/JKs/W3	IB	B/16E (7)op <sup>1</sup>	1,p118;2,p32;3,p58; 6,p20;7,p70;11,p162; 12;17,p35;18;21,p17
12 (-)	TELLURIDE CREEK COAL	coal	cnc,stff	sed	Os	Tvs/W3	IB	B/16E (7)op	2,p32;7,p72;11,p162; 12;17,p35;18
13 (-)	BULLION CREEK 2	Au,Ag,Pt,Cu,Pb	flv,unc	plc	Qs	/AX	IB	B/15E (4)ip	1,p13;2,p14;5,p105; 7,p47;8,p46;12;14,p17; 15,p113;16,p52; 17,p35;18;20,p65;24
14 (9)	SHEEP CREEK 2	Cu	flv,unc	plc	Qs	/W2	IB	B/15E (7)ip	1,p17;7;17,p35;18
15 (-)	TELLURIDE CREEK	Au	flv,unc	plc	Qs	/W3	IB	B/16E (7)op	1,p16;7;17,p35;18; 25,p85
16 (-)	VULCAN CREEK	Au	flv,unc	plc	Qs	/W2	IB	B/15E (7)ip	1,map 894;7;17,p35;18

No.	OF. No.	INAC Name(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt (115)	Status	References
17	( )	KIMBERLEY CREEK 2	Au	fly,unc	plc	Qs	/JKs?W3	IB	B/16E (4)op <sup>1</sup>	1,p16;3,p55;4;7,p50;12, 17,p35;18;25,p85
18	(4)	JARVIS	Cu	fly,unc	plc	Qs	/JKs?W3	IB	B/16E (7)op <sup>1</sup>	1,p17;7;17,p35;18
19	( )	CANADA CREEK	Au	fly,unc	plc	Qs	/AX	IB	B/15W (7)ip	1,p1;17,p35;18

## **Explanation of Mineral Deposit Information for Mount St. Elias (115B / C [E1/2]) map area**

4 mineral deposit(s) locality and map reference number (on map): location accuracy may vary.  
N.B. locations of placers (plc) mostly are very approximate and are "spot placed" only to indicate creeks or sections of creeks where placers have been mined or occur.

OE Name main reference number

卷之三

**Deposit(s) Name(s)** name (alternate name) of deposit(s). N.B. for placers (plc) only: name is creek or river where deposit(s) occur.  
**INAC (No)** INAC (Indian and Northern Affairs Canada), mineral occurrence file number.

Abbreviations: Abbreviations are listed in descending order of abundance: elements present in trace amounts are included but not numbered.

Deposit(s) Form Abbreviations		Deposit(s) Type Abbreviations	
Ag	silver	Au	gold
An	anhydrite	Cu	copper
cnc	concordant	stb	stratobound
dsc	discordant	stf	stratiform
gy	gypsum	stw	stockwork
Ni	nickel	mas	massive
Pb	lead	dis	disseminated
Pt	platinum	vn	vein
PGE	platinum group elements	flv	fluvial deposit
Zn	zinc	unc	unconsolidated
fit			
unk			
unknown			
evp	evaporite		
plc	placer		
exhalative			
sedimentary			
magmatic			
volcanic			
hyd	hydrothermal		
rpl	replacement		

Deposit(s) Form Abbreviations

Host | Infill(s) | 1:250 000 scale map unit(s) host to mineral deposit(s)

## Plut.:Cov./Terrane

(Fig. 2 map unit(s) host to mineral deposit(s). Plut. = Plutonic Rocks; : = intruded into: Cov. = Cover Rocks; / = deposited on: Terrane = Terrane.  
 e.g. Tvs/JKs/W3 = Tvs / = deposited on JKs / = which was deposited on W3.  
 N.B. for placer deposits and float only; e.g. JKs?/W3 = Qs (not depicted in Fig. 2) / = deposited on JKs? / = which was deposited on W3.

Og = Oligocene subvolcanics: Tvs = Tertiary sediments: JKs = Jura-Cretaceous flysch (Gravina-Nutzotin Belt): W2,W3 = Wrangellia Terrane, segments 2, 3:  
 AX = Alexander Terrane.

**Mgeo Belt** Morphogeological Belts (Fig. 2 inset) in which mineral deposit(s) occur (IB = Insular Belt).

**NTS** National Topographic System map sheet reference.  
 (115) = 1:1 000 000 scale: B / C = 1:250 000 scale: 01 to 16 = 1:50 000 scale; E = East, or W = West half.

## Status Abbreviations

- (1) Being produced: commodity(s) is (are) being extracted for sale.
- (2) Never produced: reserves (grade + 3 dimens.) or resources known.
- (3) Past producer : reserves (grade + 3 dimens.) or resources known.
- (4) Past producer : reserves or resources exhausted, or not known.
- (5) Grade, 2 dimensions (length and width) known: reserves unknown.
- (6) Grade, 1 dimension (length or width) known : by drill or chip sample.
- (7) Outcrop showing; or commodity(s) known, but unpublished: ± grab sample assays.
- (8) Float; provenance known or unknown: ± grab sample assays.

ip = inside, or op = outside of Kluane National Park Reserve: op<sup>1</sup> = op, but on lands withdrawn from staking due to Native Land Claims.

**References** (e.g. 7,p50 = Read and Monger (1976) page 50; NB. p = initial page of reference cited; (1976) = year of publication.  
 N.B. references listed are main sources only; for references such as INAC publications (not given), individual assessment report numbers and summaries, and claims information, the reader should see INAC (1989) or literature cited there.

- 1 McConnell (1905).
- 2 Cairnes (1915b).
- 3 Kindle (1953).
- 4 Wheeler (1963).
- 5 Muller (1967).
- 6 Cameron and Birmingham (1970).
- 7 Read and Monger (1976).
- 8 Carriere et al. (1981).
- 9 INAC, (1987).
- 10 INAC, (1988).
- 11 INAC, (1989).
- 12 Canmindex, 115B (1989).
- 13 Morin (1989).
- 14 Skinner, (1961).
- 15 Findlay, (1969a).
- 16 Craig and Laporte (1972).
- 17 Campbell and Dodds (1978).
- 18 Campbell and Dodds (1982b).
- 19 Dodds and Campbell (1988).
- 20 Findlay (1969b).
- 21 Campbell (1967).
- 22 INAC Overview, (1990).
- 23 Archer Cathro Ltd. p. com.(1990).
- 24 Debicki (1982).
- 25 Debicki (1983).

Table 4. MINERAL DEPOSITS SW DEZADEASH MAP AREA (115A)

OF. INAC No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s)	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (115)	Status References
1 (1)	JACKPOT (PIRATE CREEK)	Cu,Ag,Zn,Pb	vn,bx,mas,dis	hyd,rpl	PRv	W2	IB	A/03E	(6)op	7,p108;8,p72;9,p62; 10,p44;11,p24;117; 27,p43;43,p35;44;45,p1
2 (3)	MOHAWK (KANE)	Ag,Pb,Zn,Au,Sb,Cu	stw,vn,bx	hyd,rpl	Φ?;Pv	Φg?;W2	IB	A/03E	(4)op <sup>1</sup>	9,p65;10,p60;12,p166; 16;17;18,p140;43,p35; 44;45,p147
3 (6)	SNO (MUSH)	Cu,Au	vn,bx	hyd,rpl	Pv	W2	IB	A/03W	(7)ip	3,p56;9,p64;10,p45;17; 19,p37;43,p35;44
4 (7)	BATES (IRON CREEK)	Pb,Ag,Au	vn	hyd,rpl	Ev?	AX	IB	A/04E	(6)ip	3,p56;9,p65;15,p159; 16;17;43,p35;44
5 (8)	FENTON	Cu	vn	hyd?,rpl?	Ev	AX	IB	A/04W	(7)ip	9;15,p159;17;43,p35; 44;47
6 (9)	SANDY (CAVE)	Cu,Ag	stb,dis,vn	vlc?;hyd,rpl	Pv?	W2	IB	A/06W	(7)ip	9,p63;10,p53;15,p159; 17;43,p35;44
7 (10)	SHAFT	Cu	dis	unk	Psv	W2	IB	A/05E	(6)ip	9;15,p159;17;43,p35; 44;47
8 (12)	HUSKY	Cu	stb,dis,vn	vlc?;hyd?,rpl?	uTN	W2	IB	A/06W	(7)ip	9,p59;10,p56;15,p159; 17;43,p35;44
9 (13)	WREN	Cu	vn	unk	uTN	W2	IB	A/06W	(7)ip	9;15,p159;43,p35;44;47
10 (14)	KEL	Cu	stb,dis,vn	vlc?;hyd?,rpl?	uTN	W2	IB	A/06W	(7)ip	9,p60;10,p45;15,p159; 17;43,p35;44
11 (20)	JOHOBBO (YUKON STAR)	Cu,Ag	mas,dis,vn	vlc;hyd,rpl	uTN	W2	IB	A/05E	(3)ip	6,p85;7,p10;9,p59; 10,p45;15,p159;17; 19,p28;20,p27;21,p24; 22,p55;23,p35;24,p1; 25,p29;43,p35;44
12 (20)	BORNITE CREEK (KINDLE)	Cu,Ag	dis,vn	vlc;hyd,rpl	uTN	W2	IB	A/05E	(7)ip	3,p57;9,p59;15,p159; 43,p35;44
13 (26)	DECOELI	Cu,Ab	vn	hyd,rpl	PMvs	W3	IB	A/13W	(7)op	15,p159;17;43,p35;44; 47

OF. No.	INAC (No.)	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (115)	NTS Status	References
14 (27) KLOO (ELLEN)	Cu,Au	stf?;mas	vlc?;exh?	PMvs	W3	IB	A/13W (5)op	10,p46;15,p159;17; 22,p54;43,p35;44;46,p7		
15 (28) SOUTHER-STANCUI	Cu,Mo,Pb,Zn,Ag,Au	stw,bx,vn	"b.p.";Nw?	Mg:(Tvs,AX)	IB	A/12W (7)ip	10,p59;15,p159;17; 26,p66;43,p35;44;45,p1			
16 (5) MIKE (PHOTO)	Cu,Co,Au,Sb	vn,bx	hyd,rpl	Pv	W2	IB	A/03W (6)op <sup>1</sup>	4,p73;9,p62;10,p45; 15,p159;17;43,p35;44		
17 ( ) BATES LAKE (ISLAND)	Ab	vn	hyd,rpl	Kg?;Ob?	Kg?;AX;AX?	IB	A/05E (7)ip	3,p38;9,p67;15,p159; 17;43,p35;44;45,p1		
18 (23) STRIDE	Cr,Fe,Pt,Ti	stf;pod,dis	mgm	I Kub,b?	Kg;Jks/W3	IB	A/12W (7)ip	3,p37;15,p159;17; 28,p129,p185;41; 43,p35;44;45,p1		
19 (2) DALTON	Cu	bx,vn,dis	prp?;hyd?;rpl?	Or?;Pv?	Og?;W2	IB	A/03E (6)op	9;15,p159;43,p35;44; 45,p1		
20 (35) BURGER KING	Au,Ag	vn	hyd?;rpl?	Or?;?;UPp?	Og?;?;AX;AX	IB	A/03E (7)op	12,p169;13,p306;16; 43,p35;44;45,p1		
21 (25) FERGUSON	Au	vn	hyd,rpl?	? AX	AX	IB	A/12W (7)ip	9;15,p159;43,p35; 30,p12;31,p11;47		
22 ( ) TATSHENSHINI	Cu,Mo	dis	prp	Kg?	Kg?;W2	IB	A/03E (7)op <sup>1</sup>	9;43,p35;44;45,p1;47		
23 (11) BELOUD	Cu	vn	hyd?;rpl?	Pv	W2	IB	A/06W (7)ip	9;15,p159;43,p35;44		
24 (15) SHORTY	Fe	vn	hyd,rpl	Kgd	Kg;Jks/W3	IB	A/06E (7)ip	3,p49;15,p159; 43,p35;44		
25 ( ) COLTON 1	Cu,Au	vn	hyd?;rpl?	PMvs	W3	IB	A/13W (7)op	15,p159;43,p35;44; 46,p10		
26 ( ) COLTON 2	Cu	ft,(unk)	(unk)	PMvs?	W3	IB	A/13W (8)op	15,p159;43,p35;44		
27 (19) MILLHOUSE	Cu,Ag,Au	ft,(unk)	(unk)	? /JKs/W3	IB	A/11W (8)ip	15,p159;43,p35;44;47			
28 (24) SUGDEN	coal	cnc,eff	sed	Os	Tvs/(W2,AX?)	IB	A/12W (7)ip	2,p17;3,p58;5,p70; 9,p72;15,p195;17; 42,pA50;43,p35;44		
29 ( ) SUGDEN CREEK	Au,Pt	flv,unc	p/c	Qs	/Tvs/AX;AX	IB	A/12W (4)ip	3,p54;9,p52;17;43,p35; 44;48		
30 ( ) CONTACT CREEK	Pt	flv,unc	p/c	Qs	/Kg;Jks/W3	IB	A/12W (7)ip	3,p37;41;43,p35;44; 45,p1		

OF.	INAC No.	Deposit(s) Name(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt (115)	Status References
31	( )	GOAT CREEK	Au	flv,unc	plc	Qs	/Jks/W3	IB	A/11W (4)ip 48
32	( )	VICTORIA CREEK	Au	flv,unc	plc	Qs	/Og?;Jks/W3	IB	A/11W (4)ip 3,p48;43,p35;44;45,p1
33	( )	DOLLIS (SQUAW) CREEK	Au,Cu,Ag,Pb	flv,unc	plc	Qs	/W2;AX	IB	A/03E (1)op 3,p50;9,p51;17;30,p6; 31,p5;32,p6;33,p4; 34,p10;35,p18;36,p7; 37,p2;38,p74;39,p90; 40,p36;43,p35;44; 49,p83;50
34	( )	BATES RIVER 1	Au	flv,unc	plc	Qs	/AX	IB	A/04E (7)ip 3,map;9;43,p35;44
35	( )	MUSH CREEK	Au	flv,unc	plc	Qs	/W2	IB	A/06W (7)ip 3,p48;9;43,p35;44
36	( )	SHAFT CREEK	Au	flv,unc	plc	Qs	/Tvs?;AX	IB	A/05E (7)ip 3,map;9;43,p35;44
37	( )	GOATHERD CREEK	Au	flv,unc	plc	Qs	/AX	IB	A/05W (7)ip 3,p54;9;43,p35;44
38	( )	MARBLE CREEK	Au	flv,unc	plc	Qs	/AX	IB	A/05W (7)ip 3,p54;9;43,p35;44
39	( )	PARK CREEK	Au	flv,unc	plc	Qs	/Mg;(Tvs,AX)	IB	A/12W (7)ip 3,map;9;43,p35;44; 45,p1
40	( )	ARCHIBALD CREEK	Au	flv,unc	plc	Qs	/Jks/W3;W3	IB	A/13W (7)op 3,map;43,p35;44;45,p1
41	( )	KIMBERLEY CREEK 1	Au	flv,unc	plc	Qs	/Jks/W3;W2	IB	A/13W (4)op <sup>1</sup> 1,p16;3,p55;9,p50; 43,p35;44;49,p83;50
42	( )	WOLVERINE CREEK	Au	flv,unc	plc	Qs	/AX	IB	A/04E (4)ip 3,p48;9;43,p35;44
43	( )	TATSHENSHINI RIVER	Au	flv,unc	plc	Qs	/W2	IB	A/03E (7)op 3,p52;9,p53;43,p35;44
44	( )	SILVER CREEK	Au	flv,unc	plc	Qs	/AX;W2	IB	A/03E (7)ip 3,p50;9,p51;17;43,p35; 44
45	( )	IRON CREEK	Au	flv,unc	plc	Qs	/AX	IB	A/04E (7)ip 3,p53;9,p50;17;30,p6; 31,p5;43,p35;44
46	( )	BATES RIVER 2	Au	flv,unc	plc	Qs	/AX	IB	A/04E (4)ip 3,p53;9,p47;17;19,p19; 20,p22;43,p35;44
47	( )	BELOUD CREEK	Au,Cu	flv,unc	plc	Qs	/W2	IB	A/06W (4)ip 3,p49;9,p47;10,p45;17; 43,p35;44
48	( )	ALDER CREEK	Au	flv,unc	plc	Qs	/Jks/W3	IB	A/06E (7)ip 3,map;43,p35;44

OF. No.	INAC Deposit(s) Name(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (115)	Status	References
49 ( )	SHORTY CREEK	Au,Cu	flv,unc	plc	Qs	/JKsW3	IB	A/06E	(4)ip	3,p49;10,p45;17; 43,p35;44,p8
50 ( )	DALTON POST (TATSHENSHINI RIVER)	Au	flv,unc	plc	Qs	/W2	IB	A/03E	(7)op	3,p52;9,p53;43,p35;44

#### Explanation of Mineral Deposit Information for SW Dezadeash (115A) map area

④ mineral deposit(s) locality and map reference number (on map): location accuracy may vary.  
 N.B. locations of placers (plc) mostly are very approximate and are "spot placed" only to indicate creeks or sections of creeks where placers have been mined or occur.

##### OF. No. map reference number.

INAC (No) INAC (Indian and Northern Affairs Canada) mineral occurrence file number.

Deposit(s), Name(s) name (alternate name) of deposit(s). N.B. for placers (plc) only; name is creek or river where deposit(s) occur.

Commodity(s) Abbreviations elements present are listed in decreasing order of abundance; elements present in trace amounts are included, but not specified.

Ab	asbestos	Co	cobalt	Fe	iron	Pt	platinum	Zn	zinc
Ag	silver	Cr	chromium	Mo	molybdenum	Sb	antimony		
Au	gold	Cu	copper	Pb	lead	Ti	titanium		

##### Deposit(s) Form Abbreviations

cnc	concordant	stf	stratiform	mas	massive	bx	breccia	flv	fluvial deposit
disc	discordant	stw	stockwork	dis	disseminated	vn	vein	unc	unconsolidated
stb	stratabound	pod	podiform						

##### Deposit(s) Type Abbreviations

hyd	hydrothermal	mgm	magmatic	exh	exhalative	sed	sedimentary	unk	unknown
rpl	replacement	vc	volcanic	pp	porphyry	pic	placer		

Host Unit(s) 1:250 000 scale map unit(s) host to mineral deposit(s).

## Plut.:Cov./Terrane

(Fig. 2 map unit(s) host to mineral deposit(s). Plut. = Plutonic Rocks; : = intruded into: Cov. = Cover Rocks; / = deposited on: Terrane = Terrane.  
e.g. Mg:(Tvs,AX) = Mg: = intruded into (:) = both Tvs and AX.  
N.B. for placer deposits only; e.g. Kg:JKs/W3 = Qs (not depicted in Fig. 2) / = deposited on Kg := intruded into JKs which was / = deposited on W3.

Mg = Miocene subvolcanics: Og = Oligocene subvolcanics: Kg = Late Cretaceous granitoids or Alaskan-type, ultramafic complexes:  
Tvs = Tertiary volcanics or sediments: JKs = Jura-Cretaceous flysch (Gravina-Nutzotin Belt): W2, W3 = Wrangellia Terrane; segments 2, 3: AX = Alexander Terrane.

**Mgeo Belt** Morphogeological Belts (Fig. 2 inset) in which mineral deposit(s) occur (IB = Insular Belt).

**NTS** National Topographic System map sheet reference.  
(115) = 1:1 000 000 scale: A = 1:250 000 scale: 01 to 16 = 1:50 000 scale; E = East, or W = West half.

## Status Abbreviations

- (1) Being produced; commodity(s) is (are) being extracted for sale.
- (2) Never produced; reserves (grade + 3 dimens.) or resources known.
- (3) Past producer : reserves (grade + 3 dimens.) or resources known.
- (4) Past producer : reserves or resources exhausted, or not known.
- (5) Grade, 2 dimensions (length and width) known; reserves unknown.
- (6) Grade, 1 dimension (length or width) known : by drill or chip sample.
- (7) Outcrop showing; or commodity(s) known, but unpublished; ± grab sample assays.
- (8) Float; provenance known or unknown: ± grab sample assays.

ip = inside, or op = outside of Kluane National Park Reserve: op¹ = op, but on lands withdrawn from staking due to Native Land Claims.

**References** (e.g. 9,p50 = Read and Monger (1976) page 50; NB. p = initial page of reference cited; (1976) = year of publication.

N.B. references listed are main sources only; for references such as INAC publications (not given), individual assessment report numbers and summaries, and claims information, the reader should see INAC (1989) or literature cited there.

- 1 McConnell (1906).
- 2 Campbell, J.D. (1967).
- 3 Kindle (1953).
- 4 Findlay (1969a).
- 5 Cameron and Birmingham (1970).
- 6 Kirkham (1971).
- 7 Craig and Laporte (1972).
- 8 Sinclair and Gilbert (1975).
- 9 Read and Monger (1976).
- 10 Carrierre et al. (1981).
- 11 INAC, (1985).
- 12 INAC, (1986).
- 13 INAC, (1987).
- 14 INAC, (1988).
- 15 INAC, (1989).
- 16 Morin (1989).
- 17 Canmindex, 115A (1989).
- 18 Sinclair et al. (1975).
- 19 Skinner (1961).
- 20 Skinner (1962).
- 21 Green and Godwin (1963).
- 22 Findlay (1967).
- 23 Green (1965).
- 24 Wamock (1962).
- 25 Green and Godwin (1964).
- 26 Souther & Stancui (1975).
- 27 Findlay (1969b).
- 28 Sturrock (1975).
- 29 Sturrock et al. (1980).
- 30 Bostock (1936).
- 31 Bostock (1937).
- 32 Bostock (1933).
- 33 Bostock (1934).
- 34 Bostock (1939).
- 35 Bostock (1941).
- 36 Cockfield (1928).
- 37 Cockfield (1930).
- 38 Mandy (1933).
- 39 Mandy (1934).
- 40 Watson (1948).
- 41 Evenchick et al. (1987).
- 42 Ridgway et al. (1989).
- 43 Campbell and Dodds (1978).
- 44 Campbell and Dodds (1982c).
- 45 Dodds and Campbell (1988).
- 46 INAC Overview, (1990).
- 47 Archer Cathro Ltd. p.com.(1990).
- 48 Dabicki (1982).
- 49 Dabicki (1983).
- 50 LeBarge and Morison (1990).

Table 5. MINERAL DEPOSITS TATSHENSHINI RIVER (114P) AND NE YAKUTAT (114O)<sup>1</sup>

OF.	MINF No. (No)	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (114)	Status References
1	(1)	DUCK'S FACE	Gy	unk	unk	uPv, uPpc?	AX	IB	P/13E (7)op 7,8,p17;9,13,p173; 23,p70;24,p1
2	(2)	WINDY CRAGGY	Cu,Co,Au,Ag	stf,mas,stw,stg	vlc,exh	uPv, uPpc	AX	IB	P/12E (2)op 8,p17;9,11,p159; 13,p173;14,p131; 16,p109;17,p195; 18,p197;19,p120,p723; 23,p65;24,p2;25,p455; 33,p3
3	(3)	TATS	Cu,Ag,Co,Zn,Au	stb,mas	vlc,exh	uPv	AX	IB	P/12E (6)op 8,p17;9,11,p159; 13,p173;14,p135; 16,p109;23,p65;24,p5
4	(4)	ALSEK RIVER (TURNBACK)	Cu	unk	unk	Psp?	AX	IB	P/13W (7)op 8,p17;9
5	(5)	O'CONNOR RIVER (SNOW)	Gy,An	dsc,mas	sed,avp;hyd	uPe	AX	IB	P/10E (2)op 8,p17;9,21,p279; 23,p65;24,p9
6	(6)	KIM (AUNT JEMIMA)	Zn	dis,vn	skn?;hyd,rpl	Dc?	Qg?;AX	IB	P/10E (7)op 8,p17;9,22,p1;23,p65; 24,p11
7	(7)	MAID OF ERIN (L722)	Cu,Ag,Au,Zn,Bi,Fe	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/10E (4)op 1,p29;3,p40;8,p17;9; 22,p123,p65;24,p12; 29,p117;30,p237;32
8	(8)	STATE OF MONTANA (L283)	Cu,Ag,Au,Bi	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/10E (4)op 1,p29;3,p40;5,p14; 8,p17;9,22,p1;23,p65; 24,p14;29,p117; 30,p237;32
9	(9)	VICTORIA (L903)	Zn,Pb,Ag,Cu,Au	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/10E (6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;24,p16;32
10	(10)	ADAMS (L286)	Pb,Zn,Ag,Cu,Mo	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/10E (6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;24,p18;32
11	(11)	LAWRENCE (L955)	Zn,Pb,Ag,Cu,Au	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/09W (6)op 1,p29;3,p40;8,p17;9; 22,p124,p20;32
12	(12)	SIMCOE (THREE GUARDSMEN)	Cu,Fe,Au,Mo	pod,mas,dis	skn;rpl	uPs	QgAX	IB	P/09W (6)op 1,p29;3,p40;8,p17;9; 22,p124,p22;32

OF. No.	MINF No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s) Type	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt (114)	Status References
13 (13)	MILDRED (THREE GUARDSMEN)	Cu,Ag,Fe	pod,mas,dis	skn;tp;hyd?	uPs	0g;AX	IB	P/09W (6)op	1,p29;3,p40;8,p17;9; 22,p124,p24;32	
14 (14)	CANADIAN VERDEE	Ag,Cu,Zn,Bi,Au,Fe	pod,mas,dis	skn;tp;hyd?	uPs	0g;AX	IB	P/09W (7)op	1,p29;3,p40;8,p17;9; 22,p124,p26;32	
15 (15)	GOLD CORD (STAMPEDE)	Au,Ag,Cu,Zn	vn,dis	hyd	0g	0g;AX	IB	P/07E (6)op	2,p14;3,p59;8,p17;9; 17,p191;22,p124,p28; 26,p40	
16 ( )	CALLISON	Pb,Cu	unk	unk	?	AX	IB	P/10W (7)op	8,p17;9	
17 (17)	WINDSOR L804	Cu,Zn	dis?	skn?;tp?	uPs	0g;AX	IB	P/10E (7)op	8,p17;9;22,p123,p65; 24,p32;32	
18 (18)	HUM BIRD - DISCOVERY	Ag,Pb,Zn,Cu	vn,bx,dis	hyd;tp!	Pvb	AX	IB	P/10W (7)op	8,p17;9;23,p65;24,p33; 27,p71	
19 (19)	BORNITE (CAT)	Cu,Pb,Zn;Mo	dis;bx	skn;prp?	uPs; 0g, 0f?	0g;AX	IB	P/10E (6)op	8,p17;9;22,p123,p65; 24,p36;32	
20 (20)	WAR EAGLE L901	Fe	dis?	skn?;tp?	uPs	0g;AX	IB	P/10E (7)op	8,p17;9;22,p123,p65; 24,p36;32	
21 (21)	SHEEP	Cu,Zn	unk	unk	PRv?	W2?;AX?	IB	P/14E (6)op	8,p17;9;24,p37;32	
22 (22)	KELSAL 24	Cu,Zn,Mo	vn,dis;lt	hyd	PMvs	W3,TU?	IB?	P/16W (6)op	8,p17;9;24,p38;32	
23 (23)	KELSAL 32	Cu	vn,dis;lt	hyd	PMvs	W3,TU?	IB?	P/16W (6)op	8,p17;9;24,p39	
24 (24)	HUM BIRD - CREEK	Ag,Zn,Pb,Cu	vn,dis	hyd;tp!	Pvb	AX	IB	P/10W (6)op	8,p17;9;23,p65;24,p40; 27,p71	
25 (25)	HUM BIRD - SOUTH	Ag,Cu,Zn,Pb	vn,dis	hyd;tp!	Pvb	AX	IB	P/10W (6)op	8,p17;9;23,p65;24,p42; 27,p71	
26 (26)	HUM BIRD - CAMP	Ag,Cu,Zn,Pb	vn,dis	hyd;tp!	Pvb	AX	IB	P/10W (6)op	8,p17;9;23,p65;24,p44; 27,p71	
27 (27)	LUNAR (MAG)	Ag,Pb,Zn,Fe	dis?;vn	skn;hyd;tp!	uPc?	AX?;0g;AX	IB	P/10E (6)op	8,p17;9;23,p65;24,p46	
28 (28)	NADAHINI MOUNTAIN	Cu,Ab,Fe	vn,dis?	hyd;tp!	Mb, PRub?;b?	AX	IB	P/10E (7)op	3,p25;8,p17;9;23,p65; 24,p47;32	
29 (29)	HIBERNIAN	Cu,Ag,Pb,Zn,Fe	stg,vn,dis	skn;tp!	uPs	0g;AX	IB	P/10E (6)op	1,p29;8,p17;9;23,p65; 24,p48;32	

OF. No.	MINF No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (114)	NTS Belt	Status References
30	(30)	MOUNT MANSFIELD (BEARI)	Cu,Pb,Au	vn,dis,fl	hyd,pl	PMvs	W3,TU?	IB?	P/15E (7)op	3,p59;8,p17;9,24,p50
31	(31)	C AND E NORTH	Ni,Cu,PGE,Zn	dis,vn	mgn;hyd,rl	PTb,ub, PRv	W2	IB	P/15W (7)op	8,p17;9,24,p51;32
32	(32)	C AND E SOUTH	Zn,Sb,Pb,Sr	dis	unk	PRv	W2	IB	P/15W (7)op	8,p17;9,24,p52;32
33	(33)	WC 17 (TATS LAKE)	Ag,Au	unk	unk	ODcs	AX	IB	P/12E (7)op	8,p17;9,24,p53
34	(34)	TATS CREEK (ALSEK)	Cu,Ag,Au	unk	unk	JKg	JKg:AX	IB	P/12E (7)op	8,p17;9,23,p65;24,p54
35	(35)	BARBICAN MOUNT	Ag,Cu,Ag	vn,sg,dis	hyd,mc?	uPpc,v?	AX	IB	P/12E (7)op	8,p17;9,13,p174; 14,p135;24,p56
36	(36)	KLEHINI RIVER SW	Zn,Ag,Cd	vn	hyd	Og	Og:AX	IB	P/10E (7)op	8,p17;9,24,p58
37	(37)	ALSEK TRIBUTARY	Cu,Au,Ag	unk	unk	ODcs?	AX	IB	P/12W (7)op	8,p17;9,13,p174; 14,p135;23,p65;24,p60
38	(38)	FAULT CREEK (TARR)	Pb,Zn,Ag,Au	vn	hyd	ODcs	AX	IB	P/10W (7)op	8,p17;9,24,p61
39	(39)	CAMP CREEK	Cu,Ag,Au	unk	skn;hyd	uPs	Og:AX	IB	P/10E (7)op	8,p17;9,22,p123,p65; 24,p63;29,p117; 30,p237
40	(40)	KLEHINI RIVER NE	Zn,Cu,Pb,Ag	unk	skn;hyd?	uPs	Og:AX	IB	P/10E (7)op	8,p17;9,22,p123,p65; 24,p65;29,p117; 30,p237
41	(41)	ALSEK (TATS GLACIER)	Cu,Ag,Au,Fe	mas,vn	vic?;hyd?	uPv	AX	IB	P/12E (7)op	8,p17;9,13,p173; 14,p135;24,p67;32
42	(42)	INSPECTOR CREEK	Cu,Ag	mas	skn;pl?	uPs	Og:AX	IB	P/10E (7)op	8,p17;9,22,p123,p65; 24,p68;32
43	(43)	BASEMENT - JUNE 24	Ba,Ag,Cu,Pb,Zn,Au	stb,dis	unk	Psp?	AX	IB	P/06W (7)op	8,p17;9,24,p70
44	(44)	BASEMENT - JUNE 21	Au,Cu,Ag	stb,vn,stg,dis	unk	Psp?	AX	IB	P/06W (7)op	8,p17;9,24,p72
45	(45)	BASEMENT MAIN	Cu,Co	mas,pod	vic	Psp?	AX	IB	P/06W (6)op	8,p17;9,24,p74
46	(46)	HUM BIRD - DOME	Ag,Cu,Zn,Pb,Au	vn,dis	hyd,pl	Pvb	AX	IB	P/10W (6)op	8,p17;9,23,p65;24,p76; 27,p71

OF.	MINF No.	Deposit(s) (No)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s) Unit(s)	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (114)	Status References	
47	(47)	SAM - NORTH GLACIER	Cu,Zn,Pb,Au,Ag	pod,dis	skn;hyd?	Pvb	Og?;AX	IB	P/10W	(7)op	8,p17;9:22,p1:23,p65; 24,p78;32	
48	(48)	SAM - MAIN GLACIER	Cu,Pb,Zn,Ag,Au	fl(stg,mas,dis)	(skn;pl,hyd?)	Pvb?	/AX	IB	P/10W	(8)op	8,p17;9:22,p1:23,p65; 24,p79;32	
49	(49)	KEL	Cu,Ag,Au,Zn	vn	hyd,rl?	PMvs	W3,TU?	IB?	P/09W	(6)op	8,p17;9:24,p81	
50	(50)	VALIA	Cu,Pb,Zn,Ag,Fo	stf,mas,sig	vlc	Pvb	AX	IB	P/07W	(7)op	8,p17;9:24,p83	
51	(51)	OUNT BIGGER	Pb,Zn,Cu,Ag	vn,stg	hyd,rl?	Pvb	AX	IB	P/07W	(6)op	8,p17;9:24,p84	
52	(52)	PENDANT GLACIER	Zn,Cu,Ag,Au,Pb	mas,vn,dis	exh?	ODcs?	AX	IB	P/12E	(7)op	8,p17;9:24,p86	
53	(53)	PAMPERO RIDGE	Zn,Cu,Ag	dis?,vn	vlc?,hyd?	uPv	AX	IB	P/13E	(7)op	8,p17;9:13,p173; 14,p135;24,p88	
54	(54)	GRAMPS CRAG	Cu,Zn,Ag	dis?	unk	uPv,	uPpc?	AX	IB	P/13E	(7)op	8,p17;9:13,p173;24,p90
55	(55)	AEOLIAN STEEPLE	Zn,Ag,Cu	unk	vlc?	uPv	AX	IB	P/12E	(7)op	8,p17;9:13,p173; 14,p135;23,p65;24,p92	
56	(56)	VEGA (L145)	Cu,Ag,Zn,Au	dis	skn;pl,hyd?	uPs	Og;AX	IB	P/10E	(7)op	8,p17;9:22,p1:23,p65; 24,p94;32	
57	(57)	ALU	Cu,Mo,Wo	dis,stw,vn	skn;pl,hyd?	SDc	JKg;AX	IB	P/03E	(7)op	8,p17;9:22,p1:24,p96	
58	(58)	SAC	Cu,Pb	dis	unk	Or?	Og?;JKg;AX	IB	P/06E	(7)op	8,p17;9:22,p1:24,p98	
59	(59)	HAINES ROAD	Ba	vn	hyd?,rl?	uPs?	AX?;W2?	IB	P/15E	(7)op	8,p17;9:24,p99	
60	(60)	SQUAW VALLEY	Ab	vn?	hyd?,rl?	Mb?, P'rb,ub?	AX?;W2?	IB	P/14E	(7)op	8,p17;9:24,p100	
61	(61)	RIME (X, MUS)	Au,Ag,Cu,Zn,Co	stb,mas,dis?	vlc?,exh?	uPpc,	uPv?	AX	P/12E	(6)op	8,p17;9:11,p159; 13,p173;14,p135; 16,p109;17,p191; 18,p197;23,p65; 24,p103	
62	(62)	HERBERT (MOUTH) WEST	Au,Ag,Cu,Co,Pb	dsc,pod,mas,dis	vlc;hyd?	Pvb	AX	IB	P/07E	(7)op	8,p17;9:12;15,p365; 16,p109;17,p191; 18,p197;23,p65;	
63	(63)	HERBERT (MOUTH) EAST	Ag,Cu,Zn,Au,Co,Pb	stb,mas,dis	vlc	Pvb	AX	IB	P/07E	(7)op	8,p17;9:12;15,p365; 16,p109;17,p191; 18,p197;23,p65; 24,p105	

OF.	MINE No.	Deposit(s) (No)	Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s) Unit(s)	Host Unit(s)	Plut.:Cov. Terrane	Mgeo Belt	NTS (114)	Status References
64	(64)	LOW HERBERT	Cu,Ag,Au,Zn,Pb,Ba	stf?,dis	vfc	Pvb	AX		IB	P/07E	(6)op	8,p17;9;12;15,p365; 16,p109;17,p191; 18,p197;23,p65; 24,p107
65	(65)	HIGH HERBERT NORTH	Ba,Cu,Ag,Au,Pb,Zn	vn,stg,dis	vic;hyd,rpl	Pvb	AX		IB	P/08W	(6)op	8,p17;9;12;15,p365; 16,p109;17,p191; 18,p197;23,p65; 24,p109
66	(66)	JARVIS SOUTH	Cu	unk	unk	Pvb	AX		IB	P/08W	(7)op	8,p17;9;15,p365; 24,p111
67	(67)	HIGH JARVIS	Zn,Ag,Au,Cu,Pb	stf,mas	hyd?	ODcs	AX		IB	P/07E	(6)op	8,p17;9;15,p365; 24,p112
68	(68)	GRIZZLY HEIGHTS 1	Au,Ag	vn,bx	hyd,rpl	Pvb,(Dc,p)	AX		IB	P/07E	(6)op	8,p17;9;15,p365; 23,p65;24,p114
69	(69)	KUD	Cu,Ag,Au	vn?,dis?	hyd?,rpl?	uP?	AX		IB	P/14E	(7)op	8,p17;9;24,p116
70	(70)	RED MOUNTAIN (FAIR)	Cu,Pb,Zn,Ag,Au,As	mas;sig,vn,dis	skn;vfc?;hyd	uPpc;v; Mf?	Mag: (AX;Tvs/AX)		IB	P/11E	(6)op	8,p17;9;17,p191;22,p1; 24,p117;29,p117; 30,p237
71	(71)	LOW JARVIS (JUMAR)	Cu,Zn,Ag,Ba	stf,mas,dis	vfc,exh	Pvb	AX		IB	P/08W	(6)op	8,p17;9;12;15,p365; 16,p109;17,p191; 18,p197;22,p123,p65; 24,p119;31,p1
72	(72)	EMPIRE (L288)	Cu,Ag	dis?	skn;hyd,rpl	uPs	Obj:AX		IB	P/10E	(7)op	1,p29;3,p40;8,p17;9; 22,p123,p65;24,p120; 32
73	(73)	ARIZONA (L285)	Cu,Ag,Pb	dis?	skn;hyd,rpl	uPs	Obj:AX		IB	P/10E	(7)op	1,p29;3,p40;8,p17;9; 22,p123,p65;24,p122; 32
74	(74)	GILROY (L730)	Cu,Ag,Zn,Fe	mas?,dis?	skn;hyd,rpl	uPs	Obj:AX		IB	P/10E	(7)op	1,p29;3,p40;8,p17;9; 22,p123,p65;24,p124; 32
75	(75)	MOCKING BIRD (HARTFORD)	Cu,Ag,Zn	dis?	skn;hyd,rpl	uPs	Obj:AX		IB	P/10E	(7)op	1,p29;3,p40;8,p17;9; 22,p123,p65;24,p126; 32

OF. No.	MINF (No.)	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s)	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (114)	NTS Belt (114)	Status	References
76 (76)	NEW YORK (L287 (CARIBOO))	Cu,Ag,Au	mas,dis?	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;24,p128; 32		
77 (77)	EVENING (WHITE HORSE)	Ag,Cu,Zn	dis	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;24,p130; 32		
78 (78)	FRISCO (L154)	Ag,Zn,Cu,Pb,Au	dis?	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;24,p132; 32		
79 (79)	COLUMBIA (FAIRFIELD)	Cu,Ag,Au	mas,dis?	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;24,p134; 32		
80 (80)	SADDLE 3/4	Au,Ag	vn,dis	hyd	0g	0gAX	IB	P/10E	(7)op	8,p17;9;22,p1;23,p65; 24,p136;32		
81 (81)	SADDLE 2	Cu,Ag,Au,Fe	dis	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(7)op	8,p17;9;22,p1;23,p65; 24,p138;32		
82 (82)	KLEHINI RIVER (KR1) (BETA)	Au,Ag,Cu	vn,dis	hyd,rpl?	0g	0gAX	IB	P/10E	(6)op	8,p17;9;22,p1;23,p65; 24,p140;32		
83 (83)	KLEHINI RIVER (KR4) (BETA)	Au,Ag,Cu	vn,stw,dis	hyd,rpl?	0g	0gAX	IB	P/10E	(6)op	8,p17;9;22,p1;23,p65; 24,p142;32		
84 (84)	KLEHINI RIVER (KR7) (DELTA)	Cu,Ag,Au	vn,dis	hyd	0g	0gAX	IB	P/10E	(6)op	8,p17;9;22,p1;23,p65; 24,p144		
85 ( )	JARVIS	Pb,Ag,Au	pod?,dis?	skn;hyd,rpl	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;32		
86 ( )	EAST ARM GLACIER	Cu	stf?,mas,dis	vc,c,exh?	uPpc?v?	AX	IB	P/12E	(6)op	8,p17;9;19,p1;20,p723		
87 ( )	TORONTO (L289)	Zn,Cu,Ag	unk	skn	uPs	0gAX	IB	P/10E	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;32		
88 ( )	WOODCHOPPER	Pb,Zn,Ag	unk	skn	uPs	0gAX	IB	P/09W	(6)op	1,p29;3,p40;8,p17;9; 22,p1;23,p65;32		
89 ( )	PANTHER (CAT)	Cu,Ag,Mo	bx,stw,stg	hyd,rpl?,prp?	0g	0gAX	IB	P/10E	(6)op	3,p40;8,p17;9;22,p1; 23,p65;32		
90 ( )	TIKKE GLACIER	Au,Cu,Ag	vn,stg,dis	unk	Psp?	AX	IB	P/06W	(7)op	8,p17;9,32		
91 ( )	WATSON	Cu,Au	vn,dis,flt	hyd,rpl	PMvs	W3,TU?	IB?	P/15E	(7)op	3,p60;8,p17;9;32		

OF. No.	MINF No.	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt (114)	NTS Belt (114)	Status References
92 ( )		BLANCHARD LAKE	Cu	dis,flt	skn	Mc	W3,TU?	CB	P/16W	(7)op 3,p60;8,p17;9;32
93 ( )		SONORA L727	Zn,Pb,Cu,Fe	mas,dis	skn	uPs	0g;AX	IB	P/10E	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
94 ( )		CALGARY	F <sub>e</sub>	vn,dis	hyd,tpl?	uPs	AX	IB	P/10E	(7)op 1,p29;3,p40;8,p17;9; 23,p65;32
95 ( )		WONDERFUL L726	Pb,Zn,Cu,Ag	pod?;mas	skn?	uPs	0g;AX	IB	P/10E	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
96 ( )		CUSTER	Zn,Pb,Cu,Ag,Au	unk	skn?	uPs	0g;AX	IB	P/10E	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
97 ( )		BLUE BIRD	Cu,Ag	unk	skn?	uPs	0g;AX	IB	P/10E	(7)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
98 ( )		TAGISH	Pb	unk	skn?	uPs	0g;AX	IB	P/10E	(7)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
99 ( )		SOLDIERS THREE	Pb,Zn,Cu,Ag	mas?	skn?	uPs	0g;AX	IB	P/10E	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
100 ( )		ELISE L902	Cu	mas,vn	skn?;hyd,tpl	uPs	0g;AX	IB	P/10E	(7)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
101 ( )		HORRIBLE L810	Cu,Au,Ag	dis	hyd?;tpl?	0f?	0g;AX	IB	P/10E	(7)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
102 ( )		LE ROI	Pb,Ag	mas,vn	skn?;hyd?;tpl?	uPs	0g;AX	IB	P/09W	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
103 ( )		ATLIN L904	Cu,Zn	unk	skn?	uPs	0g;AX	IB	P/10E	(6)op 1,p29;3,p40;8,p17;9; 22,p123,p65;32
104 ( )		PARTON	Au,Zn	vn	hyd?	OSwp?	AX	IB	P/10E	(6)op 8,p17;9;32
105 ( )		MULE CREEK	Au,Ag,Cu	mas?	unk	Ps,PRv?	W2	IB	P/15E	(6)op 8,p17;9;32
106 ( )		JOE	Zn,Pb	unk	unk	uPv?,pc?	AX	IB	P/11E	(7)op 8,p17;9;17,p194;32
107 ( )		STONE	Cu,Zn,Pb,Fe	pod,vn	skn	uPs	0g;AX	IB	P/09W	(7)op 3,p40;8,p17;9;32
108 ( )		CHILKAT	Cu,Ni,PGE	dis	mgn	PRub	W2	IB	P/15W	(7)op 3,p25;8,p17;9;32
109 ( )		GRIZZLY HEIGHTS	Zn,Pb,Ag,Ba	flt,(mas)	(unk)	?	/AX	IB	P/07E	(8)op 8,p17;9;15,p365

OF.	MINF No.	Deposit(s) Name(s) (No.)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Deposit(s) Unit(s)	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo NTS Belt (114)	Status References
110 ( )	BUCKWELL GLACIER	Cu,Zn,Pb,Co,Ag,Au	fit,(stf,mas)	(vlo)	? /AX	IB	P/07E	(8)op	8,p17;9;15,p365;32	
111 ( )	HENSHI CREEK 1	Cu,Zn,Co,Ag	fit,(stf,mas)	(vlo)	uPpc,v /AX	IB	P/11W	(8)op	8,p17;9;13,p173;19,p1; 20,p723	
112 ( )	HENSHI CREEK 2	Cu,Zn,Co,Ag	fit,(stf,mas)	(vlo)	uPpc,v /AX	IB	P/11W	(8)op	8,p17;9;19,p1;20,p723	
113 ( )	HENSHI CREEK 3	Cu,Zn,Co,Ag	fit,(stf,mas)	(vlo)	uPpc,v /AX	IB	P/11W	(8)op	8,p17;9;19,p1;20,p723	
114 ( )	SOUTH RANGE CREEK	Cu	fit,(mas?)	(vlo?)	? /AX	IB	P/13E	(8)op	8,p17;9;19,p1;20,p723	
115 ( )	PARTON RIVER	Cu,Ag,Au	fit,(dis)	(unk)	? /AX	IB	P/15W	(8)op	8,p17;9;32	
116 ( )	SELTAT PEAK	Cu	fit,(unk)	(skn)	? /AX	IB	P/09W	(8)op	8,p17;9;32	
117 ( )	GP	Au	fit,(unk)	(unk)	PMsv? /W1?	IB	P/03E	(8)op	8,p17;9;32	
118 ( )	CANADIAN BOULDERADO	Cu,Zn,Pb,Ag,Ba	fit,(stf,mas)	(vlo)	Pvb? /AX	IB	P/08W	(8)op	8,p17;9;15,p365; 17,p191	
119 ( )	COAL CREEK	coal	crc,stf	sed	Ps Tws/AX	IB	P/14W	(7)op	7;8,p17;9	
120 ( )	KUSAWAK RANGE	coal	crc,stf	sed	Ps Tws/AX	IB	P/10E	(7)op	8,p17;9	
121 ( )	SQUAW CREEK COAL	coal	crc,stf	sed	Ps Tws/AX	IB	P/14E	(7)op	3,p61;8,p17;9;28,p7	
122 ( )	TATS CREEK COAL 1	coal	crc,stf	sed	Ps Tws/AX	IB	P/12E	(7)op	7;8,p17;9	
123 ( )	TATS CREEK COAL 2	coal	crc,stf	sed	Ps Tws/AX	IB	P/12E	(7)op	7;8,p17;9	
124 ( )	TATS CREEK COAL 3	coal	crc,stf	sed	Ps Tws/AX	IB	P/12E	(7)op	7;8,p17;9	
125 ( )	TATS LAKE COAL	coal	crc,stf	sed	Ps Tws/AX	IB	P/12E	(7)op	7;8,p17;9	
126 ( )	TALBOT CREEK	coal	crc,stf	sed	Ps Tws/AX	IB	P/15W	(7)op	3,p61;8,p17;9	
127 ( )	DATLASAKA	bentonite	stf	sed	Qs? /AX	IB	P/15W	(7)op	8,p17;9;32	
128 (4)	SQUAW (DOLLS) CREEK	Au,Cu,Ag,Fe,Pb	fly,unc	plc	Qs /AX,W2)	IB	P/14E	(1)op	3,p36;4,p50;6,p51; 8,p17;9;24,p7;28,p7; 34,p74;35,p90	

OF. No.	MINF (No)	Deposit(s) Name(s)	Commodity(s)	Deposit(s) Form	Deposit(s) Type	Host Unit(s)	Plut.:Cov. /Terrane	Mgeo Belt	NTS (114)	Status References
129 (16)	GOLD RUN (BLIZZARD) CREEK	Au	flv.unc	plc	Qs	/AX,W2?	IB	P/15W	(4)op	3,p39;8,p17-9;24,p30; 35,p93
130 ( )	KLEHINI RIVER	Au	flv.unc	plc	Qs	/AX	IB	P/08W	(7)op	3,p39;8,p17-9

\* No mineral deposits have been recorded from NE Yakutat map area; (reference; 8,p17;10).

#### Explanation of Mineral Deposit Information for Tatshenshini River (114P) map area

⊗ 4 mineral deposit(s) locality and map reference number (on map): location accuracy may vary.  
N.B. locations of placers (plc) mostly are very approximate and are "spot placed" only to indicate creeks or sections of creeks where placers have been mined or occur.

OF. No. map reference number.

MINF (No) MINF (MINFILE: British Columbia Ministry of Energy Mines and Petroleum Resources) No (number) for NTS 114P Tatshenshini River, Dec. 1988 edition.  
Deposit(s) Name(s) name (alternate name) of deposit(s). N.B. for placers (plc) only; name is creek or river where deposit(s) occur.  
N.B. e.g. L727 = Crown-granted mineral claim lot number; e.g. (L722) = original lot number, now reverted.

Commodity(s) Abbreviations elements present are listed in decreasing order of abundance; elements present in trace amounts are included, but not specified.

Ab	asbestos	Au	gold	Co	cobalt	Mo	molybdenum	Sb	antimony
An	anhydrite	Ba	barium	Cu	copper	Ni	nickel	Sr	strontium
Ag	silver	Bi	bismuth	Fe	iron	Pb	lead	Wo	tungsten
As	arsenic	Cd	cadmium	Gy	gypsum	PGE	platinum gp. elements	Zn	zinc

#### Deposit(s) Form Abbreviations

cnc	concordant	stf	stratiform	stw	stockwork	mas	massive	unc	unconsolidated
disc	discordant	pod	podiform	stg	stringer	dis	disseminated	fit	float
stb	stratabound	bx	breccia	vn	vein	flv	fluvial deposit	unk	unknown

#### Deposit(s) Type Abbreviations

hyd	hydrothermal	vlc	volcanic	sed	sedimentary
rpl	replacement	exh	exhalative	evp	evaporite

Host Unit(s) 1:250 000 scale map unit(s) host to mineral deposit(s).

#### Plut.:Cov./Terrane

(Fig. 2 map unit(s) host to mineral deposit(s). Plut. = Plutonic Rocks: : = intruded into: Cov. = Cover Rocks: / = deposited on: Terrane = Terrane.  
e.g. Mg:(AX;Tvs/AX) = Mg: = intruded into (:) = both AX and Tvs which was / = deposited on AX.  
N.B. for placer deposits and float only; e.g. /(AX,W2) = Qs (not depicted in Fig. 2) / = deposited on (,) = both AX and W2.

Mg = Miocene subvolcanics: Qg = Oligocene granitoids and or subvolcanics: JKg = Late Jurassic to Earliest Cretaceous granitoids: Tvs = Tertiary volcanics and or sediments: W1?, W2, W3 = Wrangellia Terrane; segments 1?, 2, 3: TU = Taku Terrane: AX = Alexander Terrane.

**Mgeo Belt** Morphogeological Belts (Fig. 2 inset) in which mineral deposit(s) occur (IB = Insular Belt; CB = Coast Belt).

NTS National Topographic System map sheet reference.  
(114) = 1:1 000 000 scale: O / P = 1:250 000 scale: 01 to 16 = 1:50 000 scale; E = East, or W = West half.

#### Status Abbreviations

- (1) Being produced; commodity(s) is (are) being extracted for sale.
- (2) Never produced; reserves (grade + 3 dimens.) or resources known.
- (3) Past producer : reserves (grade + 3 dimens.) or resources known.
- (4) Past producer : reserves or resources exhausted, or not known.
- (5) Grade, 2 dimensions (length and width) known: reserves unknown.
- (6) Grade, 1 dimension (length or width) known : by drill or chip sample.
- (7) Outcrop showing; or commodity(s) known, but unpublished: ± grab sample assays.
- (8) Float; provenance known or unknown: ± grab sample assays.

ip = inside, or op = outside of Kluane National Park Reserve.

**References** (e.g. 6,p50 = Read and Monger (1976) page 50; N.B. p = initial page of reference cited; (1976) = year of publication.  
N.B. references listed are main sources only; for references such as BCMEMPR publications (not given), newspaper releases (e.g. Northern Miner), and individual assessment reports, etc., the reader should see BCMEMPR Minfile 114P (1988), or Northern B.C. Mineral Inventory (Archer, Cathro & Associates (1981) Limited).

- 1 McConnell (1914).
- 2 Eakin (1919).
- 3 Watson (1948).
- 4 Kindle (1953).
- 5 Mulligan (1966).
- 6 Read and Monger (1976).
- 7 J.McDougall,p.comm.(1977).
- 8 Campbell and Dodds (1979).
- 9 Campbell and Dodds(1983a).
- 10 Campbell and Dodds (1983b).
- 11 MacIntyre (1983).
- 12 Still (1984).
- 13 MacIntyre (1984).
- 14 Grammon and Chandler (1986).
- 15 MacIntyre and Schroeter (1985).
- 16 Gehrels et al. (1986).
- 17 Schroeter and MacIntyre (1986).
- 18 MacIntyre (1986).
- 19 Day (1985).
- 20 Day et al. (1987).
- 21 White (1986).
- 22 Dodds and Campbell (1988).
- 23 Dodds (1988).
- 24 BCMEMPR Minfile, 114P (1988).
- 25 Peter (1989).
- 26 Mandy (1932).
- 27 Schroeter (1978).
- 28 Cockfield (1928).
- 29 Ellingør & Ray (1989).
- 30 Ray et al. (1990).
- 31 Fields (1986).
- 32 NBCMI, 114P (1989).
- 33 Geddes Res. Ltd. (1990).
- 34 Mandy (1933).
- 35 Mandy (1934).

## REFERENCES

**BCMEMPR Minfile, 114P (1988).**

1988: Minfile NTS No. 114P, Tatshenshini River; British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Survey Branch, December 1988 edition, 145 p.

**Berg, H.C., Jones, D.L., and Coney, P.J.**

1978: Map showing pre-Cenozoic tectonostratigraphic terranes of southeastern Alaska and adjacent areas; United States Geological Survey, Open File Report 78-1085, 2 sheets.

**Berg, H.C., Jones, D.L., and Richter, D.H.**

1972: Gravina-Nutzotin Belt - tectonic significance of an Upper Mesozoic sedimentary and volcanic sequence in southern and southeastern Alaska; United States Geological Survey Professional Paper 800-D, p. D1-D24.

**Bostock, H.S.**

1933: The mining industry of Yukon, 1932; in Geological Survey of Canada, Summary Report 1932, Part A II, p. 1AII-14AII.

1934: The mining industry of Yukon, 1933; in Geological Survey of Canada, Summary Report 1933, Part A, p. 1A-8A.

1935: The mining industry of Yukon, 1934; Geological Survey of Canada, Memoir 178, 10 p.

1936: Mining industry of Yukon, 1935; Geological Survey of Canada, Memoir 193, 12 p.

1937: Mining industry of Yukon, 1936; Geological Survey of Canada, Memoir 209, 13 p.

1938: Mining industry of Yukon, 1937; Geological Survey of Canada, Memoir 218, 21 p.

1939: Mining industry of Yukon, 1938; Geological Survey of Canada, Memoir 220, 21 p.

1941: Mining industry of Yukon, 1939 and 1940; Geological Survey of Canada, Memoir 234, 40 p.

1952: Geology of northwest Shakwak Valley, Yukon Territory; Geological Survey of Canada, Memoir 267, 54 p.

1957: Yukon Territory: selected field reports of the Geological Survey of Canada 1898 to 1933; Geological Survey of Canada, Memoir 284, 650 p.

**Boucher, G. and Fitch, T.J.**

1969: Microearthquake seismicity of the Denali fault; Journal of Geophysical Research, v. 74, p. 6638-6648.

**Brew, D.A. and Ford, A.B.**

1978: Megalineament in southeastern Alaska marks southwest edge of Coast Range batholith complex; Canadian Journal of Earth Sciences, v. 15, no. 11, p. 1763-1772.

1981: The Coast plutonic complex sill, southeastern Alaska; in The United States Geological Survey in Alaska: Accomplishments during 1979, United States Geological Survey Circular 823-B, p. B96-B98.

1985: Preliminary reconnaissance geologic map of the Juneau, Taku River, Atlin, and part of the Skagway 1:250 000 quadrangles, southeastern Alaska; United States Geological Survey, Open-File Report 85-395, 23 p.

**Brew, D.A. and Morrell, P.M.**

1983: Intrusive rocks and plutonic belts of southeastern Alaska, U.S.A.; in Circum-Pacific plutonic terranes, J.A. Roddick (ed.); Geological Society of America, Memoir 159, p. 171-193.

**Brew, D.A., Johnson, B.R., Grybeck, D., Griscom, A., Barnes, D.F., Kimball, A.L., Still, J.C., and Rataj, J.L.**

1978: Mineral resources of the Glacier Bay National Monument Wilderness Study Area, Alaska; United States Geological Survey Open-File Report 78-494, 670 p.

**Brooks, A.H.**

1900: A reconnaissance from Pyramid Harbour to Eagle City, Alaska, including a description of the copper deposits of the upper White and Tanana rivers; United States Geological Survey, Annual Report 21, part 2, p. 331-391.

**Cairnes, D.D.**

1915a: Upper White River district, Yukon; Geological Survey of Canada, Memoir 50, 191 p.

1915b: Exploration in southwestern Yukon; Geological Survey of Canada, Summary Report 1914, p. 10-37.

**Cameron, A.R. and Birmingham, T.F.**

1970: Radioactivity in western Canadian coals; Geological Survey of Canada, Paper 70-52, 35 p.

**Campbell, F.A.**

1956: Nickeliferous sulphide deposits at Quill Creek and White River, Yukon; M.A. thesis, Queen's University.

1960: Nickel deposits in the Quill Creek and White River areas, Yukon; Canadian Institute of Mining and Metallurgy, Transactions, v. 63, p. 662-668.

**Campbell, J.D.**

1967: Guide to coal deposits, Yukon and Mackenzie Territories: a compilation; Research Council of Alberta, Report 66-6, 33 p.

**Campbell, S.W.**

1976: Nickel-copper sulphide deposits in the Kluane Ranges, Yukon Territory; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, Open File Report EGS 1976-10, 17 p.

1977: Geology of the Wellgreen property; in North of 60 - Mineral Industry Report 1976, Yukon Territory, Indian and Northern Affairs Canada, Report EGS 1977-1, p. 55-67.

1981: Geology and genesis of copper deposits and associated host rocks in and near the Quill Creek area, southwestern Yukon; Ph.D. thesis, University of British Columbia, Vancouver, 215 p.

**Campbell, R.B. and Dodds, C.J.**

1975: Operation Saint Elias, Yukon Territory; in Report of Activities, Part A; Geological Survey of Canada, Paper 75-1A, p. 51-53.

1978: Operation Saint Elias, Yukon Territory; in Current Research, Part A; Geological Survey of Canada, Paper 78-1A, p. 35-41.

1979: Operation Saint Elias, British Columbia; in Current Research, Part A; Geological Survey of Canada, Paper 79-1A, p. 17-20.

1982a: Geology of SW Kluane Lake map area (115G and F (E1/2)), Yukon Territory; Geological Survey of Canada, Open File 829.

- 1982b: Geology of Mount St. Elias map area (115B and C (E1/2)), Yukon Territory part; Geological Survey of Canada, Open File 830.
- 1982c: Geology of SW Dezadeash map area (115A), Yukon Territory; Geological Survey of Canada, Open File 831.
- 1983a: Geology of Tatshenshini River map area (114P), British Columbia; Geological Survey of Canada, Open File 926.
- 1983b: Geology of Yakutat map area (114O), British Columbia part; Geological Survey of Canada, Open File 927.
- 1983c: Terranes and major faults of the Saint Elias Mountains, Yukon Territory, British Columbia, and Alaska; Geological Association of Canada, Annual Meeting Victoria, Programs with Abstracts, p. A10.

**Campbell, R.B. and Elsbacher, G.H.**

- 1974: Operation Saint Elias, Yukon Territory; in Report of Activities, Part A; Geological Survey of Canada, Paper 74-1A, p. 11-12.

**Canmindex, 115A.**

- 1989: Canmindex, sheet no. 115A: (Canadian Mineral Occurrence Index) of the Geological Survey of Canada (after Picklyk et al., 1978); Geological Survey of Canada, computerized file.

**Canmindex, 115B.**

- 1989: Canmindex, sheet no. 115B: (Canadian Mineral Occurrence Index) of the Geological Survey of Canada (after Picklyk et al., 1978); Geological Survey of Canada, computerized file.

**Canmindex, 115G/F**

- 1989: Canmindex, sheet nos. 115G & F: (Canadian Mineral Occurrence Index) of the Geological Survey of Canada (after Picklyk et al., 1978); Geological Survey of Canada, computerized file.

**Carlson, P.R., Plafker, G., and Bruns, T.R.**

- 1985: Map and selected seismic profiles of the seaward extension of the Fairweather Fault, eastern Gulf of Alaska; United States Geological Survey, Map MF-1722, 2 sheets.

**Carriere, J.J., Sinclair, W.D., and Kirkham, R.V.**

- 1981: Copper deposits and occurrences in Yukon Territory; Geological Survey of Canada, Paper 81-12, 62 p.

**Christopher, P.A., White, W.H., and Harakal, J.E.**

- 1972: K-Ar dating of the 'Cork' (Burwash Creek) Cu-Mo prospect, Burwash Landing area, Yukon Territory; Canadian Journal of Earth Sciences, v. 9, p. 918-921.

**Cockfield, W.E.**

- 1928: Dezadeash Lake area, Yukon; in Geological Survey of Canada, Summary Report, 1927, Part A, p. 1A-7A.
- 1930: The mining industry of Yukon, 1929; in Geological Survey of Canada, Summary Report, 1929 Part A, p. 1A-15A.

**Coney, P.J., Jones, D.L., and Monger, J.W.H.**

- 1980: Cordilleran suspect terranes; Nature, v. 288, p. 329-333.

**Clague, J.J.**

- 1979a: The Denali Fault System in southwest Yukon Territory - A Geologic hazard ?; in Current Research, Part A; Geological Survey of Canada, Paper 79-1A, p. 169-178.

- 1979b: An assessment of some possible flood hazards in Shakwak Valley, Yukon Territory; in Current Research, Part B; Geological Survey of Canada, Paper 79-1B, p. 63-70.
- 1981: Landslides at the south end of Kluane Lake, Yukon Territory; Canadian Journal of Earth Sciences, v. 18, p. 959-971.
- Clague, J.J. and Rampton, V.N.**
- 1982: Neoglacial Lake Alsek; Canadian Journal of Earth Sciences, v. 19, no. 1, p. 94-117.
- Clarke, G.K.C.**
- 1982: Glacier outburst floods from "Hazard Lake", Yukon Territory, and the problem of flood magnitude prediction; Journal of Glaciology, v. 28, no. 98, p. 3-21.
- Clarke, G.K.C. and Mathews, W.H.**
- 1981: Estimates of the magnitude of glacier outburst floods from Lake Donjek, Yukon Territory, Canada; Canadian Journal of Earth Sciences, v. 18, no. 9, p. 1452-1463.
- Clarke, G.K.C., Schmok, J.P., Ommann, C.S.L., and Collins, S.G.**
- 1986: Characteristics of surge-type glaciers; Journal of Geophysical Research, v. 91, no. B7, p. 7165-7180.
- Craig, D.B. and Laporte, P.J.**
- 1972: North of 60 - Mineral Industry Report 1969 and 1970, v. 1 - Yukon Territory and southwestern sector, District of MacKenzie; Indian and Northern Affairs Canada, Northern Economic Development Branch, Report EGS 1972-1, 188 p.
- Craig, D.B. and Milner, M.W.**
- 1975: North of 60 - Mineral Industry Report 1971 and 1972, v. 1 of 3 - Yukon Territory; Department of Indian Affairs and Northern Development, Northern Economic Development Branch, Report EGS 1975-76, 169 p.
- Davis, A. and Plafker, G.**
- 1985: Comparative geochemistry and petrology of Triassic basaltic rocks from the Taku terrane on the Chilkat Peninsula and Wrangellia; Canadian Journal of Earth Sciences, v. 22, p. 183-194.
- Day, S.J.**
- 1985: A petrographic and geochemical comparison of massive sulphide boulders in outwash from East Arm Glacier, St. Elias Mountains, British Columbia, with the Windy-Craggy deposit; B.Sc. thesis, University of British Columbia, Vancouver, 79 p.
- Day, S.J., Broster, B.E., and Sinclair, A.J.**
- 1987: Sulphide erratics applied to subglacial exploration: St. Elias Mountains, British Columbia; Canadian Journal of Earth Sciences, v. 24, p. 723-730.
- Debicki, R.L.**
- 1982 Yukon Mineral Industry 1941-1959; Indian and Northern Affairs Canada, Yukon, 136 p.
- 1983: Yukon Placer Mining Industry 1978-1982; Indian and Northern Affairs Canada, Exploration and Geological Services and Northern Affairs Program, Yukon, 203 p.
- Debicki, R.L. and Gilbert, G.W.**
- 1986 Yukon Placer Mining Industry 1983-1984; Indian and Northern Affairs Canada, Placer Mining Section, Mining Engineering Division, Northern Affairs Program, Yukon, 153 p.
- Denton, G.H. and Armstrong, R.L.**
- 1969: Miocene-Pliocene glaciations in southern Alaska; American Journal of Science, v. 267, p. 1121-1142.

- Dodds, C.J.**  
 1988: Geological mapping in Tatshenshini River map area, British Columbia; in Current Research, Part E; Geological Survey of Canada, Paper 88-1E, p. 65-72.
- Dodds, C.J. and Campbell, R.B.**  
 1988: Potassium-argon ages of mainly intrusive rocks in the Saint Elias Mountains, Yukon and British Columbia; Geological Survey of Canada, Paper 87-16, 43 p.
- Downey, M.E.**  
 1978: Geology, K-Ar and Rb-Sr geochronometry and chemistry of the Bock's Brook stock in the Kluane Ranges, southwestern Yukon Territory; B.Sc. thesis, University of British Columbia, Vancouver, 67 p.
- Downey, M.E., Armstrong, R.L., and Parrish, R.R.**  
 1980: K-Ar, Rb-Sr and fission track geochronometry of the Bock's Brook Stock, Kluane Ranges, southwestern Yukon Territory; in Current Research Part B; Geological Survey of Canada, Paper 80-1B, p. 189-193.
- Eaklin, H.M.**  
 1919: The Porcupine gold placer district, Alaska; United States Geological Survey, Bulletin 699, 29 p.
- Eckstrand, O.R.**  
 1984: Canadian mineral deposit types: a geological synopsis; Geological Survey of Canada, Economic Geology Report 36, 86 p.
- Elsbacher, G.H.**  
 1975: Operation Saint Elias, Yukon Territory: Dezadeash Group and Amphitheatre Formation; in Report of Activities, Part A; Geological Survey of Canada, Paper 75-1A, p. 61-62.  
 1976: Sedimentology of the Dezadeash flysch and its implications for strike-slip faulting along the Denali Fault, Yukon Territory and Alaska; Canadian Journal of Earth Sciences, v. 13, p. 1495-1513.
- Elsbacher, G.H. and Hopkins, S.L.**  
 1977: Mid-Cenozoic paleogeomorphology and tectonic setting of the St. Elias Mountains, Yukon Territory; in Report of Activities, Part B; Geological Survey of Canada, Paper 77-1B, p. 319-335.
- Erdmer, P.**  
 1989: The Nisling Schist in eastern Dezadeash map area, Yukon; in Current Research, Part E; Geological Survey of Canada, Paper 89-1E, p. 139-144.  
 1990: Studies of the Kluane and Nisling assemblages in Kluane and Dezadeash map areas, Yukon; in Current Research, Part E; Geological Survey of Canada, Paper 90-1E, p. 107-111.
- Ettlinger, A.D. and Ray, G.E.**  
 1989: Precious metal enriched skarns in British Columbia: an overview and geological study; British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Branch, Paper 1989-3, 128 p.
- Evenchick, C.A., Friday, S.J., and Monger, J.W.H.**  
 1987: Potential hosts to platinum group element concentrations in the Canadian Cordillera; Geological Survey of Canada, Open File 1433.
- Eyles, C.H. and Eyles, N.**  
 1989: The upper Cenozoic White River "tillites" of southern Alaska: Subaerial slope and fan-delta deposits in a strike-slip setting; Geological Society of America Bulletin, v. 101, p. 1091-1102.

- Farrar, E., Clark, A.H., Archibald, D.A., and Way, D.C.**  
 1988: Potassium-argon age of granitoid plutonic rocks, southwest Yukon Territory, Canada; Isochron West, no. 51, p. 19-23.
- Fields, M.**  
 1986: Geology of the Still's Ridge barite, zinc, copper, silver prospect, St. Elias Mountains, northwestern British Columbia (114P/07E): B.Sc. thesis, University of British Columbia, Vancouver, 62 p.
- Findlay, D.C.**  
 1967: The mineral industry of Yukon Territory and southwestern District of MacKenzie, 1966; Geological Survey of Canada, Paper 67-40, 104 p.
- 1969a: The mineral industry of Yukon Territory and southwestern District of MacKenzie, 1967; Geological Survey of Canada, Paper 68-68, 131 p.
- 1969b: The mineral industry of Yukon Territory and southwestern District of MacKenzie, 1968; Geological Survey of Canada, Paper 69-55, 71 p.
- Fraser, J.A.**  
 1915: Rainy Hollow; British Columbia Minister of Mines Annual Report 1915, p. 64-65.
- Gammon, J.B. and Chandler, T.E.**  
 1986: Exploration of the Windy Craggy massive sulphide deposit, British Columbia, Canada; in Geology in the Real World - The Kingsley Dunham Volume, R.W. Nesbitt and Ian Nichol (ed.), The Institution of Mining and Metallurgy, p. 131-141.
- Gardner, M.C., Bergman, S.C., Cushing, G.W., MacKevett, E.M., Jr., Plafker, G., Campbell, R.B., Dodds, C.J., McClelland, W.D., and Mueller, P.A.**  
 1988: Pennsylvanian pluton stitching of Wrangellia and the Alexander Terrane, Wrangell Mountains, Alaska; Geology, v. 16, p. 967-971.
- Geddes Res. Ltd.**  
 1990: The Windy Craggy Project - a proposed new copper mine in British Columbia; Geddes Resources Limited, March 1990, 7 p.
- Gehrels, G.E. and Berg, H.C.**  
 1984: Geologic map of southeastern Alaska; United States Geological Survey, Open-File Report 84-886, p. 1-28.
- Gehrels, G.E. and Saleeby, J.B.**  
 1987: Geologic framework, tectonic evolution, and displacement history of the Alexander Terrane; Tectonics, v. 6, no. 2, p. 151-173.
- Gehrels, G.E., Dodds, C.J., and Campbell, R.B.**  
 1986: Upper Triassic rocks of the Alexander terrane, SE Alaska and the Saint Elias Mountains of B.C. and Yukon; Geological Society of America, Abstracts with Programs 1986, v. 18, no. 2, p. 109.
- Green, L.H.**  
 1965: The mineral industry of Yukon Territory and southwestern District of MacKenzie, 1964; Geological Survey of Canada, Paper 65-19, 94 p.
- 1966: The mineral industry of Yukon Territory and southwestern District of MacKenzie, 1965; Geological Survey of Canada, Paper 66-31, 137 p.

**Green, L.H. and Godwin, C.I.**

1963: The mining industry of Yukon Territory and southwestern District of MacKenzie, 1962; Geological Survey of Canada, Paper 63-38, 71 p.

1964: The mining industry of Yukon Territory and southwestern District of MacKenzie, 1963; Geological Survey of Canada, Paper 64-36, 94 p.

**Hayes, C.W.**

1892: An expedition through the Yukon district; National Geographic Magazine, v. 4, p. 117-162.

**Horner, R.B.**

1983: Seismicity in the St. Elias region of northwestern Canada and southeastern Alaska; Bulletin of the Seismological Society of America, v. 73, no. 4, p. 1117-1137.

1989: Low-level seismic monitoring at the Windy Craggy deposit in northwestern British Columbia; in Current Research, Part E; Geological Survey of Canada, Paper 89-1E, p. 275-278.

**Hudson, T.**

1983: Calc-alkaline plutonism along the Pacific rim of southern Alaska; in Circum-Pacific plutonic terranes, J.A. Roddick (ed.); Geological Society of America, Memoir 159, p. 159-169.

**Hudson, T. and Plafker, G.**

1982: Paleogene metamorphism of an accretionary flysch terrane, eastern Gulf of Alaska; Geological Society of America Bulletin, v. 93, p. 1280-1290.

**Hudson, T., Plafker, G., and Lanphere, M.A.**

1977: Intrusive rocks of the Yakutat-St. Elias area, south-central Alaska; United States Geological Survey, Journal of Research, v. 5, no. 2, p. 155-172.

**Hughes, O.L., Rampton, V.N., and Rutter, N.W.**

1972: Quaternary geology and geomorphology, southern and central Yukon (northern Canada), 24th International Geological Congress Guidebook, Field Excursion A11, 59 p.

**Hulbert, L.J., Duke, J.M., Eckstrand, O.R., Lyndon, J.W., Scroates, R.F.J., and Cabri, L.J.**

1988: Geological environments of the platinum group elements; Geological Survey of Canada, Open File 1440, 148 p.

**INAC, (1985)**

1985: Yukon Exploration and Geology 1983; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 317 p.

**INAC, (1986)**

1986: Yukon Exploration 1984; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 288 p.

**INAC, (1987)**

1987: Yukon Exploration 1985-1986; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 451 p.

**INAC, (1988)**

1988: Yukon Exploration 1987; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 364 p.

**INAC, (1989)**

1989: Yukon Exploration 1988; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 304 p.

**INAC Overview, (1990)**

1990: Yukon Mining and Exploration Overview, 1989; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, 16 p.

**Jacobson, B.**

1979: Geochronology and petrology of the Tkope River batholith in the Saint Elias Mountains, northwestern B.C.; B.Sc. thesis, University of British Columbia, Vancouver, 47 p.

**Jacobson, B., Parrish, R.R., and Armstrong, R.L.**

1980: Geochronology and petrology of the Tkope River Batholith in the Saint Elias Mountains, northwestern British Columbia; in Current Research, Part B; Geological Survey of Canada, Paper 80-1B, p. 195-206.

**Johnston, R.A.A.**

1915: A list of Canadian mineral occurrences; Geological Survey of Canada, Memoir 74, 275 p.

**Kindle, E.D.**

1953 Dezadeash map-area, Yukon Territory; Geological Survey of Canada, Memoir 268, 68 p.

**Kirkham, R.V.**

1971: Geology of copper and molybdenum deposits; in Report of Activities, Part A; Geological Survey of Canada, Paper 71-1 Part A, p. 85-88.

**Lahr, J.C., Plafker, G., Stephens, C.D., Fogleman, K.A., and Blackford, M.E.**

1979: Interim report on the St. Elias Earthquake of 28 February 1979; United States Geological Survey, Open File Report 79-670, 35 p.

**Lahr, J.C., Stephens, C.D., Hasegawa, H.S., and Boatwright, J.**

1980: Alaskan seismic gap only partially filled by 28 February 1979 earthquake; Science, v. 207, p. 1351-1353.

**LeBarge, W.P. and Morison, S.R.**

1990: Yukon Placer Mining and Exploration 1985-1988; Indian and Northern Affairs Canada, Northern Affairs Program, Yukon, 151 p.

**Lerbekmo, J.F., and Campbell, F.A.**

1969: Distribution, composition, and source of the White River ash, Yukon Territory; Canadian Journal of Earth Sciences, vol. 6, p. 109-116.

**Lerbekmo, J.F., Westgate, J.A., Smith, D.W.G., and Denton, G.H.**

1975: New data on the character and history of the White River volcanic eruption, Alaska; in Quaternary studies: selected papers from IX INQUA Congress, R.P. Suggate and M.M. Cresswell (ed.), Royal Society New Zealand Bulletin 13, p. 203-209.

**Loney, R.A. and Himmelberg, G.R.**

1983: Structure and petrology of the La Perouse gabbro intrusion, Fairweather Range, southeastern Alaska; Journal of Petrology, v. 24, part 4, p. 377-423.

**Lowey, G.W.**

1980: Depositional themes in a turbidite succession, Dezadeash Formation, (Jura-Cretaceous), Yukon; M.Sc. thesis, University of Alberta, Calgary, 149 p.

**MacIntyre, D.G.**

1983: A comparison of the geologic setting of stratiform massive sulphide deposits of the Gataga district with the Midway and Windy-Craggy deposits, northern British Columbia (94F, L; 104O/16; 114P/12); in Geological Fieldwork 1982, British Columbia Ministry of Energy, Mines and Petroleum Resources, Geological Branch, Paper 1983-1, p. 149-170.

- 1984: Geology of the Alsek-Tatshenshini Rivers area (114P); in Geological Fieldwork 1983, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Branch, Paper 1984-1, p. 173-184.
- 1986: The Geochemistry of basalts hosting massive sulphide deposits, Alexander terrane northwest British Columbia (114P); in Geological Fieldwork 1985, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Branch, Paper 1986-1, p. 197-210.
- MacIntyre, D.G. and Schroeter, T.G.**
- 1985: Mineral occurrences in the Mount Henry Clay area (114P/7,8); in Geological Fieldwork 1984, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Branch, Paper 1985-1, p. 365-379.
- MacKevett, E.M., Jr.**
- 1978: Geologic map of the McCarthy quadrangle, Alaska; United States Geological Survey, Map I-1032.
- MacKevett, E.M., Jr. and Plafker, G.**
- 1974: The Border Ranges fault in south-central Alaska; United States Geological Survey, Journal of Research, v. 2, no. 3, p. 323-329.
- MacKevett, E.M., Jr., Robertson, E.C., and Winkler, G.R.**
- 1974: Geology of the Skagway B-3 and B-4 quadrangles, southeastern Alaska; United States Geological Survey Professional Paper 832, 33 p.
- MacLean, T.A.**
- 1914: Lode mining in the Yukon: an investigation of quartz deposits in the Klondike Division; Canada Department of Mines, Mines Branch, Publication no. 222, p. 204.
- Mandy, J.T.**
- 1932: Gold Cord: in Lode-gold deposits of British Columbia; British Columbia Department of Mines, Bulletin No. 1, p. 40-41.
- 1933: Tatshenshini River section; in British Columbia Minister of Mines Annual Report for 1932, p. A74-A79.
- 1934: Tatshenshini River section; in British Columbia Minister of Mines Annual Report for 1933, p. A90-A93.
- McConnell, R.G.**
- 1905: The Kluane mining district; in Geological Survey of Canada, Summary Report 1904, Report A, p. 1A-18A.
- 1906: Headwaters of White River; in Geological Survey of Canada, Summary Report 1905, p. 19-26.
- 1914: Rainy Hollow mineral area, B.C.; in Geological Survey of Canada, Summary Report 1913, Sessional Paper no. 26, p. 29-33.
- Melnyk, W.D.**
- 1972: Geology of the Moose Creek area, St. Elias Range; B.A.Sc. thesis, University of Saskatchewan.
- Miller, D.J.**
- 1960: Giant waves in Lituya Bay, Alaska; United States Geological Survey Professional Paper 354-C, p. 51-86.

- Moffit, F.H. and Knopf, A.**  
 1910: Mineral resources of the Nabesna-White River district, Alaska; United States Geological Survey, Bulletin 417, 64 p.
- Monger, J.W.H. and Berg, H.C.**  
 1984: Lithotectonic terrane map of western Canada and southeastern Alaska; in Lithotectonic terrane maps of the North American Cordilleran, N.J. Silberling and D.L. Jones (ed.); United States Geological Survey, Open-File Report 84-523.
- Morin, J.A.**  
 1989: Yukon gold-silver file; Indian and Northern Affairs Canada, Exploration and Geological Services Division, Yukon, Open File 1989-3, 227 p.
- Morin, J.A., Sinclair, W.D., Craig, D.B., and Marchand, M.**  
 1977: North of 60 - Mineral Industry Report 1976, Yukon Territory; Indian and Northern Affairs Canada, Northern Natural Resources and Environment Branch, Report EGS 1977-1, 264 p.
- Muller, J.E.**  
 1967: Kluane Lake map-area, Yukon Territory (115G, 115F E1/2); Geological Survey of Canada, Memoir 340, 137 p.
- Mulligan, R.**  
 1966: Geology of Canadian tin occurrences; Geological Survey of Canada, Paper 64-54, 22 p.
- NBCMI, 114P (1989).**  
 1989: Northern British Columbia Mineral Inventory, 114P Tatshenshini River; Archer, Cathro and Associates (1981) Limited, private publication, February 28 1989 edition, 98 p.
- Peter, J.M.**  
 1989: The Windy Craggy copper-cobalt-gold massive sulphide deposit, northwestern British Columbia; in Geological Fieldwork 1988, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Survey Branch, Paper 1989-1, p. 455-466.
- Picklyk, D.D., Rose, D.G., and Laramee, R.M.**  
 1978: Canadian Mineral Occurrence Index (CANMINDEX) of the Geological Survey of Canada; Geological Survey of Canada, Paper 78-8, 27 p.
- Plafker, G.**  
 1987: Regional geology and petroleum potential of the northern Gulf of Alaska continental margin; in Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins - Beaufort Sea to Baja California, D.W. Scholl, A. Grantz, and J.G. Vedder (ed.), American Association of Petroleum Geologists, Earth Science Series 6, p. 229-268.
- Plafker, G. and Campbell, R.B.**  
 1979: The Border Ranges fault in the St. Elias Mountains; United States Geological Survey Circular 804-B, p. B102-B104.
- Plafker, G. and Hudson, T.**  
 1980: Regional implications of Upper Triassic metavolcanic and metasedimentary rocks on the Chilkat Peninsula, southeastern Alaska; Canadian Journal of Earth Sciences, v. 17, p. 681-689.
- Plafker, G., Blome, C.D., and Silberling, N.J.**  
 1989: Reinterpretation of lower Mesozoic rocks on the Chilkat Peninsula, as a displaced fragment of Wrangellia; Geology, v. 17, p. 3-6.

**Plafker, G., Hudson, T., and Richter, D.H.**

1977a: Preliminary observations on late Cenozoic displacements along the Totschunda and Denali fault systems; United States Geological Survey Circular 751-B, p. B67-B69.

**Plafker, G., Jones, D.L., and Pessagno, E.A., Jr.**

1977b: A Cretaceous accretionary prism and melange terrane along the Gulf of Alaska margin; United States Geological Survey Circular 751-B, p. B41-B43.

**Plafker, G., Richter, D.H., and Hudson, T.**

1977c: Reinterpretation of the origin of inferred Tertiary tillite in the northern Wrangell Mountains Alaska; United States Geological Survey Circular 751-B, p. B52-B54.

**Plafker, G., Hudson, T., Bruns, T., and Rubin, M.**

1978: Late Quaternary offsets along the Fairweather Fault and crustal plate interactions in southern Alaska; Canadian Journal of Earth Sciences, v. 15. p. 805-816.

**Plafker, G., Nokleberg, W.J., and Lull, J.S.**

1985: Summary of the 1984 TACT geologic studies in the northern Chugach Mountains and southern Copper River basin; United States Geological Survey Circular 967, p. 76-79.

**Plafker, G., Nokleberg, W.J., and Lull, J.S.**

1989: Bedrock geology and tectonic evolution of the Wrangellia, Peninsular, and Chugach terranes along the Trans-Alaska Crustal Transect in the Chugach Mountains and southern Copper River Basin, Alaska; Journal of Geophysical Research, v. 94, no. B4, p. 4255-4295.

**Plafker, G., Jones, D.L., Hudson, T., and Berg, H.C.**

1976: The Border Ranges fault system in the Saint Elias Mountains and Alexander Archipelago; United States Geological Survey Circular 733, p. 14-16.

**Rampton, V.N.**

1979a: Surficial geology and geomorphology, Koidern Mountain, Yukon Territory; Geological Survey of Canada, Map 5-1978.

1979b: Surficial geology and geomorphology, Burwash Creek, Yukon Territory; Geological Survey of Canada, Map 6-1978.

1979c: Surficial geology and geomorphology, Generc River, Yukon Territory; Geological Survey of Canada, Map 7-1978.

1979d: Surficial geology and geomorphology, Congdon Creek, Yukon Territory; Geological Survey of Canada, Map 8-1978.

1981: Surficial materials and landforms of Kluane National Park, Yukon Territory; Geological Survey of Canada, Paper 79-24, 37 p.

1982a: Surficial geology and geomorphology, Kusawak Ranges, British Columbia; Geological Survey of Canada, Map 13-1981.

1982b: Surficial geology and geomorphology, Frederick Lake, Yukon; Geological Survey of Canada, Map 15-1981.

1982c: Surficial geology and geomorphology, Pine Lake, Yukon; Geological Survey of Canada, Map 16-1981.

- Ray, G.E., Ettlinger, A.D., and Meinert, L.D.**  
 1990: Gold skarns: their distribution, characteristics and problems in classification; in Geological Fieldwork 1989, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Survey Branch, Paper 1990-1, p. 237-246.
- Read, P.B.**  
 1976: Operation Saint Elias, Yukon Territory: pre-Cenozoic volcanic assemblages in the Kluane Ranges; in Report of Activities, Part A; Geological Survey of Canada, Paper 76-1A, p. 187-193.
- Read, P.B. and Monger, J.W.H.**  
 1975: Operation Saint Elias, Yukon Territory: the Mush Lake Group and Permo-Triassic rocks in the Kluane Ranges; in Report of Activities, Part A; Geological Survey of Canada, Paper 75-1A, p. 55-59.
- 1976: Pre-Cenozoic volcanic assemblages of the Kluane and Alsek ranges, southwestern Yukon Territory; Geological Survey of Canada, Open File 381, 96 p.
- Richter, D.H.**  
 1976: Geologic map of the Nabesna quadrangle, Alaska; United States Geological Survey, Miscellaneous Investigations Map I-932.
- Richter, D.H. and Matson, N.A. Jr.**  
 1971: Quaternary faulting in the eastern Alaska Range; Geological Society of America Bulletin, v. 82, p. 1529-1539.
- Richter, D.H., Lanphere, M.A., and Matson, N.A., Jr.**  
 1975: Granitic plutonism and metamorphism, eastern Alaska Range, Alaska; Geological Society of America Bulletin, v. 86, p. 819-829.
- Ridgway, K.D., Cameron, A.R., Sweet, A.R., and Decelles, P.G.**  
 1989: Evolution of a strike-slip basin as defined by sedimentology, palynology and organic petrology, St. Elias Mountains, Yukon Territory, Canada; in Geological Society of America, Abstracts with Programs, v. 21, no. 6, p. A50.
- Schroeter, T.G.**  
 1978: HUM BIRD (114P/10W); in Geological Fieldwork 1977, British Columbia Ministry of Mines and Petroleum Resources, Geological Division, p. 71.
- Schroeter, T.G. and MacIntyre, D.G.**  
 1986: Tatshenshini map-area (114P); in Geological Fieldwork 1985, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Branch, Paper 1986-1, p. 191-196.
- Sharp, R.P.**  
 1943: Geology of the Wolf (Steele) Creek area, St. Elias Range, Yukon Territory, Canada; Geological Society of America Bulletin, v. 54, p. 625-650.
- Sinclair, A.J., Bentzen, A., and McLeod, J.A.**  
 1979: Geology of the White River native copper deposit, Yukon Territory; Indian and Northern Affairs Canada, Publication No. QS-Y0001-000-EE-A1, 27 p.
- Sinclair, W.D. and Gilbert, G.W.**  
 1975: North of 60 - Mineral Industry Report 1973, Yukon Territory; Indian and Northern Affairs Canada, Northern Natural Resources and Environment Branch, Report EGS 1975-7, 177 p.
- Sinclair, W.D., Maloney, J.M., and Craig, D.B.**  
 1975: North of 60 - Mineral Industry Report 1974, Yukon Territory; Indian and Northern Affairs Canada, Northern Natural Resources and Environment Branch, Report EGS 1975-9, 216 p.

- Sinclair, W.D., Morin, J.A., Craig, D.B., and Marchand, M.**  
1976: North of 60 - Mineral Industry Report 1975, Yukon Territory; Indian and Northern Affairs Canada, Northern Natural Resources and Environment Branch, Report EGS 1976-15, 210 p.
- Skinner, R.**  
1961: Mineral industry of Yukon Territory and southwestern District of MacKenzie, 1960; Geological Survey of Canada, Paper 61-23, 52 p.
- 1962: Mineral industry of Yukon Territory and southwestern District of MacKenzie, 1961; Geological Survey of Canada, Paper 62-27, 48 p.
- Souther, J.G.**  
1974: Cordilleran Volcanic Project: Tertiary volcanic rocks of St. Elias Mountains; in Report of Activities, Part A; Geological Survey of Canada, Paper 74-1A, p. 39-41.
- 1977: Volcanism and tectonic environments in the Canadian Cordillera - A second look; Geological Association of Canada Special Paper 16, p. 3-24.
- Souther, J.G. and Stancul, C.**  
1975: Operation Saint Elias, Yukon Territory: Tertiary volcanics; in Report of Activities, Part A; Geological Survey of Canada, Paper 75-1A, p. 63-70.
- Stanley, A.D.**  
1969: Observations of the surge of the Steele Glacier; Canadian Journal of Earth Sciences, v. 6, p. 819-830.
- Stauder, W.**  
1960: The Alaska earthquake of July 10th, 1958: Seismic studies; Bulletin of the Seismological Society of America, v. 50, p. 293-322.
- Stephens, C.D., Lahr, J.C., Fogelman, K.A., and Horner, R.B.**  
1980: The St. Elias, Alaska, earthquake of February 28, 1979: regional recording of aftershocks and short-term, pre-earthquake seismicity; Bulletin of the Seismological Society of America, v. 70, no. 5, p. 1607-1633.
- Still, J.**  
1984: Stratiform massive sulfide deposits of the Mount Henry Clay area, Southeast Alaska; United States Bureau of Mines, Alaska Field Operations Center, Open-File Report 118-84.
- Stover, C.W., Reager, B.G., and Wetmiller, R.J.**  
1980: Intensities and isoseismal map for the St. Elias earthquake of February 28, 1979; Bulletin of the Seismological Society of America, v. 70, no. 5, p. 1635-1649.
- Sturrock, D.L.**  
1975: The Pyroxenite Creek Ultramafic Complex: an Alaska-type ultramafic intrusion in the southwest Yukon; B.Sc. thesis, University of British Columbia, Vancouver, 90 p.
- Sturrock, D.L., Armstrong, R.L., and Maxwell, R.B.**  
1980: Age and Sr isotopic composition of the Pyroxenite Creek Ultramafic Complex, southwestern Yukon Territory: an Alaskan-type ultramafic intrusion; in Current Research, Part B, Geological Survey of Canada, Paper 80-1B, p. 185-188.
- Tarr, R.S. and Martin, L.**  
1912: The earthquakes at Yakutat Bay, Alaska, in September, 1899; United States Geological Survey Professional Paper 69, 135 p.

**Tempelman-Kluit, D.J.**

1974: Reconnaissance geology of Aishihik Lake, Snag and part of Stewart River map-areas, west-central Yukon; Geological Survey of Canada, Paper 73-41, 97 p.

**Thatcher, W. and Plafker, G.**

1977: The Yakutat Bay, Alaska, earthquakes; seismograms and crustal deformation; Geological Society of America, Abstracts with Programs, v. 9, p. 515.

**Theberge, J.B.**

1980: Klavane Pinnacle of the Yukon; J.B. Theberge (ed.), published by Doubleday Canada Limited, Toronto, Canada and Doubleday & Company Inc., Garden City, New York, 175 p.

**Tobin, D. and Sykes, L.**

1968: Seismicity and tectonics of the northeast Pacific Ocean; Journal of Geophysical Research, v. 73, p. 3821-3845.

**Tocher, D.**

1960: The Alaska earthquake of July 10, 1958: Movement on the Fairweather fault and field investigations of southern epicentral region; Seismological Society of America Bulletin, v. 50, p. 267-292.

**Tyrrell, J.B.**

1901: Yukon District; Geological Survey of Canada, Annual Report 1898, v. XI, p. 36-46.

**Warnock, G.F.**

1962: Geology of the Johobo Mines Limited, Yukon Territory; M.S. thesis, University of Arizona, Tucson.

**Watson, K. DeP.**

1948: The Squaw Creek-Rainy Hollow area, northern British Columbia; British Columbia Department of Mines, Bulletin 25, 74 p.

**Wheeler, J.O.**

1963: Kaskawulsh (Mount St. Elias, east half), Yukon Territory; Geological Survey of Canada, Map 1134A.

**Wheeler, J.O. and McFeely, P.**

1987: Tectonic assemblage map of the Canadian Cordillera and adjacent parts of the United States of America; Geological Survey of Canada, Open File 1565.

Wheeler, J.O., Brookfield, A.J., Gabrielse, H., Monger, J.W.H., Tipper, H.W., and Woodsworth, G.J.

1988: Terrane map of the Canadian Cordillera; Geological Survey of Canada, Open File 1894.

**White, G.V.**

1986: Preliminary report O'Connor River gypsum deposit (114P/10E); in Geological Fieldwork 1985, British Columbia Ministry of Energy, Mines, and Petroleum Resources, Geological Branch, Paper 1986-1, p. 279-282.