

DESCRIPTIVE NOTES

PREVIOUS GEOLOGICAL MAPPING
The area was last mapped by Folinsbe (1949; GSC 1:250,000 scale map #977A). There has been no systematic regional mapping in the eastern half of the Lac de Gras sheet since then.

DESCRIPTION OF MAP UNITS
All map units recognized are either Archean metasedimentary and granitoid rocks or younger intrusives (diabase, kimberlite). There is evidence of basement to the above-mentioned units in the map area. Lithologic units in the map legend are described below in inferred stratigraphic order.

Metasedimentary rocks
Greywacke
Green-brown to rusty brown weathering metagreywacke outcrops throughout the map area. Rocks of this unit are dominantly thin-bedded (1-10 cm beds), although thicker beds (to 1 metre) occur. The metagreywackes consist of massive psammite beds and graded psammite - pelite sequences. Bedding is preserved in porphyroblastic schists, but is less common in migmatitic greywackes. Migmatites occur only in smaller rafts and panels (N.B. panel is a non-genetic term to describe narrow outcrops of greywacke which extend along strike) of greywacke within hornblende + biotite tonalites north of Eagle Lake.

Granitoid intrusive rocks
Hornblende + biotite quartz diorite
A large, homogeneous pluton of equigranular, medium-grained quartz diorite occurs at the western margin of the map sheet. Preliminary petrographic studies illustrate that mineral assemblages consist principally of plagioclase + biotite + hornblende + quartz ± epidote. The diorite and epidote are thought to be of magmatic origin. Inclusions of microcline cognate xenoliths are rare or absent in this body. This pluton is weakly deformed, with poor to non-existent foliation development.

Hornblende + biotite tonalite
East of the quartz diorite pluton, an extensive body of deformed tonalite crops out north of Eagle Lake. The hornblende + biotite tonalites are distinguished in the field from the quartz diorites on the basis of lower modal hornblende, higher quartz + biotite contents and the presence of minor K-feldspar. Thin (10 cm - 10 m scale) greywacke septae of variable length (10's - 100's of m) commonly occur in this body; good examples are found along the north shore of Eagle Lake and also in association with the NW/SE trending greywacke panels at the northwest margin of the map area. The small, isolated outcrops of hornblende tonalite northeast of Paul Lake are strongly foliated. This unit typically contains hornblende-rich microcline enclaves. The enclaves are bounded, with long axes lying parallel to S₁.

Biotite tonalite
These rocks are light brown to white weathering, medium-grained and have weak to strong foliations (and are locally lineated). They typically consist of equal proportions of quartz, plagioclase and K-feldspar, with 5-10% of both muscovite and biotite. Aquamarine apatite and tourmaline are common accessories. Garnet, cordierite and sillimanite are also observed, but the latter is invariably found only in association with greywacke inclusions. These granites are fine- to coarse-grained and generally equigranular, but occasionally are weakly porphyritic. A very coarse-grained, K-feldspar-rich, tourmaline-bearing pegmatite phase forms dykes and small stocks up to 250 m in diameter. These pegmatites are widespread in the map area and occur both internal and external to the two mica granite plutons. The two mica granite is observed to intrude greywacke and both tonalite units.

Porphyritic biotite granite
Light red to pinkish-white weathering, medium- to coarse-grained, K-feldspar porphyritic granite forms a distinctive pluton in the northeast part of the map sheet. K-feldspar phenocrysts range from 1-3 cm in length and display a preferred orientation. Rocks consist of 5-10% biotite, with primary muscovite usually absent, or present only in trace amounts (<1%). One large body (consisting of recognizable multiple intrusives) of this unit is found in the region of the Coppermine River and east of Duchesne Lake. Lack of penetrative deformation is consistent with these rocks being emplaced after the main regional metamorphism and deformational event(s).

Diabase dykes
Diabase dykes, ranging in width from 15 to 100 metres are common in the area, and post-date granite intrusions. The dykes are distinguished on the basis of orientation, and are correlated with known dyke swarms in the Slave Province (Fahrig and West, 1986). Four (and possibly five) dyke swarms are represented; all are interpreted to be of Proterozoic age.

McKay (800')
A single McKay dyke, 50 metres wide and trending approximately 080°, is exposed intermittently across the central part of the map sheet. An estimated age of 2.4 Ga (K/Ar whole-rock; Fahrig and West, 1986) for this swarm suggests it is the oldest dyke in the map area.

Contwoyo (045')
Three Contwoyo dykes, 20-40 m wide and trending approximately 045° were noted in the map area. Preliminary U-Pb baddeleyite studies yield an age of 2.23 Ga (A. LeCheminant and O. van Breemen, pers. comm.).

Lac de Gras (190')
Three Lac de Gras dykes, 35-60 m wide, trending approximately 190°, outcrop intermittently in the map area. These dykes are easily distinguished by their well-developed optical texture. The Lac de Gras dykes are observed in the field to cut Contwoyo dykes. Preliminary U-Pb baddeleyite studies yield an age of 2.03 Ga (A. LeCheminant and Heaman, 1989).

Mackenzie (335')
Six dykes of the Mackenzie swarm, with trends of 330°-340° and up to 100 m wide, occur in the map area. Cross-cutting relationships observed in the field suggest the Mackenzie dykes are younger than the Lac de Gras, Contwoyo and McKay dykes, consistent with their U-Pb baddeleyite age of 1.27 Ga (LeCheminant and Heaman, 1989).

'305'
Thin dykes, 15-20 metres wide, trending 300° to 310° can be traced intermittently over a distance of 25 km in the northeast part of the map sheet. The age of this dyke is unknown and its relationship to the other swarms (specifically the Mackenzie) is equivocal.

Kimberlite
Several kimberlite pipes occur in the area. The 'Point Lake' pipe is shown on the map; exact locations of other pipes remain proprietary information at this time. A Rb/Sr three point isochron yielded an Eocene age of 52 ± 1.2 Ma (Northern Miner, 1993), however the exact location of the specific kimberlite dated (other than being in the Dia Mel/BHP claim block) is unknown. Kimberlite-derived mudstones with Cretaceous to Paleocene dinoflagellates, pollen and spores (Northern Miner, 1993) potentially suggest two periods of kimberlite emplacement. The observation that kimberlites in the Lac de Gras area have either a positive or a negative remnant magnetic polarity (Buckie, 1992) is consistent with two periods of kimberlite emplacement.

ARCHEAN INTRUSIVE RELATIONSHIPS & AGE CORRELATIONS
U-Pb geochronology is presently not available for Archean rocks in the Paul Lake map sheet. Relative ages (based on intrusive relationships) are suggested to be metagreywacke > quartz diorite > hornblende + biotite tonalite > biotite tonalite > two mica granite > porphyritic biotite granite. On the basis of mineralogy, preliminary geochemistry and relative degree of deformation, all granitoid rocks in the map area are considered correlative of the 'younger' Slave granitoid suite (King et al., 1992), intruded at ca. 2625-2580 Ma (van Breemen et al., 1992). Further details and tentative correlations can be found in Kjørgaard and Wylie (in press). The metagreywackes in the map area are typical of the metasedimentary rocks of the Yellowknife Supergroup (Henderson, 1970).

STRUCTURAL GEOLOGY
Fabric elements
Fabric elements measured in the field are summarized as follows and described more completely below:
S₁ is primary compositional layering in metagreywacke. It is defined by textural and mineralogical variation at the outcrop scale.
S₂ is the dominant regional cleavage, generally oriented parallel or sub-parallel to S₁ in the metagreywackes.
S₃ is a locally observed, symmetric crenulation cleavage.
F₁ folds are isoclinal. Frequent changes in younging direction of S₃ are observed at the mesoscale and S₃ verges both clockwise and counter-clockwise. Fold closures are not observed due to pervasive lichen cover and shoreline exposures that parallel the regional strike of S₃.
F₂ is a steeply plunging open fold with limbs that extend beyond the limits of the map area, and a wavelength of >20 km. The crenulation cleavage (S₃) is believed to be genetically related to F₂ folding, as it lies parallel to the axial trace.
Structural development
The deformation history of the Paul Lake sheet is polyphase. Tectonic fabrics exhibit varying degrees of development and heterogeneity. S₁ is dominantly steeply dipping, becoming moderate in the F₂ hinge zone. The main cleavage (S₂) is associated with cryptic isoclinal folds (F₁) in metagreywacke. S₂ forms a discrete, spaced cleavage in metagreywackes and is oriented sub-parallel to S₃.
Large, kink-belted porphyroblasts, wrapped by S₃, suggests that D₂ deformation is preceded by static porphyroblastesis of cordierite, andalusite, staurolite and biotite. These porphyroblasts contain straight inclusion trails, suggesting a relict cleavage related to either a previous, but unrecognized folding event, or the early stages of S₁ fabric development. A more locally developed crenulation cleavage (S₃) appears to be genetically related to the large, open fold (F₂) in the map area.
Moderately-dipping stretching lineations trending approximately 020°, C and S fabric in granitoids, local transposition of S₂ and the presence of sheath folds suggest the development of localized high strain zones in the limbs of the F₂ fold. Further evidence can be found in folded pegmatite veins, bedding-parallel shearing and localized ramping of greywackes. Strain partitioning is only observed within the greywackes, close to rheological interfaces with syn-kinematic granitoids.
F₂ Geometry
The main greywacke outcrops lie in a folded belt (F₂). This greywacke belt parallels the northwest arm of Lac de Sauvage, passes through the closure and follows the long axis of Paul Lake. In the Lac de Sauvage limb, the dominant bedding orientation is about 135/70°. In the Paul Lake limb, it is 255/75°. Bedding orientations between Lac du Sauvage and Paul Lake are consistent with ductile folding. The statistical mean fold axis for S₂ trends 008° and plunges 70°. Discontinuous biotite tonalite outcrops of variable size are also found following the trend of the F₂ fold in the lineament along the northwest arm of Lac du Sauvage and between Paul Lake and Eagle Lake. The biotite tonalite outcrops are inferred to have formed a continuous unit before intrusion of two mica granite.

Zones of high strain
Three areas of high strain (Paul Lake; north shore Eagle Lake; northwest arm Lac du Sauvage) are evident from field and structural studies. All occur within or adjacent to areas of metagreywacke outcrop and regional lineaments, but are also observed in the adjacent tonalites.
One, and possibly two generations of high strain structures are observed in the limbs of the regional F₂ fold. The two high strain zones (HSZ) in the northwest part of the map sheet at Eagle Lake and Paul Lake, appear to have an overprinting relationship to one another. The Paul Lake HSZ is associated with transposition of S₂ and development of sheath folds, oriented sub-parallel to the long axis of Paul Lake. The Eagle Lake HSZ is oriented parallel to an asymmetric crenulation cleavage that cuts the sheath folds in the Paul Lake area. The zone itself has a consistent stretching lineation (defined by hb + plag + qtz) oriented at about 020/60°. The regional significance of the overprinting fabric is currently equivocal in the absence of solid kinematic evidence to link the crenulation cleavage with the Eagle Lake HSZ.
A third high strain zone lies in the northwest arm of Lac du Sauvage. It is indicated by progressive strain of porphyroblasts, development of C and S fabrics and shear bands in rocks adjacent to the lineament. Ramping of S₂ and folding of pegmatite veins in greywackes are consistent with ductile kinematics.

METAMORPHISM
In pelitic rocks, there is a subtle variation in mineral assemblages suggesting that metamorphic grade increases slightly northward. Porphyroblasts of biotite, cordierite and andalusite (var. chaoite) are ubiquitous in pelitic layers. Garnet occurrence is variable, but its restriction to narrow, bedding-parallel zones, suggests its occurrence is a function of bulk composition, rather than metamorphic grade. Sillimanite (var. fibrolite) is common throughout the region (typically overprinting the porphyroblasts), as are occasional blades of accreted sillimanite. Metagreywackes are generally sillimanite-bearing in the main contact zones with granitic rocks, whereas those found in narrow panels within tonalite tend to be sillimanite-absent and migmatitic in character. The metamorphic assemblages observed in the greywackes are similar to those observed in other areas of the Slave Province (Thompson, 1978) and are consistent with low-P, high-T metamorphism.

ECONOMIC GEOLOGY
Numerous (> 50) kimberlites have recently been discovered in the Lac de Gras area of the central Slave Province, many diamondiferous. In 7809, the 'Point Lake' pipe has been bulk sampled, with 101 carats recovered from a 180 tonne sample (grade = 0.3 carats/tonne); approximately 25% of the stones are of gem quality, with a few of the stones in the 1-3 carat range (BHP/DiaMet press release, June 1992).

REFERENCES
Buckie, J.
1992: Detection of kimberlite pipes in the Lac de Gras area with helicopter-borne electromagnetics and magnetics; in Brophy, J.A., ed., Exploration Overview 1992, Indian and Northern Affairs Canada, Geological Division, Yellowknife, p.17-18.
Fahrig, W. and West, T.D.
1986: Diabase dyke swarms of the Canadian Shield; Geological Survey of Canada, Map 1627A.
Folinsbe, R.F.
1949: Lac de Gras, District of Mackenzie, Northwest Territories; Geological Survey of Canada, Map 977A.
Henderson, J.B.
1970: Stratigraphy of the Yellowknife Supergroup, Yellowknife Bay-Prosperous Lake area, District of Mackenzie; Geological Survey of Canada, Paper 70-26, 12 p.
King, J.E., Davis, W.J. and Reif, C.
1992: Late Archean tectono-magmatic evolution of the central Slave Province, Northwest Territories; Canadian Journal of Earth Science, v. 29, p. 2156-2170.
Kjørgaard, B.A. and Wylie, R.J.S.
in press: Geology of Paul Lake area, Lac de Gras - Lac du Sauvage region of the central Slave Province, District of Mackenzie, Northwest Territories; in Current Research 1994-C, Geological Survey of Canada.
LeCheminant, A.N. and Heaman, L.M.
1989: Mackenzie igneous events, Canada: Middle Proterozoic hotspot magmatism associated with ocean opening. Earth and Planetary Science Letters, v. 98, p. 38-48.
Northern Miner.
1993: BHP-Dia Met age date kimberlites. Northern Miner, v. 79, #29, p. 1.
Thompson, P.H.
1978: Archean regional metamorphism in the Slave Province - a new perspective on some old rocks; in Fraser, J.A. and Heywood, W.W., eds., Metamorphism of the Canadian Shield; Geological Survey of Canada Paper 78-10, p. 85-102.
van Breemen, O., Davis, W.J. and King, J.E.
1992: Temporal distribution of granitoid plutonic rocks in the Archean Slave Province, northwest Canadian Shield. Canadian Journal of Earth Sciences, v. 29, p. 2186-2199.

ACKNOWLEDGEMENTS
The following are thanked for their assistance: Bedrock mapping, S. Alvarado (1993) K. Buchan, T. LeCheminant, T. Peterson, P. Orr, T. Sagriff (1992), Fieldlog and AutoCad consultation, digitizing assistance, K. Baker, P. Arscott and B. Brodicar, CGD. Digital output using ARC/INFO software, Vic Dohar, Cartography Unit, GICD.

LEGEND

QUATERNARY
Q Undifferentiated glacial sediments



PHANEROZOIC
Kimberlite

PROTEROZOIC
Mackenzie diabase dykes
'305' diabase dyke
Lac de Gras diabase dykes
Contwoyo diabase dykes
McKay diabase dyke

ARCHEAN
A2mg Porphyritic biotite granite
A2mg Muscovite - biotite granite
Abt Biotite tonalite
Ahtb Hornblende - biotite tonalite
Aqd Quartz diorite
Agw Greywacke

Lithological boundary (inferred, assumed)
Bedding (upright, overturned, tops unknown, vertical)
First (main) cleavage (S₁) (inclined, vertical)
Second cleavage
Lineation (intersection, stretching)
Fold axis (F₂)

OPEN FILE 2739
GEOLOGY
PAUL LAKE AREA, LAC DE GRAS
NORTHWEST TERRITORIES
Scale 1:50 000 - Échelle 1/50 000



78 D16	78 D18	78 D13
78 D10	78 D9	78 D12
78 D7	78 D8	78 D6

2739

Geology by B.A. Kjørgaard and R.J.S. Wylie, Geological Survey of Canada
Digital cartography by V. Dohar, Geological Survey of Canada
Part of the Canada - N.W.T. Mineral Initiative Agreement and NATMAP Slave Province Project

OPEN FILE
DOSSIER PUBLIC
2739
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
1993

Recommended citation:
Kjørgaard, B.A. and Wylie, R.J.S.
1993: Geology of the Paul Lake Area (78 D9), Lac de Gras, N.W.T., Geological Survey of Canada, Open File 2739, scale 1:50,000.