GEOLOGICAL MAP OF THE OTTER AND DUCKER LAKES AREA

































Adjoins GSC Open File 2428; Aspler et al. 1992

TABLE 2 GOSSANS

SCHEMA	SAMPLE	LOCATION	DESCRIPTION	METHOD	Au	Ag	Cu	Pb	Zn	Mo	Ni	Co	Mn	Fe	As	Sb	Bi	U
					FA/AA	FA/AA	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm
1	93-I-2-B	65H4a 615967	pillow basalt; rust; fracture	3697	3F/A	3	<1	13	6	3	1	4	13	0.1	<1	<2	<1	<1
2	93-I-2-H2		pillow basalt; rust; fracture; gabbro	3682	4	<1	16	10	7	<1	13	6	40	0.2	<1	<2	<1	<1
3	93-I-2-I1	65H4c 594331	pillow basalt; rust; fracture	3677	3	<1	10	7	5	<1	6	3	20	0.1	<1	<2	<1	<1
4	93-I-2-I	65G1-1 647818	50 m Hurwitz gabbro	3678	5	3F/A	14	7	7	<1	<1	1	10	0.1	<1	<2	<1	<1
5	93-I-2-J	65G1-2 647818	50 m Hurwitz gabbro	3679	3	<1	9	6	5	<1	<1	1	10	0.1	<1	<2	<1	<1
6	93-I-2-K	65G1-3 647818	50 m Hurwitz gabbro; quartz veinlets	3680	3	<1	9	6	5	<1	<1	1	10	0.1	<1	<2	<1	<1
7	93-I-2-N	65H4c 605668	as in Maguse ironstones	3681	<1	<1	17.4	4.8	7	<1	<1	3	10	1.2	0.1	<2	<1	<1
8	93-I-2-O	65H4c 594331	as in Maguse ironstones	3682	<1	<1	4.4	2.5	3	<1	<1	1	10	0.4	<1	<2	<1	<1
9	93-I-2-P	65H3b 594030	sulphides in magnetite IF	3683	<1	3	<1	63.8	39	13	1.2	<1	<1	1.4	0.1	<2	<1	<1
10	93-I-2-Q		sulphides in magnetite IF	3684	<1	3	<1	63.8	39	13	1.2	<1	<1	1.4	0.1	<2	<1	<1
11	93-I-2-R	65H5 615345	pillow basalt; rust; fracture	3685	<1	<1	10	6	6	<1	3	1	10	0.1	<1	<2	<1	<1
12	93-I-2-T	65H4 615345	pillow basalt; rust; fracture	3686	<1	<1	10	6	6	<1	3	1	10	0.1	<1	<2	<1	<1
13	93-I-2-U	65H4 615345	gossan in Hurwitz gabbro	3687	<1	<1	10	6	6	<1	3	1	10	0.1	<1	<2	<1	<1
			50 m Hurwitz gabbro															

UNIT			DESCRIPTION	COMMENT/INTERPRETATION
Mackenzie Diabase			Northeast-trending diabase dykes; non-foliated, unmetamorphosed	1267 ± 3 Ma baddeleyite; LeCheminant and Heaman, 1989
HURWITZ GROUP	GABBRO	HG	Numerous gabbro sills; coarse-grained, pebbly. Local rusty-brown to buff weathering concordant with bedding in adjacent units. Circular to linear zones common. Typically within Ameto Formation, Ducker Fm, and Tavani Formation, rarely in Watterson Formation. Thickest in Ameto Formation and Ducker Fm (1000 m). Thin (<100 m) in Tavani Formation and Watterson Formation. Sills also present in Kinga Formation and Henik Group (Aspler et al., 1993 b).	Evidence of injection pre- Hurwitz Group deformation in form of multiple-injection episodes in isolated zones (Aspler et al., 1993 b). Gabbro sills 2711 ± 0.6 Ma (Heaman and LeCheminant in press); 2171 ± 1.8 Ma (Heaman and LeCheminant in press).
	TAVANI FORMATION	HT3	Mixed sandstone and mudstone with metre-scale interbeds consisting of arkose to mudrock fining-upward sequences	Marine flooding of coastal plain; development of widespread, thin, fine-grained deposits. Interbedded sandstone and mudstone with abundant soft-sediment deformation, because from eustatic collapse.
		HT2	In structural basement north of Ducker Lake, interbedded mondstor laminated and thrombolitic dolostone and arkose to quartz arenite. Thickness ranges from negligible to 3 cm; local sill hapless (see Fig. 3 in Miller and Reading, 1993). Common soft-sediment deformation structures. Stromatolites include: Boxonia, Jacutophyton, Linella, Baicalia, Conophyton, Gymnosolen, Kussiella, Minjaria, Tungussia, Vesicularites.	Shallow marine to subaerial coastal plain
		HT1	Tan and grey weathering siltstone, local parallel and cross-stratified heavy mineral laminae, rare interbeds of grey and pink mudstone breccia	Local lagoons between Tavani coastal plain and Watterson carbonate ramp; local tidal influence; local interbedding with arkose sheetfloods and algal mounds.
	DUCKER FM	HD	Dissentinuous unit with pink to dirty-pink fining-upward cycles of grey and tan arkose to mudrock. Locally, thick beds of arkose with parallel and cross-stratification. Base sharp, commonly with arkose sheetfloods and algal mounds. Thickness changes, where absent. Whatever passes directly and conformably to Tavani.	Intertidal to supratidal carbonate ramp
	WATTERSON FORMATION	HW3	Orange-weathering, recessive laminated dolostone, local m-scale siltstone to mudrock fining-upward cycles	Intertidal to supratidal carbonate ramp
		HW2	Mixed siliciclastic-carbonate unit; arkose to mudrock fining-upward sequences are predominant (typically 80%). Brown massive and microbial laminated dolostone (10%). Green and red Fe-rich mudrock (10%). Local thin (<1 m) beds of pebble-rich arkose. Base of unit at Hurwitz Gabbro contact forms a angle unconformity.	Shallow water-base mass flows alternating with subtidal to supratidal
		HW1	Rose, flesh and orange stromatolitic and microbial laminated dolostone; local mm-scale, fenestral, intraclastic cross, laterally-linked, bulbous to nodular hemispheroids. Mixed carbonate and siliciclastic rocks; arkose to mudrock fining-upward sequences. Local thin (<1 m) beds of pebble-rich arkose; arkose to slate fining-upward sequences interbedded with orange dolomitic laminite.	Intertidal to supratidal carbonate ramp; tidal influence; local interbedding with arkose sheetfloods and algal mounds. Base of unit at Hurwitz Gabbro contact forms a angle unconformity.
	AMETO FM	HA	At base: red, blue, green clate with ramum-scale parallel-oriented arkose. Up-section: arkose to mudrock fining-upward cycles. At top: 0.1-2m- thick dolomitic laminates	Below wave-base sedimentation interrupted by storm-induced sediment gravity flows; local tidal influence; local carbonate ramp
	HENIK GP.	HKw	Discontinuous unit of bedded white chert, magnesian ghost and chert breccia conformably overlying Whiterock Member and Arkosic Formation. Discordant hematitic breccias at base of Whiterock Member.	Previously considered siliciclastic; re-interpreted as chert by Aspler and Chiarenzelli (in press) and Aspler et al. (1992)
		HKm	Supermature quartz arenite. Lower part of section commonly massive, upper 200 m with ubiquitous wave ripples. Parasilicate calcrustions based on comments; winter strata replace indicates depths of 2 cm to 2 m for the entire sequence (Aspler et al., in press).	Tidal - hydrographically-open, tide-free, fresh-water lake or series of lakes
		HKm	Supermature quartz arenite, grey outcrops to quartz arenite, interbeds of white quartz arenite at top. Local trace fossils and cross-stratified. Heavy mineral bands. At base, pebble conglomerate and quartz pebble sandstone. Thickness ranges from 0 to 1320 m.	High energy unchannelled sheet floods draining low relief, stable hinterland to shallow, fresh-water lake or series of lakes; local tidal influence in northern part of Otter Lake
Agd	Granodiorite		Well-foliated granodiorite intrusive into Henik Group. Border zone of Henik Group metamorphosed to amphibolite facies (Aspler et al., 1992).	2711 ± 2 Ma; Rb-Sr whole-rock isochron (Van Breemen and Davis, 1973)
Aga	Gneissic diorite to Granodiorite		Foliated, gneissic diorite to granodiorite. With gneissic banding and lenticular mafic enclaves that engulf granitoid sheetlets.	Deep-level equivalent to Agd, but possibility of being basement to Henik Group cannot be precluded
HENIK GP.	Volcanic/ Sedimentary Sequence	Avs	Green mafic flows to pod-pillowed mafic tuffs and gabbro sills. Avsfl: Whitpor-mouth tuffs; porphyries (in part ±0.1% Ana). Dm-scale graded sandstone to mudstone and mm-scale, graded siltstone to mudstone couplets. Avsm: Mafic to felsic tuff bedded and tuff-bedded tuffaceous siltstone. Sandstone and gravel-bearing mass flow deposits.	Mafic and felsic volcanism; pyroclastic/epiclastic sedimentation. Intervening siliciclastic mass flows with subaerial sedimentation between mass flow pulses.
	Sedimentary Sequence	As	Massive sandstone and dm-m-scale sandstone to mudstone fining-upward sequences	2711 ± 2 Ma; Rb-Sr whole-rock isochron (Van Breemen and Davis, 1973)

MARGINAL NOTES

This open file summarizes the initial results of 1:50,000-scale mapping in the Otter and Ducker lakes area. It extends work south from Bray and Montgomery lakes (Aspler et al., 1992) and represents the continuation of a project designed to better understand Archean and Proterozoic supracrustal rocks in the south-central District of Keewatin (Aspler et al., 1989; 1993 a; b; 1994; Aspler and Bursey, 1990).

STRATIGRAPHY AND SEDIMENTATION

The local stratigraphy of the Archean **Henik Group** (Eade, 1974) part of the Ennadai-Rankin greenstone belt, consists of a siliciclastic turbidite sequence (unit As) and a mixed mafic and felsic volcanic sequence with magnetite iron formation-bearing turbidites (unit **Avs**). Both sequences are intruded by Archean granitic plutons (unit **Agd**). Near the base of unit Avs, mafic sequences are intruded by Archean granitic plutons (unit **Agd**). Near the base of unit Avs, mafic volcanic rocks, in part pillowed, are predominant. Up section are lenses of magnetite iron formation-bearing sandstone to mudstone turbidites and, toward the top, magnetite-jasper iron formation. The number and thickness of iron formation-bearing lenses diminishes southward such that at the Kognak River, iron formation-bearing turbidites form a single 50-metre-thick bed within felsic and mafic tuffs.

The Early Proterozoic **Hurwitz Group** is preserved in a northeast-trending belt of outliers that represent the remnants of a large intracratonic basin thought to have formed due to compressive intraplate stress arising from the pre- ca 2.1 Ga convergence of the Buffalo Head terrane (Ross et al., 1991) with the western margin of the Churchill Province (Fig. 1; Aspler et al., 1992). Lower Hurwitz Group strata are arranged in a conformable onlap sequence of radial basin expansion. The Maguse Member is the lowest unit exposed in the present map area (Table 1). North of the map area, in The Grey Hills, the Maguse Member is about 320 metres thick. It thins southward and pinches out conformably beneath quartz arenites of the Whiterock Member southeast of Otter Lake; further to the west the Maguse Member is represented by rare lenses less than 5 metres thick. Quantitative analysis of wave-ripples within the Whiterock Member indicate average water depths of 2 cm to 2 m over an area of about 100,000 km^2. The Whiterock Member was deposited in a large, hydrographically open, tide-free lake or series of lakes that onlapped fluvial and eolian deposits of the Maguse Member within a low relief continental depression (Aspler et al., in press). Several new exposures of Hawk Hill Member chert and chert breccia, likely sinter deposits (Aspler et al., 1992) were mapped as thin (<10 m) discontinuous lenses above the Whiterock Member east and south of Otter Lake.

The upper Hurwitz Group stratigraphy defines a marine to terrestrial offlap sequence. The abrupt switch from shallow water lacustrine sedimentation of Whiterock Member quartz arenites to deep water mudrock and arkose of the Ameto Formation is attributed to sudden basin deepening and marine waters flooding through gaps in basement arches. With the source of quartz sand and marine waters flooding through gaps in basement arches. With the source of quartz sand and marine waters flooding through gaps in basement arches. With the source of quartz sand