

DESCRIPTIVE NOTES

The Elbow Lake map area (NTS 63K16) comprises 865 km², bounded by latitudes 54° 44' and 55° 02' and longitudes 101° 07' and 100° 30'. It is located within the Early Proterozoic Fin Flats...

Supracrustal rocks in the area belong to the Aniak Group (unit 1) and consist mainly of mafic metavolcanic rocks and related intrusions. Within the Elbow Lake project, these rocks have been subdivided into several units...

Diabase supracrustal rocks (possible Aniak Group affinity) (unit 2) are exposed in the northeast portion of the map area. A large part unit 2 was compiled from McGynn (1993).

A number of large mafic-ultramafic intrusions (unit 4) occur within the map area. The Long Bay, East Claw Lake and Elbow-Claw Lake gabbro complexes (units 4a, 4b and 4c) represent an early mafic to intermediate group (unit 2a) and a mafic to intermediate group (unit 2b).

The East Elbow Lake stock (unit 5a) is an oval intrusion which covers an area of 10 km² just east of the northeast corner of Elbow Lake. Preliminary U-Pb results from this body indicate it is ca. 1850 Ma.

The Elbow Lake tonalite (unit 6) covers an area of 250 km² and is located within the map area. Portions of the Elbow Lake pluton have been described by Baldwin (1980) and Hunt (1970).

The name Webb Lake plutone complex (unit 7) is used to refer to a texturally diverse group of mafic-ultramafic plutonic rocks covering an area of about 35 km² near Webb Lake.

The Big Rat Lake pluton (unit 14) covers an area of about 32 km² on the southwest side of the map area. Dioritic units 14f and 14g are more mafic phases (unit 14a to 14e) and appear to be dykes which intrude the more mafic phases.

The Gaucher Lake pluton (unit 15) is an oval body covering an area of about 30 km² in the north central portion of the map area. It is zoned from a monzonitic to quartz monzonitic rim (unit 15a) to a granitic core (unit 15c).

The one North Star Lake pluton (unit 16) covers an area of 24 km² at the east-central edge of the map area. The western half of this intrusion, which lies in the map area, is zoned from a mafic to intermediate dioritic rim (unit 16a) through quartz diorite and granodiorite (units 16b and 16c) to minor areas of late granite (unit 16d).

The Elbow Lake batholith (unit 8) is an elongate, north-south trending body which extends for over 54 km from beneath the Shield margin to within the Keeseyne belt. Of the 500 km² it covers, over 80% is located in the map area.

Comptonically banded to migmatitic gneisses occur on the northeastern side of the Garts Lake batholith. These have been roughly subdivided into various mappable components. This includes a number of coarse grained amphibole gabbro bodies (unit 8a) which have been described by McGynn (1993).


The northern portion of the Garts Lake batholith consists of a group of probably magmatic mafic phases (unit 8a to 8e). The U-Pb age of 1877 Ma for the Elbow-Claw Lake gabbro complex (unit 8) (Hunt and Whalen, unpublished data) indicates that these phases are among the oldest in the area.

The northeastern portion of the Elbow Lake batholith consists of a group of probably magmatic mafic phases (unit 8a to 8e). The U-Pb age of 1877 Ma for the Elbow-Claw Lake gabbro complex (unit 8) (Hunt and Whalen, unpublished data) indicates that these phases are among the oldest in the area.

REFERENCES

Baltes, A.H. 1980. Geology of the Fin Flats area, Manitoba. Manitoba Mineral Resources Division, Geological Report 79-1.
Baldwin, D.A. 1980. Porphyritic intrusions and related mineralization in the Fin Flats volcanic belt, Manitoba. Energy and Mines, Miscellaneous Report 1990K-3: 230.
Galley, A.G., and Franklin, J.M. 1987. Geological setting of gold, copper, tungsten and molybdenum occurrences in the Phantom Lake region. In Summary of Investigations 1987, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Miscellaneous Report 87-4, p. 115-124.

OPEN FILE DOSSIER PUBLIC 2709 GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA 1993



LEGEND: 15 Tonalite in shear zones and faults. 14a Heterogeneous 'banded' intrusive rocks. 14b Grey, foliated, medium grained, equigranular, hornblende-biotite quartz diorite. 14c Beige to grey, foliated, medium grained, equigranular, hornblende-biotite quartz diorite to granodiorite. 14d Medium pink, foliated, medium grained, porphyritic, hornblende-biotite granodiorite. 14e Dark pink to red, foliated, coarse grained, equigranular to subporphyritic, hornblende-biotite granodiorite. 14f Dark pink, foliated, equigranular, fine to medium grained, biotite +/- hornblende granite. 14g Dark pink, foliated, equigranular, medium to coarse grained, biotite granite. 13a Grey, foliated, fine to medium grained, equigranular, biotite +/- biotite diorite to quartz diorite. 13b Grey to pink, foliated, medium to coarse grained, equigranular, hornblende-biotite quartz diorite to granite. 13c Pink, foliated, fine to medium grained, equigranular, biotite-hornblende granite. 12a Beige to grey, foliated, fine to coarse grained, equigranular, hornblende-biotite monzonite to quartz monzonite. 12b Pink to grey, foliated, medium to coarse grained, equigranular, hornblende-biotite granodiorite. 12c Pink, foliated, medium to coarse grained, K-feldspar phytic, biotite-hornblende granite to granodiorite. 11a Black, massive, medium to coarse grained, pyroxene and quartz gabbro. 11b Grey to black, foliated, medium to coarse grained, equigranular, biotite +/- biotite gabbro to quartz diorite with abundant (1-20%) mafic mineral inclusions, including pyroxene. 11c Grey to beige, foliated, medium to coarse grained, equigranular, hornblende +/- biotite diorite to granite. 11d Pink to grey, foliated, medium to coarse grained, equigranular, hornblende-biotite monzonite to quartz diorite. 11e Grey, massive, foliated porphyritic, hornblende-biotite, granodiorite, minor equigranular coarse grained granite equivalent to unit 11d (see Figure 11d). 10a Grey to beige, well foliated, medium to coarse grained, quartz phytic, biotite granite. 10b Dark grey to pink, well foliated, very fine to fine grained, equigranular, biotite-hornblende granite. 9a Black, medium to coarse grained, foliated, hornblende-biotite +/- quartz diorite. 9b Grey, medium to coarse grained, equigranular, biotite-hornblende, biotite to quartz diorite gneiss with abundant (1-20%) mafic mineral inclusions, including pyroxene, rutile and amphibole gneiss. 9c Grey to grey, fine to coarse grained, quartz porphyritic, biotite granite to granodiorite. 9d Heterogeneous 'banded' intrusive rocks, mixture of foliated, fine to medium grained, quartz biotite to quartz diorite and mafic volcanic rocks in a matrix of, and cut by, foliated mafic mineral inclusions. 9e Heterogeneous 'banded' intrusive rocks, mixture of foliated, fine to medium grained, quartz biotite to quartz diorite and mafic volcanic rocks in a matrix of, and cut by, foliated mafic mineral inclusions. 9f Grey to beige, coarse grained, equigranular, hornblende-biotite quartz diorite to tonalite. 9g Grey to pink, massive to foliated, fine to coarse grained, hornblende-biotite diorite to quartz diorite, coarse grained, followed contacts with biotite to suggest anagmatic intrusion. 9h Grey to beige, coarse grained, equigranular, hornblende-biotite quartz diorite to tonalite. 9i Dark pink to grey, coarse grained, foliated, hornblende-biotite granodiorite to granite, including quartz phytic portions and dykes (1840-2717 Ma; Hunt and Whalen, unpublished data). 9j Dark pink to red, foliated, coarse grained, equigranular to subporphyritic, hornblende-biotite granodiorite. 9k Pink, foliated to mylonitic, biotite granite pegmatite and quartzite. 9l Grey, medium grained, equigranular to plagioclase phytic, biotite +/- hornblende granodiorite to quartz diorite (1877 +/- 3 Ma; Hunt and Whalen, unpublished data). 9m Grey, foliated, fine to coarse grained, equigranular, hornblende-biotite quartz diorite to diorite. 8a Pink to beige, foliated, coarse to very coarse grained, quartz megacrystic biotite-hornblende tonalite. 8b Beige, foliated, coarse to very coarse grained, biotite megacrystic biotite-hornblende tonalite. 8c Beige, foliated, fine to medium grained, quartz phytic biotite-hornblende tonalite (structure 8). 7a Grey, the gabbro, foliated, plagioclase phytic, biotite-hornblende quartz diorite. 7b Beige to pink, slightly to strongly foliated, very fine to medium grained, quartz megacrystic biotite-hornblende tonalite. 7c Beige to pink, slightly to strongly foliated, very fine to medium grained, quartz megacrystic biotite-hornblende tonalite. 7d Pink, foliated, fine to medium grained, quartz phytic, biotite-hornblende tonalite to quartz diorite. 7e Pink to grey, foliated, medium to coarse grained, quartz phytic, biotite-hornblende tonalite to quartz diorite. 7f Pink to grey, foliated, medium to coarse grained, biotite phytic, biotite-hornblende tonalite. 6 Beige to pink, foliated, medium to coarse grained, quartz megacrystic, biotite-hornblende tonalite (1850-257 Ma; Hunt and Whalen, unpublished data). 5 Beige to coarse grained, quartz megacrystic, biotite-hornblende tonalite. 4a As Long Lake gabbro complex. 4b East Claw Lake gabbro complex. 4c Elbow-Claw Lake gabbro complex. 3a Felsic to intermediate intrusions, quartz porphyry, the granitic hornblende, mylonite, quartz diorite. 3b Mafic to intermediate intrusions, diabase, gabbro diorite, pyroxene, mafic gneiss. 2 Mixture of compositionally banded amphibole gabbro and granodiorite and gneissic gneiss or by abundant dykes and sheets of tonalite in granodiorite outcrop-gneiss (equivalent to units 2a and 2c). 1a An assemblage. 1b Ocean floor assemblage. 1c Ocean island basalt assemblage. 1d Undefined. Geological contact (dashed, inferred). Bedding, top known (dotted). Bedding, top unknown (dashed). Pilons, top known (dotted/continuous). Pilons, top unknown (dashed). Flow contact, top known (dotted). Flow contact, top unknown (dashed). Foliation, inferred (generation unknown, 1st, 2nd). Unconformity (top known (dotted/continuous). Unconformity, top unknown (dashed). Fault, minor unknown (dotted, dash). Fault, major unknown (dotted, dash). Normal strike and other zones. Boundary of shear zones.

OPEN FILE 2709 GEOLOGY ELBOW LAKE MANITOBA Scale 1:50 000 - Echelle 1:50 000