### DESCRIPTIVE NOTES

### PREVIOUS MAPPING

**DESCRIPTION OF MAP UNITS** 

Metasedimentary rocks (Agw)

The area (NTS 76D/10, Koala) lies in the central part of the Slave structural province, 315 km NNE of Yellowknife. Folinsbee previously mapped the area on a regional scale in 1949 (GSC 1" to 4 mile scale map

All map units recognized are either Archean metasedimentary and granitoid rocks or younger intrusives developed. hornblende-biotite tonalite, quartz diorite and diorite. The Koala batholith intrudes flanking metasedimentary pluton cooling. rocks (of variable grade), and itself is cut by a two mica granite along its eastern margin. Evidence for basement to the above-mentioned units in the map area is problematic. Small pods and lenses (1 - 10 m scale; not indicated on the map) of highly deformed rocks of intermediate composition observed in the metasediments along the north described below in inferred stratigraphic order.

A variety of metamorphosed sedimentary rocks ('greywackes' consisting of interbedded psammite/pelite) observed. narrow belt along the north shore of Lac de Gras. In these rocks thin, bedding parallel-quartz veins are common. along the northwest shore of Lac de Gras.

Based on criteria from studies of greywackes in the Contwoyto-Itchen Lakes area of the central Slave schists near the north shore of Lac de Gras. province, Bostock (1980) subdivided the metasediments into the Contwoyto and Itchen Formations. Recent work notable for thinner bedding, dominantly millimeter to 10 cm scale (although thicker beds to 50 cm occur) and been emplaced late- to post-D<sub>1</sub>. minor occurrences of graphite-bearing metasediments and beds of mafic and intermediate volcaniclastic rocks. These rock types can only be traced along strike for 10's of metres and their distribution is thus not shown on the High strain zones map. Although banded iron formation was not observed in 76D/10, it occurs to the northeast in sheet 76D/16 (along strike from the greywacke belt north of the Koala batholith). These characteristics suggest the greywackes have affinities with the Contwoyto Formation (as defined by Bostock, 1980).

### Granitoid intrusive rocks Rocks from the Koala batholith have been subdivided into mappable units (diorite/quartz diorite; quartz

diorite/hornblende-biotite tonalite; hornblende-biotite tonalite and biotite granodiorite/hornblende-biotite tonalite) on the basis of mineralogy, presence or absence of cognate xenoliths and variation in fabric development. They are tentatively correlated with the Concession plutonic suite (Bostock, 1980; King et al., 1992; Davis, 1994) in the Contwoyto-Nose Lakes map area to the north. Diorite/quartz diorite (Ad/Aqd)

### Quartz diorite/hornblende-biotite tonalite (Aqd/Ahbt)

A large pluton consisting of multiple intrusions of equigranular, medium-grained rocks dominated by quartz diorite with lesser amounts of hornblende-biotite tonalite occurs in the northeastern part of the Koala batholith. These rocks intrude adjacent metasedimentary rocks. Inclusions of cognate microdiorite or hornblendite consisting principally of plagioclase and hornblende with biotite + quartz + pyrite ± epidote ± interstitial K feldspar. The biotite and epidote are thought to be of magmatic origin. Hornblende + biotite tonalite is distinguished in the METAMORPHISM field from quartz diorite on the basis of lower modal hornblende, higher quartz + biotite contents and the presence of minor K-feldspar.

Hornblende-biotite tonalite (Ahbt)

metasedimentary panel.

#### Biotite granodiorite/hornblende-biotite tonalite (Abgd/Ahbt) Granodiorites of the Koala batholith are dominantly massive, consisting of plagioclase, K-feldspar, quartz,

biotite and ubiquitous accessory pyrite. Texturally distinct phases (e.g. orbicular biotite granodiorite) occuring are observed along the northern contact, suggesting tectonic overprinting subsequent to intrusion. Cleavage

Middle, Lower and Deep samples. Koala bulk sample infor development weakens to the south away from this contact; rocks are generally massive one km from the northern margin of the batholith.

# Two mica granite (A2mg)

A distinctive white to light grey weathered surface and abundant primary muscovite characterizes these granites. Rocks typically consist of equal proportions of quartz, plagioclase and K-feldspar, with 5 - 10% of both muscovite and biotite. Aquamarine apatite and tourmaline are common accessories. Garnet, cordierite and sillimanite are also observed, but the latter is invariably found only in association with greywacke inclusions. The two mica granite body in the southeast part of map sheet is a composite intrusion, consisting dominantly of No data is available for the Leslie and Grizzly pipes. The diamond grade of the DI-2 kimberlite pipe near the north Fabric development consists of one (variably developed) cleavage. This foliation in general persists weakly, sampled displaying localized strain partitioning throughout the pluton. The two mica granite in contact with tonalite has a massive, equigranular fabric. In contrast, a well developed S<sub>1</sub> fabric is observed in the two mica granite where it is in contact with greywacke.

Three swarms of Proterozoic diabase dykes (<1 to 50 m in width) are observed. The dykes cut all Archean units. Rocks from separate dyke swarms are distinguished on the basis of orientation, texture, mineralogy and magnetic characteristics. Dykes have been correlated with known swarms in the Slave Province (Fahrig and West, 1986; LeCheminant, 1994).

# MacKay (080°)

Two MacKay dyke were observed. These dykes are strongly plagioclase phyric. The dyke in the southern Davis, W.J. part of the map sheet is 35 to 45 metres wide and is intermittently exposed across the greywacke belt and two mica granite body on the north shore of Lac de Gras. While this dyke trends at approximately 080°, there are a number of segments which trend at approximately 045°. Two outcrops of a MacKay dyke occur in the central part of the map sheet and range from 1 - 10 m wide, trending approximately 080°. It is unknown if these thin dykes represent splays or satellite dykes from a wider (i.e. 30 - 50 m), but unobserved, MacKay dyke nearby. U-Pb baddeleyite studies on a MacKay dyke yielded an age of 2.21 Ga (LeCheminant and van Breemen, 1994). Lac de Gras (010°)

Two Lac de Gras dykes, 30 - 50 m wide, trending approximately 010°, outcrop intermittently in the map area. They are distinguished from other dykes by their well-developed ophitic texture. U-Pb baddeleyite studies yielded ages of 2.03 to 2.02 Ga (LeCheminant and van Breemen, 1994), correlative with the Booth River intrusive suite 300 km to the NNE. Mackenzie (335°)

#### Segments of at least five dykes of the Mackenzie swarm, with trends of 330°- 340° and up to 50 m wide, were observed in the map area. Previous U-Pb baddeleyite studies on dykes from the Mackenzie swarm yielded an age of 1.27 Ga (LeCheminant and Heaman, 1989).

Several kimberlite pipes occur in the area. The Panda, Koala, Leslie, Grizzly and Fox pipes (BHP/DiaMet ioint venture), along with pipe #DI-2 (DHK/Kennecott joint venture) are shown on the map. Note that all of these kimberlites lie under lakes, therefore pipe outlines as indicated on the map are schematic only and do not indicate

LeCheminant, A.N. and Heaman, L.M. the actual size (this remains proprietary information). Exact locations of other pipes on this map sheet remain proprietary information at this time. A Rb/Sr three point isochron yielded an Eocene age of  $52 \pm 1.2$  Ma (Northern Miner, 1993), however the exact location of the specific kimberlite dated (other than being in the BHP/DiaMet claim block) is unknown. Kimberlite-derived mudstones with Cretaceous to Paleocene dinoflagellate, pollen and spores (Northern Miner, 1993) potentially suggest more than one period of kimberlite emplacement. A U-Pb perovskite age of 86 ± 2 Ma (C.M.H. Jennings, quoted in Pell, 1994) from another kimberlite in the Lac de Gras

#### area (locality unspecified) supports this idea. ARCHEAN INTRUSIVE RELATIONSHIPS & AGE CORRELATIONS

Relative ages (based on intrusive relationships) are considered to be metagreywacke > diorite ≈ quartz diorite ≈ hornblende-biotite tonalite ≈ biotite granodiorite > two mica granite. On the basis of mineralogy, preliminary

Thompson, P.H. eochemistry and relative degree of deformation, all granitoid rocks in the map area are considered correlatives of the 'younger Slave granitoid suite' (King et al., 1992; Davis, 1994), intruded at ca. 2625 - 2580 Ma (van Breemen et al., 1992). Specifically, the diorite - granodiorite suite in the map sheet appears to be equivalent to the Concession suite granitoids found to the north in the Contwoyto-Nose Lakes sheets, dated at 2608 +5/-4 Ma (Davis, 1994). Previous age determinations on two mica granites from the Contwoyto-Nose Lakes sheets are in the range 2599 ± 5 Ma to 2585 ± 4 Ma (summarized in Davis, 1994). Detrital zircon studies on a metagraywacke van Breemen, O. Davis, W.J. and King, J.E. sample from the Paul Lake sheet to the east of the map area give a preliminary maximum deposition age of 2.67 Ga (M.E. Villeneuve, pers. comm., 1994). STRUCTURAL GEOLOGY Fabric elements

#### So is bedding in metasediments and is defined by textural and mineralogical variation at the outcrop scale. Primary sedimentological features observed include graded turbititic beds and scour and fill, ball and pillow and flame structures.

S<sub>1</sub> is the dominant regional cleavage, axial planar to F<sub>1</sub>. In metasediments it is defined mainly by biotite alignment, plus parallel first generation quartz stringers. S<sub>1</sub> is generally oriented parallel or sub-parallel to S<sub>0</sub> in the metasediments, but is strongly refracted across pelitic and psammitic S<sub>0</sub> beds.. S<sub>1</sub> is defined in the diorite/quartz diorite/tonalite/granodiorite suite by mildly

biotite/muscovite + equidimensional quartz/feldspar foliation. S<sub>2</sub> is a locally observed cleavage axial planar to F<sub>2</sub>. S<sub>2</sub> in the greywackes is defined by biotite, muscovite and a Solution of generation of quartz veins. Cleavage development is best observed in pelitic beds. An Solution of the regions Protein 2016 diabase dynas. It. Tryling assistance in interpretation of the regions Protein 2016 diabase dynas. It. Tryling assistance is second generation of quartz veins. Cleavage development is best observed in pelitic beds. An Solution of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the regions Protein 2016 diabase dynas. It. Tryling assistance is thanks go to our expeditor of the region and the region as a second diabase dynas. It is the region as a second diabase dynas. It is the region as a second diabase dynas as a second

### F<sub>1</sub> folds are isoclinal, but are rarely seen. However, when observed, S<sub>0</sub> is transposed parallel to S<sub>1</sub>, yielding isolated F<sub>1</sub> fold closures with attenuated and sheared off limbs. Mesoscopic and and layer-scale F<sub>1</sub> isoclines are

determined by frequent changes of S<sub>0</sub> younging directions and bedding/cleavage (S<sub>0</sub>/S<sub>1</sub>) relationships. F<sub>2</sub> are asymmetric folds; Z folds predominate, S folds are quite rare. L<sub>1</sub> mineral lineations (quartz, biotite) observed in metasedimentary rocks are related to flattening across S1.  $L_2$  intersection lineations, defined by biotite, muscovite and quartz are observed where both  $S_1$  and  $S_2$  are

(diabase, kimberlite). The geology of 76D/10 is dominated by a central granitoid batholith. Co-magmatic rock Rarely observed slicken striae are shallowly plunging. Joint surfaces in all rock types are dominantly steeply types observed (in decreasing order of abundance) in the Koala batholith include: biotite granodiorite, dipping. Sub-horizontal joints (not measured) in granitoid rocks are interpreted to be cracks developed during

#### Tectonic fabrics exhibit varying degrees of development and orientation. The deformational history in the shore of Lac de Gras may possibly represent basement olistrosomes. Lithologic units in the map legend are map area consists of two phases, D<sub>1</sub> and D<sub>2</sub>. Note that in adjacent map areas in the central Slave province (e.g. King et al., 1992, Thompson et al., 1994) which contain the 'older' Slave granitoid suite an earlier deformation event and associated cleavage is observed (S1) and the main or regional cleavage in these areas is the second cleavage (S<sub>2</sub>). The main or regional cleavage in the 76D/10 map area is S<sub>1</sub>, an earlier cleavage has not been

are present in the map area. Grey-green weathering, graded psammite - pelite assemblages of low metamorphic So is dominantly steeply dipping. However, moderately dipping beds are locally observed. The main grade (spotted slates and phyllites) predominate in the northwestern portion of the map sheet as well as in a cleavage (S1) is associated with isoclinal folds (F1), related to D1. S1 forms a discrete, spaced cleavage in the greywackes and is oriented sub-parallel to S<sub>0</sub>. The locally developed S<sub>2</sub> cleavage is associated with asymmetric Adjacent to the northwestern margin of the Koala batholith, the slates/phyllites are highly silicified and quartz

F<sub>2</sub> Z folds (rarely S folds) related to D<sub>2</sub>. In the map area, D<sub>2</sub> is inferred to be genetically related to two different net-veined. South of the Koala batholith, green-brown to rusty brown weathering pelite/psammite assemblages tectonic features. One locality is associated with a high strain zone at the northern margin of the Koala are dominanted by cordierite + andalusite schists. Rare sillimanite (1 - 3 cm porphyroblasts) schists outcrop granodiorite batholith where it is in contact with greywacke. The other area is associated with the tectonic juxtaposition of low grade biotite spotted slates/phyllites against medium/high grade cordierite + andalusite

Porphyroblast development is complex, suggestive of mineral growth pre-, syn- and post-D<sub>1</sub>. At least two by King et al. (1991) has provided additional observations on rocks of these two formations. In the Koala map generations of cordierite are present. Large, idioblastic porphyroblasts of cordierite and andalusite overgrow So sheet, greywackes spatially associated with two-mica granite along the eastern margin of the map area are and are wrapped by S<sub>1</sub> (biotite) suggesting that D<sub>1</sub> deformation is preceded by static porphyroblastesis. However, similar to those in the Paul Lake map sheet to the east (Kjarsgaard and Wyllie, 1993, 1994) and are interpreted cordierite and andalusite porphyroblasts are also observed to contain dextrally rotated non-planar inclusion trails to be an extension of the Paul Lake greywacke belt. These rocks characteristically contain mats of fibrolite plus (syn-D<sub>1</sub>). Cordierite is also seen overgrowing S<sub>0</sub>, S<sub>1</sub> and a biotite mineral lineation as well as being wrapped or minor acicular sillimanite needles plus partial melt. No graphite-bearing assemblages, banded iron formation or cut by S<sub>2</sub> (post-D<sub>1</sub>, pre- to syn-D<sub>2</sub>) Both cordierite and sillimanite are observed to overgrow and alusite; S<sub>2</sub> wraps volcaniclastic sediments are found associated with greywackes in the Paul Lake belt. These characteristics, sillimanite and F<sub>2</sub> crenulates fibrolite, suggesting sillimanite is also post-D<sub>1</sub>, pre- to syn-D<sub>2</sub>. Isograds are along with 1 to 10 cm scale thick beds suggests these greywackes have affinities with the Itchen Formation (as discordant to the main regional cleavage (S<sub>1</sub>). Cleavage development in the Koala batholith is consistent with defined by Bostock, 1980). In contrast, the two large areas of metasediments flanking the Koala batholith are emplacement syn-D<sub>1</sub> and pre-D<sub>2</sub>. Two mica granites (with local cleavage development) are interpreted to have

> Localized areas of high strain were noted during mapping. In metasediments and two mica granite, these are related to D<sub>1</sub>. Strain partitioning related to D<sub>2</sub> was observed along the northern margin of the Koala batholith; here F<sub>2</sub> Z folds and related S<sub>2</sub> cleavage are locally well developed along the contact in both granodiorites and

In the Lac de Gras greywacke belt, low grade slate/psammite or biotite spotted phyllite/psammite are interpreted to have been juxtaposed (dextrally; ± unknown dip-slip component) against medium/high grade cordierite + andalusite schist/psammite. Two lines of evidence were utilized to make this inference. On the south side of the juxtaposition zone, S2 cleavage development becomes more pronounced in the metasediments northwards; similarly the S<sub>2</sub> cleavage becomes much less significant in the schists further northwards away from the juxtaposition zone. S<sub>2</sub> cleavage development is concentrated along the juxtaposition zone. Metamorphic One small pluton consisting of variably deformed (massive to foliated), coarse- to medium-grained diorite grade increases slightly northward from the south-central shore of Lac de Gras to the juxtaposition zone and quartz diorite occurs within the eastern portion of the Koala batholith. These rocks define a weak magnetic (sub-millimetre biotite slates to biotite spotted phyllites). On the north side of the juxtaposition zone the anomaly on the regional (800 m) airborne magnetic survey map (GSC map #7200G). Mineralogy of these rocks metasediments are monotonous cordierite + andalusite schist/psammite. Presently, it is not possible to is dominated by plagioclase and hornblende, with small, but variable amounts of biotite schlieren, pyroxene, determine if metamorphic grade increases to the north or the south in these schists. However, 5 km to the west magnetite, quartz and pyrite. Hornblendite and microdiorite cognate cumulate xenoliths are relatively common. at the margin of the map area, schists along strike from the juxtaposition zone are observed to be rade increases to the south. Consistent with this observation the occurrence of schists on the adjacent southwest shore of Lac de Gras (this study, Folinsbee, 1949).

Evidence for faulting in the map area is limited. Although a number of lineaments are apparent on topographic maps and satellite images, no fault gouge was found in surficial exposures, likely a result of xenoliths are rare or absent in this pluton, in contrast to their occurrence in the diorite/quartz diorite pluton to the subsequent glaciation. However, rare slicken striae were observed. Azimuth measurements of slicken striae southwest. Rocks in this pluton are weakly deformed, appearing slightly recrystallized with poor to non-existent coupled with strike measurements on steeply dipping, and often haematized joint sets suggest faulting may be

In the metasediments, metamorphic grade is quite variable. Biotite grade spotted slates and phyllites northwest of the Koala granodiorite increase in grade to cordierite + andalusite porphyroblastic schists towards the southwest. Adjacent to the north margin of the Koala batholith, the greywackes are hornfelsed and highly silicified, but no new minerals are observed macroscopically. Metasediments north of Lac de Gras increase in Deformed (S<sub>1</sub> ± L<sub>1</sub> fabric) hornblende-biotite tonalite occurs south of the massive quartz grade northwards from biotite grade slates to biotite spotted phyllites. Metagreywackes south of the Koala diorite/hornblende-biotite tonalite pluton. Thin (10 cm - 10 m scale) greywacke septae of variable length (10's - granodiorite are cordierite + andalusite schists; the occurrence of sillimanite porphyroblasts in greywackes along 100's of m) commonly occur in this body, with garnet observed in tonalite adjacent to the metasediments. This the northwest shore of Lac de Gras indicates an increase in metamorphic grade to the southwest; consistent with unit also typically contains hornblende-rich microdiorite enclaves. The enclaves are boudinaged, with long axes observations southwest of Lac de Gras. Biotite, cordierite and andalusite porphyroblasts are well developed in lying parallel to  $S_1$ . These rocks extend east into the Paul lake sheet, north of Eagle lake; the southwestern extent the pelitic layers of these schists. Note that sillimanite in these rocks forms large (1 - 3 cm) porphyroblasts. In of this unit is uncertain due to poor exposure, however, development of the anastomosing biotite plane foliation contrast, metasediments (plus associated partial melt) within two mica granite at the eastern margin of the map weakens and more massive rocks are observed to the south in the vicinity of the southwest/northeast oriented area contain silliminite in the form of fibrolite mats plus rare acicular needles. The metamorphic assemblages observed in the metasediments are similar to those observed in other areas of the Slave Province (Thompson, 1978) and are consistent with low-P, high-T metamorphism.

Numerous (> 100) kimberlites have been discovered since the autumn of 1991 in the Lac de Gras area of within the batholoith are observed in a variety of cross-cutting relationships. Subordinate hornblende-biotite the central Slave Province. A number of the pipes are diamond-bearing. The following data for pipes from the tonalite (massive) also occurs within the area, consistent with the Koala batholith being a multiple intrusive. Poor BHP/DiaMet joint venture within the Koala map sheet is adapted from the DiaMet press release of July 11, 1994. exposure limited attempts to outline individual plutons within the batholith and understand intrusive relationships

A bulk sample of 230 tonnes from the Panda pipe yielded 270 carats for an average grade of 118c/100t; the with the flanking metasedimentary rocks. Lit par lit intrusive relationships with greywacke was observed in one diamonds were valued at US\$127/carat. A bulk sample of 1572 tonnes from the Fox pipe yielded 360 carats for area along the southern contact and at four areas along the northern contact of the batholith. Based on limited an average grade of 23c/100t; no information is available on diamond value. For the Koala pipe, which consists evidence, the southern batholith contact is suggested to be intrusive, however, well developed S<sub>1</sub> ± S<sub>2</sub> foliation(s) of several stacked sub-horizontal phases, the bulk sample was reported as Total sample as well as Upper,

wer and	Deep sample	es. Noala bulk s	ample inform	nation and derived	data are as follows:	
Zone	carats	tonnes ore	grade	value US\$/c	US\$/tonne ore	
U	98	≈332	≈ 29.5	88	26	
M	468	≈362	≈129	116	150	
L	169	≈393	≈ 43	121	52	
D	158	≈101	≈157	91	143	
	Zone U M L	Zone carats U 98 M 468 L 169	Zone     carats     tonnes ore       U     98     ≈332       M     468     ≈362       L     169     ≈393	Zone     carats     tonnes ore     grade       U     98     ≈332     ≈ 29.5       M     468     ≈362     ≈129       L     169     ≈393     ≈ 43	Zone     carats     tonnes ore     grade     value US\$/c       U     98     ≈332     ≈ 29.5     88       M     468     ≈362     ≈129     116       L     169     ≈393     ≈ 43     121	U     98     ≈332     ≈ 29.5     88     26       M     468     ≈362     ≈129     116     150       L     169     ≈393     ≈ 43     121     52

medium- to coarse-grained rocks; less common are fine grained, very coarse grained and porphyritic variants. shore of Lac de Gras on the DHK/Kennecott property is unknown; it is thought to be low as it has not been bulk

The occurrence of metasediments (flanking the Koala batholith), suggested to have affinities with the Contwoyto Formation, coupled with the occurrence of banded iron formation in map area 76D/16 to the northeast (and along strike of metasediments north of the Koala batholith) has potentially important implications for banded iron formation type gold deposits in the map area.

# REFERENCES

King, J.E., Van Nostrand, T., Bethune, K., Wingate, M.T. and Relf, C.

1980: Geology of the Itchen Lake area, District of Mackenzie; Geological Survey of Canada, Memoir 391. 1994: Geochemistry and evolution of Late Archean plutonism and its significance to the tectonic development of the Slave

craton. Precambrian Research, v. 67, p. 207-241. Fahrig, W. and West, T.D. 1986: Diabase dyke swarms of the Canadian Shield: Geological Survey of Canada, Map 1627A.

1949: Lac de Gras, District of Mackenzie, Northwest Territories; Geological Survey of Canada, Map 977A.

1991: Final field report on the Contwoyto-Nose Lakes map area, central Slave Province, District of Mackenzie, N.W.T. in Current Research 1991-C; Geological Survey of Canada, p. 99-108. King, J.E., Davis, W.J. and Relf, C. 1992: Late Archean tectono-magmatic evolution of the central Slave Province, Northwest Territories. Canadian Journal of Earth Science v. 29, p. 2156-2170.

Kjarsgaard, B.A. and Wyllie, R.J.S. 1993: Geology, Paul Lake area, Lac de Gras, Northwest Territories. Geological Survey of Canada Open File 2739, scale 1:50 000. Kjarsgaard, B.A. and Wyllie, R.J.S. 1994: Geology of Paul Lake area, Lac de Gras - lac du Savage region of the central Slave Province, District of Mackenzie,

Northwest Territories. in Current Research 1994-C; Geological Survey of Canada, p. 23-32. LeCheminant, A.N. 1994: Proterozoic diabase dyke swarms, Lac de Gras and Aylmer Lake areas, District of Mackenzie, Northwest Territories; Geological Survey of Canada, Open File 2975, scale 1:250 000

1989: Mackenzie igneous events, Canada: Middle Proterozoic hotspot magmatism associated with ocean opening. Earth and Planetary Science Letters, v. 96, p. 38-48. 1994: U-Pb ages of Proterozoic dyke swarms, Lac de Gras area, N.W.T.: evidence for progressive break-up of an Archean supercontinent. Geological Association of Canada/Mineralogical Association of Canada, Annual Meeting, Program

with Abstracts, v.19, p. A62. 1993: BHP-Dia Met age date kimberlites. Northern Miner, v. 79, #29, p. 1.

U-Pb geochronology (in progress) is presently not available for Archean rocks in the Koala map sheet.

1994: Kimberlites and diamond exploration in the central Slave province, NWT. EGS 1994-7, NWT Geology Division,


Heywood, W.W., eds., Metamorphism of the Canadian Shield: Geological Survey of Canada Paper 78-10, p. 85-102. Thompson, P.H., Ross, D. and Davidson, A. 1994: Regional geology of the Winter Lake-Lac de Gras area, central Slave Province, District of MacKenzie, Nortwest Territories in Current Research 1994-C; Geological Survey of Canada, p. 1-12. 1992: Temporal distribution of granitoid plutonic rocks in the Archean Slave Province, northwest Canadian Shield. Canadian

Journal of Earth Sciences v. 29, p. 2186 - 2199. Ward, B.C., Dredge, L., and Kerr, D. 1994: Surficial geology, Lac de Gras, District of Mackenzie; Geological Survey of Canada, Open File 2928, scale 1:125 000

# **ACKNOWLEDGMENTS**

Additional glacial striae data from Ward et al., 1994. K. Baker and B. Brodaric (CGD) are thanked for FieldLog and AutoCad consultation and general assistance with file preparation for digital map production. Helicopter support in 1994 was provided by Polar Continental Shelf Project. The BHP/DiaMet joint venture are ansatomosing biotite planes ñ aligned microdiorite boudins. A local S<sub>1</sub> fabric in two mica granite is defined by a encountered in the course of mapping. Stimulating discussions with Martin Leake of BHP Minerals provided helpful insights about the geology of the area. Tony LeCheminant graciously provided an early version of his interpretation of the regions Proterozoic diabase dykes. R. Wyllie assisted with a small part of the mapping. Final

3303-33rd Street, N.W., Calgary, Alberta T2L 2A7



Any revisions or additional geological information known to the user would by welcomed by the Geological Survey of Canada Mean magnetic declination 1994, 26° 28' E, decreasing 20.7' annually. Base map digitized from topographic map published at 1:50 000 scale from Surveys, Mapping and Remote Sensing Copies of the topographical edition covering this map may be obtained from the Canada Map Office, Department of Natural Resources Canada, Ottawa, Ontario, K1A 0E9

Geology by B. A. Kjarsgaard, R. Spark and Z.J. Jakop

Digital cartography by M. Sigouin, Geological Survey of Canada

Electrostatic plot produced by the Geological Survey of Canada

Part of the Canada - N.W.T. Mineral Iniative Agreement and

NATMAP Slave Province Project

LEGEND

Biotite granodiorite with minor hornblende-biotite tonalite

Quartz diorite with minor hornblende-blotite tonalite

(c+a) cordiorite + andalusite grade greywacke

SYMBOLS

(s) silliminite grade greywacke

Cordierite + andalusite isograd

Tectonic juxtaposition

L1 (mineral lineation) .

L2 (intersection lineation)

Z fold (second generation).

S fold (second generation).

(ornaments on high temperature side). .

(ornaments on high temperature side). .

Bedding: facing unknown (inclined, vertical).

Glacial striae: direction known, unknown.

Lithological contacts (defined, assumed, inferred)

Second or crenulation cleavage, foliation (inclined, vertical)


Fold axis (fold hinge; first generation, second generation

Axial plane (first generation, second generation).

QUATERNARY

**PHANEROZOIC** 

**PROTEROZOIC** 



a subsidiary agreement under the Canada-Northwest Territories Economic Development Agreement. Project funded by the Geological Survey of Canada

DOSSIER PUBLIC GEOLOGICAL SURVEY OF CANADA COMMISSION GÉOLOGIQUE DU CANADA **OTTAWA** 11/1994

NATIONAL TOPOGRAPHIC SYSTEM REFERENCE AND INDEX

PRELIMINARY GEOLOGY DISTRICT OF MACKENZIE

OPEN FILE 2966

NORTHWEST TERRITORIES Scale 1:50 000 - Échelle 1/50 000

Universal Transverse Mercator Projection Projection transverse universelle de Mercator © Crown copyrights reserved © Droits de la Couronne réservés

This map has been produced from a scanned version of the original map Reproduction par numérisation d'une carte sur papier