


Sources of information for this compilation are geological mapping by C.A. Evenchick in 1991; and airphoto interpretation by C.A. Evenchick. Outline of Hazelton Group outcrop is from Gabrielse, 1977. Notes on the regional nature of map units and on local stratigraphy and structure are in Evenchick and Thorlerson (1993) and Evenchick (1992) respectively. Previous geological map of the region is by Eisbacher (1974); 1:250 000. This map was produced in collaboration with the BC Geological Survey, who provided the digital topographic base through the BC Mineral Potential Program.

**REFERENCES**  
Eisbacher, G.H.  
1974: Sedimentary history and tectonic evolution of the Sustut and Sifton Basins, North-Central British Columbia; Geological Survey of Canada, Paper 73-31, 57 p.  
Evenchick, C.A.  
1992: Bowser Basin facies and map units in southwest Toadogone map area, British Columbia; Geological Survey of Canada, Paper 92-1A, p. 77-84.  
Evenchick, C.A. and Thorlerson, D.J.  
1993: Geology, Spatsizi River, British Columbia (104H); Geological Survey of Canada, Open File 2719, scale: 1:250,000  
Gabrielse, H.  
1997: Geology of Toadogone (NTS4E) and Ware (B4F West Half) map areas, British Columbia; Geological Survey of Canada, Open File 483



OPEN FILE 3517  
GEOLOGY  
**LASLUI LAKES**  
BRITISH COLUMBIA  
Scale 1:50 000 - Échelle 1/50 000

Kilomètres 1 0 1 2 3 4 Kilomètres

Transverse Mercator Projection / Projection transverse de Mercator  
CS 127° 45' / Scale Factor 0.9998, NAD27 / NS C. 127° 45', facteur d'échelle 0.9998, NAD27  
© Crown copyright reserved / © Droits de la Couronne réservés

Published 1997

104H1	B4E12	B4E11
OF 2888		
104H4	B4E15	B4E14
OF 2884	OF 3517	
104H11	B4E14	B4E13
	OF 3518	OF 3516

NATIONAL TOPOGRAPHIC SERIES REFERENCE MAPS TO ACCOMPANY GEOLOGICAL SURVEY OF CANADA MAPS

**OPEN FILE DOSSIER PUBLIC 3517**  
GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA  
OTTAWA  
August 27, 1997