Natural Resources Canada | Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA

NATMAP
CARTNAT
Canada's National Geoscience Mapping Program
Le Programme national de cartographie géoscientifique du Canada

STRATIGRAPHIC SECTIONS


1
LE
TC
C
R
Sandy clayey silt, dark brown. No stratification recognizable. A mammoth molar, found on the surface of this unit yielded a radiocarbon age of 11 220 ± 60 BP (Beta 79781).
Stony diamicton (till). Most sandstone and mudstone clasts disaggregated.
Scattered limestone and Shield stones
Obscured by grading
Mudstone
2
E
A
L
GM
R
C
Stoneless silt. Black chernozem developed within the upper 50 cm
Pebble gravel, 2-30 cm thick. Contains lenses of fine sand
Silt to sandy silt, laminated
Tephra. Ranges from pure bloks of tephra up to 8 cm thick to admixtures of tephra and eolian silty sand (likely Mazama tephra)
Laminated to structureless silt. Scattered pebbles, patchy stringers and pebble/cobble lenses. Pebbles are angular to rounded, locally faceted and striated. Contact with underlying sand gradational laterally for 3 m or more
Laminated silty sand. Abrupt contact with underlying gravel, gradational contact with overlying silts
Cobble gravel, Poorly exposed except for the upper 2 m. All clasts are of montane provenance
Fractured mudstone and sandstone
Apron of sloughed sediments
3
E
TM
LE
GM
R
Stratified eolian sand and silt. Contains a thin tephra band in the middle of the unit (projected thickness from 50 m upstream). Sharp basal contact
Cobble gravel, scattered pebbly sand lenses. Sharp erosional contact with underlying till which it truncates
Diamicton (till), stiff, silty matrix, slightly stony. Has a pronounced vertical prismatic fracture pattern. Clasts are angular and mostly limestone, subordinate sandstone, porphyritic intrusive and quartzite. Contact with underlying gravel sharp. Abrupt contact with underlying sheared and deformed glaciolacustrine sand, silt, and clay. Till fabric indicates ice flow from the northwest
Laminated clayey silt, silty sand and silt fining upwards to clay and silty clay. Entire sequence has been disturbed and folded. Top 50 cm is a melange of gravel, sand, and laminated clay
Sandstone and mudstone, steeply dipping
4
LE
LE
TC
LE
TM
C
TM
LE
GM
GM
R
Slumped glaciolacustrine and eolian sediments including surface soil
Clay and silty clay. Rhythmically bedded. Beds 1-5 cm. Scattered dropstones including quartzite pebbles from the Wollaston Lakes area and pulonitic and metamorphic pebbles from the Canadian Shield
Diamicton (till). Sandy clay silt matrix. Stonier than the underlying diamicton. Pebble lithology same as underlying diamicton except for trace amounts of green argyllite likely from the Burger Group (Waterton Lakes area) and pink plagioclase crystals from the Canadian Shield. Till fabric indicates ice flow from the southeast
Silt and dark grey clay. Extremely indurated, rhythmically laminated, light grey, laminae range from 1 mm to 1 cm. Unit has been tectonized: sheared, thrust faulted, and overturned. Locally, the unit has also been intruded into the overlying diamicton. Slickensides present along some bedding planes
Diamicton (till). Matrix sandy clayey silt. Slightly stony (5-10%). Slightly flasile with an indistinct prismatic fracture. Pebble and bigger clasts are indurated sandstone, limestone, porphyritic intrusive, and quartzite, rounded to angular. Many stones are faceted and striated. White quartzite, and buff, rounded to angular. Many stones are faceted and striated. White quartzite, and buff, rounded to angular. Many stones are faceted and striated. White quartzite, and buff