



**MARGINAL NOTES**  
Introduction

This sheet of nine maps is one of three similar format for the survey area. Grouped together are elements of certain affinities and/or similar distribution patterns. The three sheets are entitled:

- 1) Base Metals and Partition Elements (Open File 3587a)
- 2) Multi-States of Elements with Thorian and Lanthanum (Open File 3587b)
- 3) Alkali Metals, Alkaline Earths, Manganese and Aluminum (Open File 3587c)

The elements depicted on these sheets are either those which show moderate geochemical data relief in the survey area, or they are representative of a closely associated suite of elements (e.g. of the rare-earth elements, only La is shown because the distribution patterns of the other 14 elements, Ce, Eu, Lu, Nd, Sm, Tb, Yb) are almost identical. Listings of concentrations of these and other elements in each sample are given in the diskette released as Open File 3587c. The diskette includes, also, a more detailed description of the methodology, analytical methods and analytical quality control.

**Rationale for Biogeochemical Surveys**

The roots of a single large tree extract elements from many cubic metres of soil, overburden, groundwater and sometimes bedrock. These elements are then transferred to aerial parts of the tree where they may become locally concentrated. In a multi-disciplinary survey program, data derived from the analysis of an appropriate vegetation sample medium permits geochemical mapping, with enhanced background to anomaly contrast of certain elements, which may assist both in mapping bedrock and in the search for concealed zones of mineralization. Because each species of plant has a different requirement for, and tolerance to, a range of chemical elements, some partitioning of elements takes place and there is selective absorption and transference into the plants. For biogeochemical exploration, conifers provide suitable and effective sample media because they are primitive plants that have a wide tolerance to many toxic elements. The outer bark is a repository for many elements that are not required for the metabolic function of the tree.

The geochemical information supplied by vegetation is different from that of soil. Just as two methods of geological survey may provide different information, so may two methods of geochemical survey. A high correlation between distribution patterns of two geochemical sample media is the exception rather than the rule. In geological environments where there is sufficient concentration of metals to form a mineral deposit, such a list of elements may be sufficient to generate biogeochemical anomalies above the mineral source (e.g. by upward diffusion) or close to it (e.g. by groundwater movement or upward movement in electrochemical cells). This, however, usually has geochemical anomalies displaced down-ice from the mineralised source. Such factors need to be taken into consideration when interpreting geochemical results.

**Reconnaissance Survey**

Lodgepole pine is the most common tree species in the Nechako area, and many metals concentrate in its outer bark. Hence, pine bark was selected as the sample medium for a reconnaissance-level biogeochemical survey in the Nechako project area. Samples were obtained by stripping approximately 100 g of outer bark from around the circumference of mature trees. The preferred sample interval was 2 cm along roads, trails, and tracks. To minimise the effects of airborne contamination from roads, samples were collected at least 100 m into the forest. Along lesser used trails and tracks the distance was reduced to approximately 50 m. Within the survey area a two-person crew sampled trees at 268 sites within a 15-day period in July, 1997. Because of the lack of roads and trails in some areas, the sampling grid is not even. However, on average the sample coverage is approximately 1 site per 7 km<sup>2</sup>.

Bark samples were returned to the GSC laboratories in Ottawa where they were air-dried then reduced to ash by controlled ignition at 470°C for 24 hours. A separate 20 g of dry material was ground to a powder and analyzed for mercury. Ash samples were submitted for the analysis of 38 elements by instrumental neutron activation (INAA) and 30 elements by inductively coupled plasma emission spectrometry (ICP-ES) at Activation Laboratories Ltd. (Ancaster, ON). The INAA analysis reports the total concentration of elements in the sample. The ICP-ES is performed on an equal weight digest of the ash, and provides data on the total or near total concentrations of most elements. Data for some elements are obtained both methods, but after review of the data from a quality control standpoint only the more accurate and precise set of values is recorded in the data listing.

**Map Production and Data Presentation**

The proportional dot maps are plotted using the Transverse Mercator projection, with a central meridian of 125°30'. They were generated using AMI (ARC/INFO Macro Language). The macro, with its corresponding input menu, prompts the user to input percentile break points and an appropriate scaling exponent for each element to be mapped. Proportional dots are then generated, using the ARC/INFO SPOTSIZE, POINTSPOT and SPOT commands, with the user specifying an appropriate minimum and maximum dot size. Exponents for individual elements were chosen to provide the best view of the analytical data. Accordingly, care should be exercised when attempting to compare different elements plotted with different exponents. Comments on distribution patterns are given in the digital file in the diskette available as Open File 3587c.

**LEGEND**

UNCONSOLIDATED COVER  
Tertiary and Quaternary Assemblages

Reference: Wilkie, C.P. (comp.) 1997. Geological compilation of the Nechako River (20F) map area, British Columbia, Geological Survey of Canada, Open File 3587, scale 1:250 000.

QUATERNARY  
Pleistocene and Holocene

- Qal Unconsolidated glacial till, fluvial deposits and poorly sorted alluvium

LAYERED BEDS  
Tertiary and Quaternary Assemblages

TERTIARY  
UPPER EOCENE TO OLIгоценE  
Eocene Group  
Massive, vesicular and amygdaloidal varieties of basalt and andesite; minor breccia, tuff, flows of gabbro, neck, plugs and dykes, minor dacite, myelite, conglomerate, sandstone and siltstone

LOWER TO MIDDLE EOCENE  
Ootsa Lake Group  
Buffs a white-colored fine banded myelite and purple-grey; quartz, and biotite phylite phases, fine grained roots with abundant sphaerulites, medium to dark grey phlogopite-biotite phylite, calcareous, biotite with vesicles and amygdaloids of calcite, calcite and chertite; basalt, volcanic conglomerates, sandstone, breccia and tuff

MESozoic  
Cretaceous Group  
Kataikwa Group  
Predominantly hornblende-biotite porphyritic andesite flows and related tuffs, dacite flows, volcanic breccias and gabbro; basalt, also includes lesser hornblende-biotite porphyritic andesite, quartz phylite, biotite, calcite, quartz and ash tuffs, ignimbrite, lava, volcanic breccia and agglomerate, ophiolite assemblage

UPPER TRASSIC AND LOWER JURASSIC  
Tule Group  
Andesite, basalt, breccia, tuff, minor argillite and limestone

MIDDLE JURASSIC TO EARLY CRETACEOUS  
Francois Intrusions  
Jkg Granite and granodiorite  
JKwgn Whether quartz monzonitic, coarse grained monzonitic quartz monzonite with local orthoclase megacrysts, variegated, also called Meander dacite

**SYMBOLS**

Geological boundary (thrust, approximated, assumed) - - - - -  
Fault (suspected/covered by drift) - - - - -  
Road (unclassified) - - - - -

REL. 8301 8302 8303 8304 8305  
8316 8317 8318 8319 8320  
8321 8322 8323 8324 8325  
8326 8327 8328 8329 8330  
8331 8332 8333 8334 8335

**NATMAP CARTMAP**  
Canada's National Cartographic Mapping Program  
Le Programme national de cartographie géoscientifique du Canada

**OPEN FILE PUBLIC**  
3587c  
GEOLOGICAL SURVEY OF CANADA  
COMMISSION GÉOLOGIQUE DU CANADA  
July 1998

Recommended citation:  
Dunn, C.E. and N.L. Hastings  
1998. Biogeochemical Survey of the Ootsa-François Lakes Area, Using Outer Bark of Lodgepole Pine (NTS 93F/13, 14, and part of 12), north-central British Columbia. Alkali Metals, Alkaline Earths, Manganese and Aluminum, Geological Survey of Canada, Open File 3587c, scale 1:250 000.