Natural Resources Canada | Ressources naturelles Canada
Indian and Northern Affairs Canada | Affaires indiennes et du Nord Canada
GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA
NATMAP CARTNAT
Canada's National Geoscience Mapping Program
Le Programme national de cartographie géoscientifique du Canada
Western Churchill Project

DESCRIPTIVE NOTES

INTRODUCTION

In the central part of the Ennadai-Rankin belt, Neoarchean iron formation units of the Henik Group have been mapped from the Dubrin Lake area (ca. 35 km north to the area west of the Henik lakes (Fig. 1; Aspler and Chiarenzelli, 1997 a). New mapping in the Noonut River area (65 H/10 and 11) confirm that these iron formations continue east for an additional 80 km, from Magnet Bay on South Henik Lake across the Noonut River area to Harling Lake. Paleoproterozoic rocks of the Hurwitz Group are intruded into the Henik Group north of Noonut River and Amedy Lake.

Henik Group: **A2mv (A2mvf); Agb; A2f; A2i (A2ii; A2iif; A2iig; A2ips); A2it; A2ipg; A2p**

The Henik Group is the total name for Archean supracrustal rocks in the central part of the Ennadai-Rankin greenstone belt (Eade, 1974). Herein we retain the informal subdivisions outlined in Aspler and Chiarenzelli (1998 a), however new data necessitate revision of provisional stratigraphic correlations (Fig. 3). In the Noonut River area, rocks previously considered part of "unit **A1**" are now thought to be entirely within unit **A2**, a mixed sedimentary-volcanic assemblage consisting of: mafic volcanic rocks (**A2mv**) and related dykes, sills and stocks (**Agb**); felsic to intermediate volcanic rocks (**A2f**); and a variety of sedimentary rocks (**A2i**) including iron formation (**A2it**). Unit **A2i** is a mixed sedimentary-volcaniclastic (**A2ii**) and magnetite-bearing iron formation (**A2iif**). Adjacent to a pluton in the southern part of the map area is a zone of paragneiss (**A2ips**) developed at the expense of unit **A2i**. Unit **A2** subdivides in the Noonut River area are interpreted within the context of a subaqueous back pile - slope - basin depositional model developed for the central part of the Ennadai-Rankin greenstone belt (Aspler et al., 1999). Recent mapping suggests deposition in a single basin that covered an area of at least 6,000 km2.

The lowermost subfacies in the Noonut River area is unit **A2mv**. It consists of predominantly mafic volcanic flows cut by gabbroic dykes and sills. The flows are non-amygdaloidal, thinly bedded and display massive, pillowed, and sheet flood morphologies. Pillow breccia and variolitic horizons are common. Felsic flows (**A2mvf**) form rare interbeds. Exposures of **A2mv** in the northwestern part of 65 I/11 are continuous with rocks previously mapped as part of unit **A1** in the northeast corner of the adjacent sheet, 65 H/12 (Aspler and Chiarenzelli, 1997 a; Aspler et al., 1997). Rocks of **A2mv** are interpreted to underlie the entire area between Amedy Lake and the Henik lakes. In the western part of the area, **A2mv** is overlain by a wedge of felsic to intermediate volcanic flows, tuffs, breccias and agglomerates (**A2f**). The breccias and agglomerates are monomictic, and form massive beds up to 10m thick. In the breccias, angular clasts up to 1m have sharp clast boundaries and are self-supporting; in the agglomerates, amoeboid clasts display mutlimm boundaries and are supported by a crystal-rich matrix.

Elsewhere in the Noonut River area, **A2mv** is overlain by unit **A2i**. Because of the discontinuous nature of outcrop, it is difficult to determine the relationship between **A2mv** and **A2i**. Locally, **A2i** is in contact with rocks of unit **A2mv** without lithologic contacts. Turbidite sandstone/mudstone are (**A2i**) and intermediate to felsic tuffs (**A2iif**) are member subfacies within **A2i**. The turbidite rocks consist of laterally continuous dm-scale sandstone to mudstone fining upward sequences and cm to mm-scale rhythmites. Rarely preserved are vertical profiles arranged in the sequence: sharp base - massive or graded arkose - parallel stratification (+/- ripple drift cross stratification) - siltstone - mudstone. Sedimentary structures indicate paleoflow to the northeast. Unit **A2i** is thickest in the area of the sandstone conglomerate to greywacke unit, and thins to the west. In the eastern Noonut River area, where sandstones form thin garnetoid beds several metres thick (Lyssky sandstone), **A2ips** that are separated by thin mudstone partings. The turbidite rocks also contain local lenses of conglomerate (**A2cg**) with intraformational clasts (mm, iron formation, mafic and felsic volcanic) and m-scale massive or parallel-stratified sandstone interbeds. In the strain zones, the tuff subfacies (**A2iif**) consists of dcm- to m- scale sheets of coarse zoned feldspar arkosite and broken eutrehits and trace biporcelidal quartz. Juvenile feldspar phyric fragments up to 10 cm (a distorted internal laminations) are locally abundant. Rarely, the tuffs contain up to 10% magnetite as fine disseminations and as cm-scale bands. Alteration is pervasive and consists of carbonate-alteration and a penetrative cleavage combines to obliterate primary textures, and the tuffs outcrop as semitic schists.

Iron formation horizons (**A2it**) are interbedded with unit **A2** turbidite and crystal tuff subfacies, both as thick (ca. 300 m) members and as thin (ca. 10 m) isolated layers. Magnetite is concentrated in the pelitic parts of dm-scale fining upward sequences and mm-scale sandstone-mudstone rhythmites, both mm/cm-banded with chert and as the top of thicker (m-scale) beds. Sedimentary structures in the sandstone and mudstone interbedded rhythmites include flow deposits, Bouma Tc-e sequences, and soft-sediment folds, faults, dykes, load casts and flame structures are also common.

Granite, granodiorite, diorite (**Agdi**)

Two felsic plutonic bodies are exposed in the Noonut River area (**Agdi**). In the south, pervasively foliated biotite-hornblende granodiorite-(cut turbidite) rocks, crystal tuffs and iron formation. Both the pluton and a ca. 2.5 km border zone of biotite-plagioclase-hornblende paragneiss (**A2ipg**) contain variably oriented and deformed late-plutonic granitic sheets. Commonly cross-cutting the host fabric, these sheets also contain a foliation concordant to the host rocks and are folded. Although plutonism may have started before, and continued during deformation, late magmatic stages were likely synchronous to post-tectonic.

In the northern part of the area, units **A2mv** and **A2f** are cut by rocks that constitute part of a large composite pluton that extends east into 65 H/2 and north into 65 H/14. The predominant lithology is locally foliated hornblende-biotite granodiorite, but dioritic to gabbroic phases are also represented. Near the margins of the pluton are supracrustal rafts which contain a pervasive internal fabric. However, both supracrustal and plutonic rocks are cut by late-stage granitic dykes that range from undeformed to well foliated. Hence emplacement of the pluton was likely late synkinetic. Along its southern margin, the pluton is unconformably overlain by the Hurwitz Group.

Megacrystic-bearing gabbro (**A2gb**)

Distinctive plagioclase megacryst-bearing northeast-trending gabbro dykes, up to 25 m wide, are sparsely distributed in the Noonut River area (**A-Fgb**). Although volumetrically insignificant, the dykes serve as important markers to distinguish local deformation events and will be valuable for testing structural correlations with other parts of the Ennadai-Rankin greenstone belt. In the south-central part of the area an undeformed dyke cuts cleanly across foliations in the surrounding supracrustal rocks and granitic sheets (Fig. 3). Elsewhere, the dykes are foliated and boudinaged. Aspler and Henik Group strata that were previously filled to new vertical attitudes. However, these dykes also contain a penetrative fabric related to "D_2" (Fig. 3;in Aspler et al., 1998). Thus geochronologic study of the dykes could potentially establish the maximum age of D_2 and the minimum age of D_1. The relationship between the gabbro dykes and the Hurwitz Group is unknown.

Hurwitz Group (**Pvw, Pvr, Pvwm; PVsa; Pvch; Pvw, Pvg**)

North of Noonut River and Amedy Lake, Archean supracrustal and plutonic rocks are unconformably overlain by Paleoproterozoic rocks of the Hurwitz Group. The Hurwitz Group is as ≥ 2.45-2.1 Ga (Heaman and Machado, 1992) and is subdivided into the Wilkinson and Kahochilla basins (Fig. 1). The Noonut River area is situated in an intracratonic basin that occupied the interior of the Hearne Province during the protracted breakup of Nenoland; a speculative Neoarchean/earliest Paleoproterozoic supercontinent (Aspler and Chiarenzelli, 1998). Continental deposits from the lower Hurwitz Group exposed in the map area include: auriferous pyritic quartz pebble conglomerate and subareose to quartz arenite (**Pvw**, Noonut Formation, fluvial); semi-pelite (**Pvr**- Padle Formation, glaciogenic/coastal marine); and a thick succession of quartz arenite, siltstone, and shale (**Pvwm**, Wilkinson Member, coastal marine) (**Pvw**, Wilkinson Member, lacustrine); and chert and chert breccia (**Pvch**, Hawk Hill Member, entire). These rocks are considered to represent the initial sag stage of basin expansion. This was terminated by abrupt basin-centre deepening and drowning, and deposition of mudstone and arkose in frontal (turbidite) of the Amedy Formation (**Pvw**, for details see Aspler et al., 1994; Aspler and Chiarenzelli, 1996 b; 1997 b). Gabbro sills (**Pvgb**) cut, and are folded with, the Hurwitz Group.

STRUCTURAL GEOLOGY

A continuing theme of the Western Churchill NATMAP Project is to separate Archean and Proterozoic thermotectonic events in the Churchill Province (Hoffman, 1988; Lucas et al., 1996). Structural geochronological control is provided by folded and faulted Hurwitz Group. North of Noonut River in 65 H/11, a northward-vergent thrust fault juxtaposes Henik Group above Hurwitz Group. In the footwall of this thrust, Noonut Formation and Wilkinson Member rocks form a panel that is overturned to the northward. On the northeast margin of the panel, the Wilkinson Member is juxtaposed above the Amedy Formation, farther north, the Hurwitz Group defines a gently south-dipping upright succession. A thrusted Wilkinson-Amedy contact, rather than a simple syncline cored by Amedy Formation, is indicated because gabbro sills within the Amedy Formation in the northeast corner of 65 H/10 continue across what would be the trace of a such a syncline.

In addition to the thrust which juxtaposes Henik Group above overturned Hurwitz Group, Paleoproterozoic deformation of Archean rocks is demonstrated by folding of the basement-cover contact and by numerous NW-trending cross faults (likely formed late in the folding history as space accommodating structures) which cut the Hurwitz Group. Paleoproterozoic deformation of Archean rocks and Hurwitz Group deformation is demonstrated by folded clasts in basal Hurwitz Group conglomerates, and by brittle deformation at the Hurwitz-Henik unconformity, where steeply-dipping Henik rocks are overlain by gently-dipping Hurwitz strata.

We recognize three structural generations in Archean rocks. The oldest (D_1) is uniquely identifiable only in the southern part of the area, and is manifested by a penetrative foliation in the granodioritic pluton and in zones of paragneiss (**A2ips** and **A2ig**) which flanks the pluton. A megacryst-bearing gabbro cuts cleanly across well-foliated granodiorite, separating "D_1" from "D_2". Identification of large-scale D_1 structures is uncertain, but relationships between paragneiss units bordering the pluton may indicate D_1 faults. Northwest of the pluton, a zone of paragneiss containing abundant granitic sheets (**A2ipg**) passes abruptly northward to a wedge of paragneiss lacking such sheets (**A2ips**). This wedge is in turn passes abruptly to greenschist-grade pluviolith (**A2f**). The sharp transitions and wedge geometry suggest that the rocks on the north side of the **A2ipg** - **A2ips** contact may be allochthonous. This contact is exposed in the eastern limit of the map area. Elsewhere in the map area, we are generally unable to make the distinction between a D_1 foliation intensified during D_2 and a foliation originating during D_2. Hence foliation measurements are mainly of unknown generation.

Between the zone of paragneiss and Hurwitz Group exposures, the Henik Group is deformed by a series of tight to isoclinal east-northeast-trending folds with steeply-dipping axial surfaces. Although plunging data demonstrate obvious, fold hinges zones are not well-exposed and no new related fabrics or stereographic data to interpret their geometry (see Aspler et al., 1998). We infer that these folds are D_2 structures because of evidence suggesting a previous episode of rifting. In at two localities, megacrystic gabbro dykes cut steeply-dipping Henik Group strata and, in contrast to the undeformed dyke cutting the granodioritic pluton, these dykes contain a foliation, and D_2 limbs of individual folds are inconsistently overturned, and bold hinges plunge steeply. However, we cannot rule out the possibility that some of the folds originated during D_1. Although the folds are tight, they are open in style, and do not display the development of a bedding-parallel cleavage, but because S_0-S_2 overprinting is only rarely preserved, we are generally unable to specify which generation of deformation this cleavage belongs to.

An east-northeast-trending fault extends from South Henik Lake to the eastern limit of mapping. For two reasons, both tenuous, this fault may be Paleoproterozoic. First, it appears to represent the continuation of the Bray Thrust which, north of Montgomery Lake, juxtaposes Archean rocks above the Hurwitz Group (Aspler and Chiarenzelli, 1997 a). Second, the fault is parallel to the Henik lakes, and the Henik lakes are parallel to the Hurwitz Group basin. If the fault is subparallel to thrusts in the Hurwitz Group (see above), although similarity in trend may be coincidence. The fault cuts steeply plunging D_2 folds in iron formation-bearing unit **A2it**. Although similar in trend, we cannot document that initiation of the folds was related to faulting and the cross-cutting relationships formed by fold bisecting during progressive fault-related strain, or if the faulting was entirely younger ("D_3?").

Northeast- and northwest-bending cross-faults cut and deflect D_2 folds in Archean rocks. Locally adjacent to these faults, D_2 folds are tight to isoclinal and are overturned. Fault rocks are 1 to 10 m thick, and are filled with shallowly-dipping axial surfaces. A cleavage related to these folds is only locally developed, but a cleavage-cleavage lineation is common. The cross-faults likely formed to accommodate constrictions related to D_2 folding, similar to those cutting the Hurwitz Group.

RECENT EXPLORATION

Pyritic quartz pebble-rich conglomerates at the base of the Hurwitz Group (Noonut Formation) have been historical targets for gold and uranium in the belt extending from north of Noonut River to Angus Lake (Roscoe, 1981). Previously considered part of the Montgomery Lake Group (see Aspler and Chiarenzelli, 1998 b) the Noonut Formation is now thought to be younger than the Montgomery Lake Group (see Fig. 3; Aspler and Chiarenzelli, 1997 a). Gold is presumably carried in pyritic, considered to be, at least partly, of paleoplacer origin. This interpretation is based on the observation that many of the pebbles are pyrite-chert composites indistinguishable from rocks mapped in the Henik Group.

Recent exploration in the Henik Group by Comaplex Minerals Corp. has identified several quartz-carbonate vein associated gold prospects (Table 1). At "Esker", gabbroic bodies cut, and are folded with, felsic rocks (quartz-feldspar porphyry) and iron formation. Mineralization is hosted by quartz-carbonate veins, associated with pervasive carbonate-pyrite alteration, principally in the gabbros near lithologic contacts. The veins contain a late, moderately southwest-plunging mineral lineation (Bonus quartz and iron carbonate) that is found only in the Esker area. The best mineralization is in hinge zones of open to tight southwest-plunging folds. Nearby pods (D_2) cm-D_3 megacrystic-bearing gabbros display pervasive sericitic alteration and a concordant penetrative fabric.

The "Ironside" and "Napartok" prospects are along the ENE-trending fault (D_2?) that extends across the center of the map area. Mineralization is hosted by quartz-carbonate veins, and by pyrite-chert rocks that are locally interbedded with turbiditic semi-pelite (c. felsic tuff). Adjacent to well-mineralized veins, magnetite is replaced by a chlorite-pyrite-arsenopyrite assemblage. At Napartok, gold is in a subvertical zone of quartz-veined iron carbonate-pyrite-albite-sulphide (pyrrhotite + pyrite + arsenopyrite) schist at the contact between variolitic mafic volcanic rocks and semi-pelite. Local preservation of variolites suggests derivation of the schist from mafic volcanic rocks. This lenses is cut by a late, steeply-plunging, quartz-feldspar porphyry dyke. Although the dyke is not mineralized, it is notable that these eighties-fiolden along strike (+ 30-35 to the NE and ESE and are responsible for the weak magnetic linear features evident in the Napartok area (Fig. 2). In core samples, pyrrhotite-pseudomorphs pyrite, and commonly displays pressure shadow textures. At "River", auriferous quartz-carbonate-biotite veins, associated with pyrrhotite and pyrite and rare massive pyrite seams, occur in magnetite iron formation intervals interbedded with turbidite semi-pelites. The veins are associated with a strong biotite-albite-pyrite alteration and shallowly SW-plunging cascade folds that overprint moderately west-plunging tight to isoclinal D_2 F_2 folds.

In summary, recently discovered gold prospects in the Henik Group are associated with zones of iron carbonate-sulphide alteration and quartz-carbonate-sulphide veins. Concentrations of alteration and veining occur within geochemically favoured competent host rocks, within or adjacent to D_2 or D_3 structures. Dating of megacrystbearing gabbros can test if these prospects are examples of Proterozoic gold in Archean rocks.

REFERENCES

Aspler, L.B., and Chiarenzelli, J.R.
1998 Sedimentology, sedimentology and physical volcanology of the Ennadai-Rankin greenstone belt, Northwest Territories, Canada: Late Archean paleogeography of the Hearne Province and tectonic implications; Precambrian Research, v. 77, p. 59-93

1998b Relationship between the Montgomery Lake and Hurwitz Group, and stratigraphic revision of the lower Hurwitz Group, District of Keewatin; Canadian Journal of Earth Sciences, v. 33, p. 1243-1255.

1997a Archean and Proterozoic Geology of the North Henik Lake area, District of Keewatin, Northwest Territories; g) Current Research 1997-C; Geological Survey of Canada, p. 145-155.

1997b Initiation of ca. 2.45-2.1 Ga intracratonic basin sedimentation of the Hurwitz Group, Keewatin Hinterland, Northwest Territories, Canada; Precambrian Research, v. 81, p. 269-287.

1996 Two Neoarchean supercontinents ?: evidence from the Paleoproterozoic; Sedimentary Geology, v. 120, p. 75-104.

Aspler, L.B., Barham, B.A., and Chiarenzelli, J.R.
1998 Results of integrated geologic and aeromagnetic mapping, and recent exploration, Henik and Hurwitz groups, Noonut River, Northwest Territories (Nunavut) g) Current Research 1998-C; Geological Survey of Canada, p. 107-199.

Aspler, L.B., Chiarenzelli, J. R., and Cousens, B.L.
1997 Geology of the Ofterbit Lake area, District of Keewatin, Northwest Territories (part of 65 G/8 and 65 H/ 12); Geological Survey of Canada Open File 3546, scale 1:50,000

Aspler, L.B., Chiarenzelli, J. R., and Bursey, T.L.
1994 Ripple marks in quartz arenites of the Hurwitz Group, Northwest Territories, Canada: Evidence for wellmixed wave in a vast, Early Proterozoic, shallow, fresh-water lake: Journal of Sedimentary Research, v. 64 A, p. 302-306.

Eade, K.E.
1974 Geology of Kognak River area, District of Keewatin, Northwest Territories; Geological Survey of Canada Memoir 377, 66p.

Heaman, L.M.
1994 2.45 Ga global mafic magmatism: Earth's oldest superplume ? p; Abstracts of the Eighth International Conference on Geochronology, Cosmochronology and Isotope Geology; M.A. Lanphere, G.B. Dalrymple, B.D. Turrin (eds.); United States Geological Survey Circular 1107, p.132.

Heaman, L.M., and LeCheminant, A.N.
1993 Paragenesis and U-Pb systematics of baddeleyite (ZrO2); Chemical Geology, v. 110, p. 95-126.
LeCheminant, A.N. and Heaman, L.M.

1989 Mackenzie igneous events, Canada: Middle Proterozoic hotspot magmatism associated with ocean opening; Earth and Planetary Science Letters, v.96, p.38-48.

Roscoe, S.M.
1981 Temporal and other factors affecting deposition of uraniferous conglomerates; g) Genesis of Uranium- and Gold-Bearing Conglomerates, P.C. Armstrong (ed.); United States Geological Survey Professional Paper 1161-A-BB, p.941-971.

Figure 1. Simplified geology of Hearne Domain and location of study area



OPEN FILE 3760
EGS 1999-3

GEOLOGY

NOOMUT RIVER AREA

NUNAVUT

Scale 1:50 000 - Échelle 1/50 000

Kilometres 1 0 1 2 3 4 Kilometres

Universal Transverse Mercator Projection
Projection transverse universelle de Mercator
*Her Majesty the Queen in Right of Canada, 1999
*Sa Majesté la Reine du chef du Canada, 1999

Geology by L.B. Aspler, J.R. Chiarenzelli and B.A. Barham, 1999

Geological compilation by L.B. Aspler, 1999

Digital cartography by R.L. Allard, Geoscience Information Division

Electrostatic plot produced by the Geoscience Information Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by the Geoscience Information Division

Mean magnetic declination 1999, 27° 49' E, decreasing 8.4' annually

LEGEND

MESOPROTEROZOIC

NANDINEMI DABASE

Pnd — Northwest-trending diabase dykes: iron-foliated, unmetamorphosed. 1260 +/- 2 Ma baddeleyite (LeCheminant and Heaman, 1989)

--- INTRUSIVE CONTACT ---

PALEOPROTEROZOIC

HURWITZ GROUP

Pmgb — HURWITZ GABBRO: Diacontinuous gabbro sills; coarse-grained, poikilitic. Local crude layering concordant with bedding in adjacent units. Chloritic shear zones common. 2111 +/- 0.6 Ma (Heaman and LeCheminant 1993). Pre-folding of Hurwitz Group

Pmk — AMETOKNOK FORMATION: Dark-coloured mudstone, siltstone and very fine-grained sandstone; sharp-based fining-upward sequences and rhythmites. Local stratiform greywacke at top

KINGIA FORMATION

Pwch — HAWK HILL MEMBER: Discontinuous unit of bedded white chert, Hanson chert and chert breccia conformably between the Wilkinson Member and Amedy Formation. Discordant hematite breccias at top of Wilkinson Member

Pwwm — WILKINSON MEMBER: Supermature quartz arenite. Lower part of section commonly massive, upper 200 m with ubiquitous wave ripples. Paleolift calculations based on symmetric, relict orbital ripples indicates depths of 2 cm to 3 m for the entire sequence (Aspler et al., 1994)

Pwm — MAGUSE MEMBER: Maroon, pink, grey subareose to quartz arenite; fining upward sequences, locally channelized. Base of member locally cross-stratified. Interbeds of red sandstone, siltstone, and grit with clasts of well rounded spherical white quartz, jasper, blue and grey chert

Pvr — PADLE FORMATION: Massive mottle & stratified polymictic conglomerate, arkose; minor iron-oxide sandstone; siltstone-mudstone rhythmites with dropstones; 56 pellets; possible kaole pseudomorphs

Pvw — NOONUT FORMATION: Subareose to quartz arenite with interbeds of quartz pebble conglomerate, in part pyritic

--- UNCONFORMITY ---

PALEOPROTEROZOIC or ARCHEAN

P-Agb — Megacrystic gabbro dykes: Northeast trending; with feldspar phenocrysts to 5 cm (locally agglomeritic); Foliated and non-foliated

--- INTRUSIVE CONTACT ---

ARCHEAN

Agdi — Well-foliated granodiorite, granite, diorite

--- INTRUSIVE CONTACT ---

HENIK GROUP

A2 — Mixed sedimentary-volcanic sequence

A2ipg — Biotite-plagioclase paragneiss developed at expense of turbiditic semi-pelite and felsic tuff protolith. **A2ipg**, with abundant variably deformed granitic sheets

A2i — Interfingering turbiditic sandstone/mudstone sets, felsic to intermediate tuffs; mafic volcanic rocks and volcaniclastic rocks. **A2i** predominantly turbiditic sandstone to mudstone fining-upward sequences and rhythmites; **A2iif**, predominantly felsic to intermediate crystal and pebbly tuff; **A2iig**, with lenses of conglomerate; **A2ii**, predominantly mafic with lenses of pebbly sandstone

A2it — Magnetite iron formation; **A2it**', defined on the basis of aeromagnetic data

A2f — Felsic to intermediate flows; tuffs; pebbly tuffs; monolithic volcanic breccia and agglomerate

Agb — Gabbroic dykes, sills and stocks; well foliated

A2mv — Mafic flows, locally pillowed; local pyritic chert layers; **A2mvf**, rarely with felsic volcanic layers

Geological boundary (defined, approximate)

Fault (defined, approximate)

Thrust fault (approximate)

Axial trace (anticline, synicline)

Outcrop

Frost heave; probable subjacent outcrop

Bedding (upright, overturned, vertical, tops unknown)

Bedding from pillow lavas (upright, overturned, tops unknown)

Cleavage, Hurwitz Group (incl/ned)

Foliation, Henik Group and related plutonic rocks, likely in part Paleoproterozoic (inclined, vertical)

Minor Fold (early generation, later generation)

Stretching lineation

Gold

Name	UTM E/N Zone 14 NAD 27	Occurrence Type/Host	Associated Minerals	Selected Sampling/ Drilling Results
Esker	592370 6629440	Quartz-iron carbonate veins in gabbro, quartz-feldspar porphyry, lesser quartz- carbonate veins in quartz-carbonate-sericite schist	Sulphides: Pyrite, lesser arsenopyrite, chalcopyrite, minor chalcocite, molybdenite Other: Iron carbonate, muscovite, albite, green mica, tourmaline, titanite	Surface samples to 55.5 gpt Au DDH97-13: 2.35 gpt Au over 70.95 m with 6.45 gpt Au over 12.9 m; 12.22 gpt Au over 2.5 m DDH97-15: 8.16 gpt Au over 13.27 m DDH97-23: 24 gpt Au over 5.0 m
Ironside	600800 6627425	Quartz-chlorite-carbonate veins in gabbro, quartz-feldspar porphyry and chlorite-sulphide replacement zones in oxide iron formation	Sulphides: Pyrite, arsenopyrite, lesser pyrrhotite, chalcopyrite Other: Iron carbonate, chlorite, biotite, calcite	Surface samples to 116 gpt Au DDH96-1: 8.78 gpt Au over 3.0 m Surface samples to 3.7 gpt Au
Napartok	594430 6623870	Quartz-iron carbonate veins, disseminated sulphides in iron carbonate-sericite-albite schist at mafic volcanic-sediment contact	Sulphides: Pyrrhotite, lesser pyrite, arsenopyrite Other: Iron carbonate, albite, sericite, muscovite, chlorite, stratoid	DDH96-6: 18.3 gpt Au over 3.8 m DDH96-7: 9.2 gpt Au over 2.5 m
River	630000 6637050	Quartz- carbonate biotite veins, disseminated sulphides in chlorite schist; massive pyrite seams in argillaceous chert-pyrr ite ironformation	Sulphides: Pyrite, pyrrhotite Other: Biotite, carbonate	Surface samples to 84 gpt Au

Table 1. Recent gold prospects, Comaplex Minerals Corp., Noonut River area

OPEN FILE DOSSIER PUBLIC
3760
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
OTTAWA
07/99

EGS 1999-3

Recommended citation:
Aspler, L.B., Chiarenzelli, J.R. and Barham, B.A.
1999: Geology of the Noonut River area, Open File 3760, scale 1:50 000
Geological Survey of Canada, Open File 3760, scale 1:50 000

Canada