

Geological Survey
of Canada



Commission géologique
du Canada

**SCOTIAN SHELF
STRUCTURE AND ISOPACH 11
ISOPACH MAP OF
WYANDOT FORMATION
TO PETREL MARKER**

**PLATE-FORME NÉO-ÉCOSSAISE
STRUCTURE ET ISOPAQUES**

**CARTE ISOPAQUE, DE LA
FORMATION DE WYANDOT JUSQU'AU
REPÈRE DE PETREL**

B. C. MacLean

Recommended citation: MacLean, B. C., 1991: Structure and isopach 11: isopach map of Wyandot Formation to Petrel marker; *in* East Coast Basin Atlas Series: Scotian Shelf; Atlantic Geoscience Centre, Geological Survey of Canada, p. 95.

Notation bibliographique conseillée: MacLean, B. C., 1991: Structure et isopaques 11: carte isopaque, de la Formation de Wyandot jusqu'au repère de Petrel; dans Série des atlas des bassins de la côte Est: plate-forme Néo-Écossaise; Centre géoscientifique de l'Atlantique, Commission géologique du Canada, p. 95.

Additional copies of this map may be obtained from the Geological Survey of Canada, Atlantic Geoscience Centre, P. O. Box 1006, Dartmouth, Nova Scotia B2Y 4A2 Canada (Ph: 902-426-2773; FAX: 902-426-4266).

Des copies supplémentaires de la carte peuvent être obtenues auprès de la Commission géologique du Canada, Centre géoscientifique de l'Atlantique, case postale 1006, Dartmouth (Nouvelle-Écosse) B2Y 4A2 Canada tél (902) 426-2773, facsimilé (902) 426-4266.

This map was generated by subtracting the Wyandot Formation depth contours (see map sheet Structure and Isopach 6, this volume) from those of the Petrel marker (see map sheet Structure and Isopach 5, this volume) and deleting all faults. The interval includes the upper part of the Dawson Canyon Formation and the Wyandot Formation. It is marginally thicker along a central southwest-northeast axis and is thickest in the southeast corner. Four salt diapirs penetrate this interval.

SELECTED BIBLIOGRAPHY

Ascoll, P.

1976: Foraminiferal and ostracod biostratigraphy of the Mesozoic-Cenozoic, Scotian Shelf, Atlantic Canada; *in* First International Symposium on Benthonic Foraminifera of Continental Margins, Part B, Paleocology and Biostratigraphy, (ed.) C. T. Schafer and B. R. Pelletier; Maritime Sediments, Special Publication No. 1, p. 653-771.

Barss, M. S., Bujak, J. P., and Williams, G. L.

1979: Palynological zonation and correlation of sixty-seven wells, eastern Canada; Geological Survey of Canada, Paper 78-24, 118 p.

Given, M. M.

1977: Mesozoic and early Cenozoic geology of offshore Nova Scotia; Bulletin of Canadian Petroleum Geology, v. 25, p. 63-91.

Jansa, L. F. and Wade, J. A.

1975: Geology of the continental margin off Nova Scotia and Newfoundland; *in* Offshore Geology of Eastern Canada, Volume 2, Regional Geology, (ed.) W. J. M. van der Linden and J. A. Wade; Geological Survey of Canada, Paper 74-30, v. 2, p. 51-106.

Mclver, N. L.

1972: Mesozoic and Cenozoic stratigraphy of the Nova Scotia shelf; Canadian Journal of Earth Sciences, v. 9, p. 54-70.

Wade, J. A. and MacLean, B. C.

1990: The geology of the southeastern margin of Canada, part 2: aspects of the geology of the Scotian Basin from recent seismic and well data; *in* Geology of the Continental Margin of Eastern Canada, (ed.) M. J. Keen and G. L. Williams; Geological Survey of Canada, Geology of Canada, no. 2, p. 190-238 (also Geological Society of America, The Geology of North America, v. I-1).

La présente carte a été obtenue en soustrayant les isobathes de la Formation de Wyandot (voir la carte Structure et isopaques 6 du présent volume) de celles du marqueur (repère) de Petrel (voir la carte Structure et isopaques 5 du présent volume) et en omettant toutes les failles. L'intervalle comprend la partie supérieure des formations de Dawson Canyon et de Wyandot. Elle est légèrement plus épaisse le long d'un axe central sud-ouest-nord-est et atteint son épaisseur maximale dans le coin sud-est. Quatre diapirs de sel pénètrent dans cet intervalle.

BIBLIOGRAPHIE CHOISIE

Ascoll, P.

1976: Foraminiferal and ostracod biostratigraphy of the Mesozoic-Cenozoic, Scotian Shelf, Atlantic Canada; *in* First International Symposium on Benthonic Foraminifera of Continental Margins, Part B, Paleocology and Biostratigraphy, (ed.) C. T. Schafer and B. R. Pelletier; Maritime Sediments, Special Publication No. 1, p. 653-771.

Barss, M. S., Bujak, J. P., et Williams, G. L.

1979: Palynological zonation and correlation of sixty-seven wells, eastern Canada; Geological Survey of Canada, Paper 78-24, 118 p.

Given, M. M.

1977: Mesozoic and early Cenozoic geology of offshore Nova Scotia; Bulletin of Canadian Petroleum Geology, v. 25, p. 63-91.

Jansa, L. F. et Wade, J. A.

1975: Geology of the continental margin off Nova Scotia and Newfoundland; *in* Offshore Geology of Eastern Canada, Volume 2, Regional Geology, (ed.) W. J. M. van der Linden and J. A. Wade; Geological Survey of Canada, Paper 74-30, v. 2, p. 51-106.

Mclver, N. L.

1972: Mesozoic and Cenozoic stratigraphy of the Nova Scotia shelf; Canadian Journal of Earth Sciences, v. 9, p. 54-70.

Wade, J. A. et MacLean, B. C.

1990: The geology of the southeastern margin of Canada, part 2: aspects of the geology of the Scotian Basin from recent seismic and well data; *in* Geology of the Continental Margin of Eastern Canada, (ed.) M. J. Keen and G. L. Williams; Geological Survey of Canada, Geology of Canada, no. 2, p. 190-238 (also Geological Society of America, The Geology of North America, v. I-1).

Publication de la Commission géologique du Canada également disponible en français