



Map 3: Eastern Grand Banks, S.E. Newfoundland Slope

Acoustic Seabed Sediment Classification

- debris flow
- high amplitude reflector
- poorly stratified
- poorly stratified eroded
- prolonged strong reflector
- smooth debris flow
- transparent reflector
- well-stratified
- well-stratified eroded

Probable Gas Occurrences

- gm
- sr,bs,wo
- sr,er
- sr,wo
- wo
- wo,bs
- wo,er
- wo,er,bs
- wo,sr,gm

Bathymetry (metres)

- 0
- 200
- 500
- 1000
- 1500
- 2000
- 2500
- 3000
- 3500
- 4000
- 4500
- 5000

Legend:

- bs=brightspot(s)
- er=enhanced reflector(s)
- gm=gas masking
- wo=wipe-out(s)
- sr=subvertical reflector(s)

Scale: 20 0 20 40 Kilometers

Projection: Lambert Conformal Conic Projection, Standard Parallels 45 N and 66 N

Map created by: Gordon Cameron

recommended citation: Cameron, G. D. M., 1999. Acoustic evidence for shallow gas and seabed classification along selected ship tracks on the deep-water margin of the Grand Banks of Newfoundland. Map 3 of 9, G.S.C. open file report 3762.