

Copies of this map may be obtained from the Geological Survey of Canada:  
601 Booth Street, Ottawa, Ontario K1A 0E8  
3305-33rd Street, N.W., Calgary, Alberta T2L 2A7  
151-665 Robinson Street, Vancouver, B.C. V6B 5A3




MAP 1987A  
SURFICIAL GEOLOGY  
**MANSON RIVER**  
BRITISH COLUMBIA

Scale 1:250 000/Échelle 1/250 000

kilometres 5 0 5 10 15 20 kilometres

Universal Transverse Mercator Projection  
North American Datum 1927  
© Her Majesty the Queen in Right of Canada, 2000

Projection transversale universelle de Mercator  
Système de référence géodésique nord-américain, 1927  
© Sa Majesté la Reine du chef du Canada, 2000



LEGEND

Coloured legend blocks indicate map units that appear on this map  
This legend is common to maps 1986A and 1987A

QUATERNARY

POST FRASER GLACIATION

NONGLACIAL ENVIRONMENT

X

**ANTHROPOGENIC DEPOSITS:** rubble, diamicton, gravel, sand and/or clay emplaced by human activity; 1 to 10 m thick; generally form flat surfaces; primarily occur in the vicinity of mine sites

O

**ORGANIC DEPOSITS:** peat and muck; 1 to 10 m thick (typically 2 to 3 m); form fens and bogs; organic deposits too small to be shown at this scale occur within other units; common within abandoned meltwater channels

Ap

**ALLUVIAL (FLUVIAL) DEPOSITS:** gravel and sand with minor silt and clay, deposited by streams; commonly stratified; generally well sorted except in alluvial fans

**Floodplain sediments:** sand and silt, commonly include organic materials and in many places underlain by gravel; 1 to 3 m thick; occur as flat surfaces close to river level; prone to flooding

At

**Terrace sediments:** stratified sand and gravel overlain by a veneer of sand and silt; 2 to 10 m thick; form terraces well above flood level

Ad

**Deltaic sediments:** stratified sand and gravel underlain by silt and clay; 2 to 5 m thick on average; occur at the mouth of streams entering lakes

Al

**Fan sediments:** poorly sorted sand and gravel, with diamicton; 2 to 15 m thick on average; form fans at the toes of slopes; composition is dependent on source materials

Au

**Alluvial sediments, undivided:** undivided floodplain, terrace, deltaic, and fan sediments

Ch

**COLLUVIAL DEPOSITS:** diamicton and rubble accumulated from various mass wasting processes, ranging from slope wash to rock fall; composition dependent on source materials

**Landslide material:** mostly unconsolidated sediments, with texture dependent on source materials; generally 1 to 10 m thick, but may exceed 10 m near the toes of large landslides; forms hummocky accumulations on lower slopes and valley floors; commonly developed in glaciolacustrine sediments

Cs

**Slope colluvium:** rock fragments in a matrix of sand, silt, and minor clay; 1 to 5 m thick; formed by reworking of unconsolidated deposits on steep (>40°) slopes; commonly gullied

Ca

**Colluvial apron and talus:** rubble and block accumulations at the bottom of steep (>40°) slopes forming aprons and cones; 1 to 10 m thick

FRASER GLACIATION (WISCONSINAN)

PROGLACIAL AND GLACIAL ENVIRONMENT

Lp

**GLACIOLACUSTRINE DEPOSITS:** well sorted, stratified sand, silt, and clay deposited in deep water of former glacial lakes; include sporadic sand and gravel deposited in a nearshore environment; sand, silt, and clay commonly occur as rhythmites with rare debris flow interbeds; outliers are common on adjacent units; contacts between subunits Lp, Lb, and Lv are gradational

**Glaciolacustrine plain:** well sorted, stratified sand, silt, and clay; generally >10 m thick; masks the underlying topography; locally incised

Lb

**Glaciolacustrine blanket:** well sorted, stratified sand, silt, and clay; 3 to 10 m thick; reflects topography of underlying units

Lv

**Glaciolacustrine veneer:** deep water deposits of well sorted, stratified sand, silt, and clay overlain, in places, by shallow water deposits of sand and gravel; occurs near limits of former glacial lakes; includes minor till outcrops; 1 to 3 m thick; reflects topography of underlying units

Gt

**GLACIOFLUVIAL DEPOSITS:** sand and gravel, well to poorly sorted, and commonly stratified; deposited by glacial meltwater; bedding disrupted locally as the result of the melting of supporting ice

**Glaciofluvial terrace sediments:** sand and gravel, stratified to massive; 1 to 10 m thick; perched above alluvial deposits or associated with meltwater channels

Gb

**Glaciofluvial blanket:** sand and gravel, stratified to massive, generally 1 to 5 m thick; sediment cover is continuous but the underlying morphology is visible

Gd

**Proglacial deltaic sediments:** sand and gravel with minor silt and clay; commonly overlies glaciolacustrine silt and clay; 5 to 10 m thick; form slightly inclined surfaces

Gh

**Ice contact deposits:** sand and gravel, stratified to massive and commonly faulted; generally greater than 3 m thick; form hummocky and kettled surfaces

GLACIAL ENVIRONMENT

Tm

**TILL:** pebbles, cobbles, and boulders in a sandy to clayey matrix; includes colluvium (reworked till) on steep slopes, and small inclusions of glaciofluvial sediments, especially in valley bottoms and near the mouth and banks of meltwater channels. The till surface is commonly fluted and drumlinized. Suffix '-c' denotes the presence of abundant meltwater channels (e.g. Tv-c)

**Thick till, rolling:** till cover; greater than 3 m thick; masks the underlying topography; bedrock outcrops are rare

Tb

**Till blanket:** continuous till cover with few bedrock outcrops; 1 to 3 m thick on average; conforms to and locally obscures topography of underlying units

c-Tb

**Pinchi Creek lens:** clayey till with a low clast content (10-15 %); 1 to 15 m thick; conforms to and locally obscures topography of underlying units; geographically restricted to the northeast of Stuart Lake, locally overlain by a glaciolacustrine veneer

Tv

**Till veneer:** discontinuous till cover with abundant bedrock outcrops; 1 m average thickness; reflects topography of underlying units, which are predominantly bedrock

PRE-QUATERNARY


R

**BEDROCK:** Sedimentary, metamorphic, volcanic, and intrusive rocks of Precambrian to Cenozoic age

**Bedrock:** outcrop; includes, in places, a thin veneer of till and colluvium which rarely exceeds 2 m

Rs

**Sleep bedrock slopes:** outcrop with predominantly loose blocks of local bedrock and few erratics; patchy cover of till and colluvium increases in abundance downslope; steeply sloping terrain (>40°); subject to rock falls



Geology by A. Plouffe, 1990-1995

Co-ordinated by A. Plouffe through the auspices of the Canada - British Columbia Agreement on Mineral Development and Nechako NATMAP Project

Digital cartography by M.M. Proulx, Geoscience Information Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by the Geoscience Information Division

Mean magnetic declination 2000, 23°28'E, decreasing 9.5' annually. Readings vary from 22°58'E in the SE corner to 23°57'E in the NW corner of the map

Elevations in metres above mean sea level

