Natural Resources Canada | Ressources naturelles Canada
Indian and Northern Affairs Canada | Affaires indiennes et du Nord Canada
GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA
NATMAP CARTNAT
Canada's National Geoscience Mapping Program
Le Programme national de cartographie géoscientifique du Canada

DESCRIPTIVE NOTES

INTRODUCTION

This map extends 1:50 000 scale coverage of Archean and Paleoproterozoic rocks southward from Hawk Hill Lake (Aspler and Bursey, 1990) to the Sealhole and Fitzpatrick lakes area (Fig.1), updating reconnaissance mapping by Eade (1973; 1974). Three major unconformity bounded sequences are exposed: the Henik Group (Researcher), the Montgomery Group (age uncertain), and the Hurwitz Group (Paleoproterozoic).

Henik Group: (unit Amf)

The Henik Group is the local name for Archean supracrustal rocks in the central part of the Ennadai-Rankin greenstone belt (Eade, 1974). North and east of Fitzpatrick Lake, the Henik group is represented by a lower greenschist facies bimodal mafic-felsic succession (unit Amf). Mafic volcanic flows are predominant (ca. 90 %), and consist of thickly bedded, fine-grained flows ($\pm$ gabbro sills). Many of these flows display exceptionally well preserved pillow structures which locally contain varioliites. Felsic flows form massive interlayers 1-10 m thick separating mafic flows several 10's of metres thick. Inter-flow siliciclastic rocks (turbiditic sandstones and mudstones) and thin (< 1 m) cotostone lenses are locally preserved. This volcanic succession is near the base of, and likely forms a tongue within, a predominantly sedimentary sub-unit of the Henik Group exposed north of the map area near Hawk Hill Lake (Aspler and Bursey, 1990) and near Otter Lake (Aspler et al., 1994b) and informally referred to as "unit A'" (Aspler and Chiarenzelli, 1998 a).

Granite, granodiorite, gneiss (units Ag; Agn)

South of Sealhole and Fitzpatrick lakes, highly strained mafic volcanic rocks define elsamembered lozenges within well-foliated biotite-hornblende granitic rocks (unit Ag). Contacts are faulted, but the volcanic rocks are inferred to be older because they are cut by granitic dykes and occur as enclaves in the granites at all scales. Granitic gneisses (unit Agn) are exposed beneath the Hurwitz Group on the western side of Sealhole Lake. The relationship of these gneisses to the granitic rocks that cut the Henik Group is unknown.

Montgomery Group (unit A-Pm)

A unit of polymictic conglomerate and arkose (unit A-Pm), up to 2000 m thick, overlies Henik Group strata east of, and extending across, Fitzpatrick Lake. Although not exposed, the lower contact of this unit is inferred to be an angular unconformity because its map trace appears to truncate folds and fabrics in the Henik Group, and because the conglomerates contain foliated clasts that are indistinguishable from subjacent basement. Near the base of the unit, massive, thickly bedded, framework-intact cobble-boulder conglomerates containing a coarse arkose matrix are predominant, and arkoses define local decimetre- to metre-scale lenses. The conglomerates contain clasts that are representative of subjacent basement sources and consist mainly of granitic fragments (up to 90 %) with variable amounts of mafic volcanic, gabbro, and felsic volcanic debris (see Figure 3 in Aspler et al., 2000a). The clasts are subangular to well rounded; maximum sizes are up to 1 metre. Significantly, many contain an internal penetrative foliation (see below). The relative abundance of conglomeratic decreases gradually up-section such that near the top of the unit, arkoses are predominant. The arkoses form decimetre- to metre-scale parallel-stratified and trough cross-stratified sandstone beds containing local pebble-cobble layers and more broadly lenticular lenses of conglomerate. Sets of 10-15 cross bedded trough cosets host angular to subrounded mudstone rip-up clasts, and these beds are interpreted to likely represent the relief necessary for the accumulation of these thick, coarse conglomerate-bearing strata, and we interpret this sequence to have been deposited on alluvial fans that drained northward from east-trending fault scarps cut into Archean basement. The lower part of the section is considered to reflect a proximal fan setting in which repeated low-viscosity flash floods produced amalgamated stacks of framework-intact, coarse sand matrix-bearing cobble-boulder conglomerate as entrenched channel, bar and diffuse sheet deposits. Local arkose lenses likely represent incompletely preserved channel- and bar-top (waning flood) wedges. An overall fining-upward trend probably reflects scarp retreat; arkoses ($\pm$ conglomerates) in the upper part of the section are considered to be medial to distal fan deposits formed by flash floods.

The relationship of this unit to the Hurwitz Group is uncertain because of faulted contacts and covered intervals. However, an unconformity is inferred because the unit is lithologically distinct from basal Hurwitz units in the area, and because it forms an isolated wedge that appears to be discordant with respect to lower Hurwitz Group strata. Hence the unit is interpreted to constitute an outlier of the Montgomery Group which, in its type area at Montgomery Lake (Fig. 1), is separated from the overlying Hurwitz Group by an angular unconformity (Aspler and Chiarenzelli, 1998 b). Whether the Montgomery Group represents a late Neoarchean molassoid sequence or a Paleoproterozoic precursor to the Hurwitz Group remains unclear.

Hurwitz Group (units PHP; PHKml; PHKmm; PHKmcc; PHKw; PHKh; PHA; PHgb; PHW)

Unconformably overlying the Henik and Montgomery groups, the Hurwitz Group is an assemblage of siliciclastic and carbonate rocks that were deposited in a continental to shallow-marine intracratonic basin that once blanketed a large part of the Hearne domain (Fig. 1). The maximum age of Hurwitz Basin is defined by ca. 2.45 Ga baddeleyite (Heaman, 1994) from the Kaminak dykes that cut basement rocks but not the Hurwitz Group (Davidson, 1970). Although previous suggestions regarding basin initiation (Noomut, Pacitek, Kinga and Ameto formations) by lithospheric stretching related to breakup of a possible Neoarchean supercontinent (Kenorland, Aspler and Chiarenzelli, 1997; Sleep and Hansen, 1983; Bekker, new data (Aspler et al., 2000b; Davis et al., 2000) indicate that uppermost units (Wattson, Tavani and Whitehorse members) are correlatives of a regionally extensive set of gabbro sills and the Hurwitz Group ([211 $\pm$ 1 Ma baddeleyite; Heaman and Cadman, 1994).

A thick lower Hurwitz Group section in the eastern and northern parts of the map area is partly a consequence of extensive Padlei Formation conglomerate, arkose and sandstone-siltstone rhythmite deposits (unit PHP). In addition, restricted subunits appear at both the base and top of the Maguse Member. The lower subunit (unit PHKml) is ca. 1200 m thick and contains arkose and polymictic pebble conglomerate sheets as well as the subarkose to quartz arenite and quartz pebble conglomerate beds that are more typical of the Maguse Member (unit PHKmm). Moving of relatively immature and mature arenites in this subunit likely reflects variation in degree of reworking and residence time at sites of temporary storage on the Maguse fluvial plain. Near the base of this subunit is a 50 m thick interval in which decimetre- to metre-scale sheets of subarkose separate beds of dolobutite ($\pm$ microbial laminite) containing centimetre-scale sandstone lenses. With the exception of similar beds near Rankin Inlet (Ryan et al., 1999), and calc-silicates west of Edehon Lake (Eade, 1973), carbonate rocks have not been reported from elsewhere in the Kinga Formation. The carbonate-bearing beds are inferred to represent the remnants of a mixed siliciclastic-carbonate sequence that was deposited during a short-lived marine incursion of the Maguse fluvial plain. Near the top of the Maguse Member (unit PHKmm) is a 1-2 m thick unit of pebbly arkose and conglomerate that is transitional into the Padlei map area (Aspler and Chiarenzelli, 1997), that consists of polymictic pebbly arkose and conglomerate containing primarily granitic and mafic volcanic clasts.The Padlei Formation and Maguse Member thin abruptly to the south and west, such that in southern Fitzpatrick Lake and southern and western Sealhole Lake, the wave ripple-marked supermature quartz arenites of the Whiterock Member (unit PHKw) directly onlap basement (see Figure 4 in Aspler et al., 2000a). Small isolated outcrops of the Hawk Hill Member (unit PHKh), consisting of layered hematitic chert and chert breccia occur above the Whiterock Member west of White Cliff Lake, southwest of Sealhole Lake and northeast of Sealhole Lake. Exposures of Ameto Formation (unit PHA) mudcrops are limited to local host heaves. Gabbro sills extend as discontinuous pods within the Hurwitz Group for a strike length of ca. 450 km, from the Rankin Inlet area southwest to Watterson Lake (Fig.1), Typically within the Ameto Formation, the sills also occur in the lower part of the Maguse Member east of Fitzpatrick Lake, and directly above the Whiterock Member at Sealhole Lake (unit PHgb). In the latter area, the sills are commonly associated with abundant sandstone dykes (see Figure 7 in Aspler et al., 2000a). Elsewhere in the Hurwitz gabbros (see Figure 7 in Aspler et al., 2000a). On the eastern shore of Sealhole Lake, such megacrysts are distributed in layers consistent to bedding in subjacent strata; some of these layers define graded beds. Feeder dykes to the gabbro sills have not hitherto been reported, but megacryst-bearing, north and northeast-trending dykes that are indistinguishable from the gabbro sills outcrop east of Fitzpatrick Lake, and we suggest that these are are examples of feeder dykes. A mixed siliciclastic-carbonate facies of the Watterson Formation (unit PHW) outcrops on a north-projecting peninsula in southern Sealhole Lake, but contacts with underlying units areaxce not exposed.

STRUCTURAL GEOLOGY

Evidence for Archean deformation, and at least one episode of Paleoproterozoic (post-Hurwitz Group) deformation, is preserved in the Sealhole-Fitzpatrick Lake area.

Archean deformation: Henik Group and allied granitic rocks

Three lines of evidence suggest pre-Paleoproterozoic deformation of Archean rocks. First, Montgomery Group and lower Hurwitz Group conglomerate units contain abundant well foliated basement-derived granitic and volcanic clasts. Second, east of Fitzpatrick Lake, east-northeast- trending folds and fabrics in the Henik Group are deflected and truncated by northwest-trending structures in the Hurwitz Group. Third, adjacent to the large granitic body south of Sealhole and Fitzpatrick lakes, granitic dykes cut mafic volcanic rocks. Some granitic dykes are folded with, and contain the same northeast-trending penetrative fabric as, the volcanic rocks, whereas other such dykes cross-cut the penetrative fabric. Hence, deformation and granite emplacement appear to have overlapped in time. Although the granitic rocks are undated, they are unconformably beneath the Hurwitz Group and are interpreted to be Archean, and thus deformation was probably Archean. Northeast-trending fabrics in Archean rocks south of Sealhole and Fitzpatrick lakes are subparallel to some structures in the Hurwitz Group, indicating that Archean and Paleoproterozoic deformation were, in part, coaxial.

Paleoproterozoic (?) deformation: Montgomery Group

Rocks of the Montgomery Group are deformed by northeast-trending folds and related fabrics that are subparallel to structures in the Hurwitz Group. We cannot specify with certainty that these structures are post-Hurwitz Group because they may represent a pre-Hurwitz Group event such as recorded in the Montgomery Group type area. However, the Hurwitz gabbro dykes that cut the Montgomery Group contain cleavage parallel to cleavage in host strata, and hence we infer that the deformation recorded in the Montgomery Group is Paleoproterozic.

Paleoproterozoic deformation: Hurwitz Group

At a simplified level, post-Hurwitz Group structures in the Sealhole-Fitzpatrick lake area define a northerly trending basement-cover synclinorium. At a detailed level however, the structure is more complex, and variably oriented folds and faults reflect heterogeneous strain from shortening of a lithologically anisotropic assemblage of non-uniform thickness.

The structural grain of the area is controlled by northeast- and north-trending folds, high-angle faults and cleavage. Folds are open, concentric and locally display a box-like morphology. Axial surfaces are steep, and hinge lines plunge gently to the north and south. Plunge reversals about northwest trends produce dome and basin (Type-1) interference structures, but a northwest-trending cleavage has not been observed, and the near-vertical, northeast-trending cleavage surfaces are undeflected about northwest trends. This interference style, best exemplified by a northwest-elongate canoe-shaped synclinr northwest of White Cliff Lake, probably represents culmination and depression growth during progressive folding rather than discrete "D" and "D" events. Near the core of the synclinorium, the Hurwitz Group is cut by a number of northeast-trending, high-angle faults that have both reverse and normal offsets. Approximately 10 km north of Fitzpatrick Lake, one fault branches into three splays that rejoin along strike farther south in Sealhole Lake. Together the faults define a duplex-like structure in which individual splays separate repeated lower Hurwitz Group panels. This duplex-like structure probably formed to accommodate constrictions in the core of the main basement-cover synclinorium. Similarly, variably oriented high-angle oblique-slip cross faults, particularly common where the sub-Whiterock Member section is thin or non-existent (e.g. southwestern Sealhole Lake), typically fan around fold hinge zones and likely were formed to accommodate fold tightening during concentric folding.

REFERENCES

Aspler, L.B., and Bursey, T.L.
1990: Stratigraphy, sedimentation, dome and basin basement-cover infolding and implications for gold in the Hurwitz Group, Hawk Hill-Griffin Mountain Lakes area, District of Keewatin; in Current Research Part C, Geological Survey of Canada Paper 90-1C, p. 219 -230.

Aspler, L.B., and Chiarenzelli, J.R.
1996a: Stratigraphy, sedimentology and physical volcanology of the Ennadai-Rankin greenstone belt, Northwest Territories, Canada: Late Archean paleogeography of the Hearne Province and tectonic implications; Precambrian Research, v. 77, p. 59 -99.
1996b: Relationship between the Montgomery Lake and Hurwitz groups, and stratigraphic revision of the lower Hurwitz Group, District of Keewatin; Canadian Journal of Earth Sciences, v. 33, p. 1243 -1255.
1997: Initiation of ca. 2.45-2.1 Ga intracratonic basin sedimentation of the Hurwitz Group, Keewatin Hinterland, Northwest Territories, Canada. Precambrian Research, v. 81, p. 269- 297.
1998: Two Neoarchean supercontinents ?: evidence from the Paleoproterozoic; Sedimentary Geology, v. 120, p. 75- 104.

Aspler, L.B., Chiarenzelli, J. R., and Bursey, T.L.
1994a: Ripple marks in quartz arenites of the Hurwitz Group, Northwest Territories, Canada: Evidence for sedimentation in a vast, Early Proterozoic, shallow, fresh-water lake; Journal of Sedimentary Research, v. 64 A, p. 282 -298.

Aspler, L.B., Chiarenzelli, J. R., Ozarko, D.L., and Powis, K.B.
1994b: Geology of Archean and Proterozoic supracrustal rocks in the Otter and Ducker lakes area, southern District of Keewatin, Northwest Territories; in Current Research 1994-C; Geological Survey of Canada, p.165- 174.

Aspler, L.B., Höfer, C., and Harvey, B.J.A.
2000a: Geology of the Henik, Montgomery and Hurwitz groups, Sealhole and Fitzpatrick lakes area, Nunavut: in Current Research 2000-C12, Geological Survey of Canada, 10p.

Aspler, L.B., Chiarenzelli, J. R, Cousens, B.L., Davis, W., McNicoll, V.J., and Rainbird, R.H.
2000b: Intracratonic basin processes from breakup of Kenorland to assembly of Laurentia: new geochronology and models for Hurwitz Basin, western Churchill Province. GeoCanada 2000-The Millennium Geoscience Summit Conference CD (extended abstract).

Davis, W.J., Aspler, L.B., Rainbird, R.H., and Chiarenzelli, J.R.
2000: Detrital zircon geochronology of the Proterozoic Hurwitz and Kluyck groups: a revised post-1.92 Ga age for deposition of the upper Hurwitz Group. GeoCanada 2000-The Millennium Geoscience Summit Conference CD (extended abstract).

Davidson, A.
1970: Precambrian geology, Kaminak Lake map-area, District of Keewatin; Geological Survey of Canada Paper 69-51, 27 p.

Eade, K.E.
1973: Geology of Nueltin Lake and Edehon Lake (west half) map-areas, District of Keewatin. Geological Survey of Canada Paper 72-21, 29 p.

Eade, K.E.
1974: Geology of Kognak River area, District of Keewatin, Northwest Territories; Geological Survey of Canada Memoir 377, 66p.

Heaman, L.M.
1994: 2.45 Ga global mafic magmatism: Earth's oldest superplume ? in Abstracts of the Eighth International Conference on Geochronology, Cosmochronology and Isotope Geology, M.A. Lauphere, G.B. Dalrymple, B.D. Turrin (ed.); United States Geological Survey Circular 1107, p.132.

Heaman, L.M., and LeCheminant, A.N.
1993: Paragenesis and U-Pb systematics of baddeleyite (ZrO$_2$); Chemical Geology, v. 110, p.95 -126.

Ryan, J.J., Aspler, L.B., Tella, S., Sandeman, H., and Studnicki-Gizbert, C.
1999: Revision of Archean and Paleoproterozoic stratigraphy at Rankin Inlet: implications for timing of multiple regional deformations, Kivalliq Region, Nunavut; 27th Annual NWT Geoscience Forum; Program and Abstracts. Indian and Northern Affairs Canada, Yellowknife, p. 60- 61.



LEGEND

PALEOPROTEROZOIC
HURWITZ GROUP

- **PHW** — *WATTERSON FORMATION: Mixed siliciclastic-carbonate facies comprising mm- and cm-scale massive dolobutite and stratified calcilutite that is rhythmically interlayered with siliciclastic siltstone and mudstone, and occurs at the tops of very fine, parallel-stratified (± ripple drift cross-stratified) sandstone-siltstone-mudstone fining-upward sequences. Local gutter casts at base of fining-upward sequences filled with intraclastic calcirudite breccia. Abundant soft sediment deformation: load casts and flame structures; calcilutite dykes with wall-parallel internal layering*

unconformity

- **PHgb** — *Discontinuous coarse-grained gabbro sills, dykes. Feldspar megacrysts, commonly in layers concordant with bedding in adjacent units. 2111±0.6 Ma (Heaman and LeCheminant 1993). Post lithification of Ameto Formation; pre-deposition of Watterson Formation. Folded with Hurwitz Group*
- **PHA** — *AMETO FORMATION: Frost-heaved mudrocks in map area, but immediately north at Hawk Hill Lake, varicoloured siltstone, mudstone, calcisiltite in fining-upward sequences*

KINGA FORMATION

- **PHKh** — *Hawk Hill Member: Discontinuous unit of bedded white chert, maroon chert and chert breccia*

Whiterock Member

- **PHKw** — *Supermature quartz arenite. Lower part of section commonly massive, upper 200 m with ubiquitous wave ripples. Paleodepth calculations based on symmetric, vortex orbital ripples indicates depths of 2 cm to 2 m for the entire sequence (Aspler et al., 1994 a)*

Maguse Member

- **PHKmu** — *Upper Maguse Member: Massive polymictic pebbly sandstone with angular to subrounded basement granules and pebbles (10-50%) dispersed in arkose matrix. Local parallel-stratified arkose and framework-intact pebble-cobble conglomerate with well-rounded clasts and arkose matrix*
- **PHKmm** — *Middle Maguse Member: Maroon, pink, grey subarkose to quartz arenite; interbeds of white quartz arenite at top. Local black parallel and cross-stratified heavy mineral bands. Quartz pebble conglomerate with clasts of well-rounded spherical white quartz, jasper, blue and grey chert*
- **PHKml** — *Lower Maguse Member: Parallel-stratified m-scale sheets of arkoses, subarkose, and quartz arenite. Polymictic framework-intact and framework-disrupted pebble-cobble conglomerate. Local graded channels. Quartz-pebble conglomerate, 50 m interval near base: dcm- to m-scale beds of subarkose alternating with carbonate-rich beds comprising cm-scale layers of massive and microbial laminated dolobutite (± chert laminae) and cm-scale siliciclastic sand lenses*
- **PHP** — *PADLEI FORMATION: Stratified polymictic conglomerate; arkose; mm-cm-scale sandstone-siltstone rhythmites*

unconformity

PALEOPROTEROZOIC OR ARCHEAN
MONTGOMERY GROUP

- **A-PM** — *At base: massive, thickly bedded, framework-intact cobble-boulder conglomerate with local arkose lenses. Clasts (to 1 m) representative of subjacent basement; many with pre-depositional tectonic fabric. At top : mainly parallel- and cross-stratified arkose with local conglomerate*

unconformity

ARCHEAN

- **Ag; Agn** — *Well-foliated biotite-hornblende granite, granodiorite* Ag; *locally gneissic* Agn

intrusive contact

HENIK GROUP

- **Amf** — *Bimodal mafic-felsic volcanic rocks. Predominantly fine-grained flows, commonly pillowed, varioltic. Interlayered with felsic flows 1-10 m thick. Local inter-flow sedimentary rocks*

Geological boundary (approximate)
Fault (defined, approximate)
Reverse fault (defined)
Normal fault (defined)
Axial trace, arrow indicates plunge (anticline, syncline)
Outcrop
Frost heave; probable subjacent outcrop
Bedding (inclined, overturned, tops unknown)
Bedding from pillow lavas
Layering in Hurwitz gabbro sills
Foliation (Hurwitz Group) (inclined)
Foliation (Montgomery Group) (inclined)
Foliation (Henik Group and related plutonic rocks; (inclined, vertical)
Axial projection line (A — A')



Figure 1. Simplified geology of Hearne domain and location of map area

LOCATION MAP

Geology by L.B. Aspler, C. Höfer and B.J.A. Harvey, 1999

Geological compilation by L.B. Aspler, 2000

Co-ordinated through the auspices of the Western Churchill NATMAP Project

Digital cartography by R.L. Allard, Earth Sciences Sector Information Division (ESS Info)

EGS 2000-14

OPEN FILE 3753

GEOLOGY

SEALHOLE AND FITZPATRICK LAKES AREA

NUNAVUT

Scale 1:50 000/Échelle 1/50 000

kilometres 1 0 1 2 3 4 kilometres

Universal Transverse Mercator Projection
North American Datum 1927
© Her Majesty the Queen in Right of Canada, 2000

Projection transverse universelle de Mercator
Système de référence géodésique nord-américain, 1927
© Sa Majesté la Reine du chef du Canada, 2000

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map supplied by the Author modified from data compiled by Geomatics Canada

Magnetic declination 2000, 6° 24' E, decreasing 11.0' annually

Elevations in metres above mean sea level



NATIONAL TOPOGRAPHIC SYSTEM REFERENCE

OPEN FILE
DOSSIER PUBLIC
3753
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
OTTAWA
12/2000

EGS 2000-14

Recommended citation:
Aspler, L.B., Höfer, C., and Harvey, B.J.A.
2000: Geology, Seahole and Fitzpatrick Lakes area, Nunavut; Geological Survey of Canada, Open File 3753, scale 1:50 000.

Canada