



GEOLOGICAL SURVEY OF CANADA  
BULLETIN 564

# PETROLEUM RESOURCE POTENTIAL OF SEDIMENTARY BASINS ON THE PACIFIC MARGIN OF CANADA

P.K. Hannigan, J.R. Dietrich,  
P.J. Lee, and K.G. Osadetz



2001



Natural Resources Canada  
Ressources naturelles Canada

Canada

GEOLOGICAL SURVEY OF CANADA  
BULLETIN 564

**PETROLEUM RESOURCE POTENTIAL OF  
SEDIMENTARY BASINS ON THE  
PACIFIC MARGIN OF CANADA**

P.K. Hannigan, J.R. Dietrich,  
P.J. Lee, and K.G. Osadetz

**2001**

©Her Majesty the Queen in Right of Canada, 2001  
Catalogue No. M42-564E  
ISBN 0-660-18288-2

Available in Canada from  
Geological Survey of Canada offices:

601 Booth Street  
Ottawa, Ontario K1A 0E8

3303-33rd Street N.W.  
Calgary, Alberta T2L 2A7

101-605 Robson Street  
Vancouver, B.C. V6B 5J3

A deposit copy of this publication is available for  
reference in public libraries across Canada

Cette publication est aussi disponible en français

Price subject to change without notice

**Cover Illustration**

Tian Bay, on the west coast of Graham Island, Queen Charlotte Islands, British Columbia, site of the first well drilled in the area (1913). Indications of dispersed bitumen, oil, gas, and tar found in Masset Formation basalts were the main incentive for drilling the Tian Bay well. Natural gas flows were encountered in the subsurface. GSC C 4609-1

**Critical readers**

*L. Currie*  
*K. Rohr*

**Authors' address**

*P.K. Hannigan, J.R. Dietrich, K.G. Osadetz*  
*Geological Survey of Canada*  
*3303-33rd Street N.W.*  
*Calgary, AB T2L 2A7*

*P.J. Lee (deceased)*

*Manuscript submitted: 1998-05*  
*Approved for publication: 2000-12*

## CONTENTS

1	Abstract
1	Résumé
1	Summary
3	Sommaire
5	Introduction
5	Scope
5	Purpose
5	Terminology
9	Method and content
9	Previous assessments
9	Tectonic setting
11	Resource assessment procedure
11	Geological play definition
11	Compilation of play data
13	Conceptual play analysis
13	Queen Charlotte assessment region
13	Exploration history and regional studies
15	Regional geology
15	Geological setting and tectonic evolution
16	Stratigraphy and structure
17	Petroleum geology
17	Reservoirs
17	Mesozoic
19	Paleogene
19	Neogene
20	Seals
20	Traps
20	Source rocks
21	Source rock maturation
21	Timing of hydrocarbon generation
22	Hydrocarbon shows
24	Petroleum assessment
24	Petroleum plays
24	Queen Charlotte Cretaceous oil and gas play
24	Queen Charlotte Miocene oil and gas play
27	Queen Charlotte Pliocene oil and gas play
29	Discussion of assessment results
32	Georgia assessment region
32	Exploration history and regional studies
34	Regional geology
34	Geological setting and tectonic evolution
34	Stratigraphy and structure
35	Petroleum geology
35	Reservoirs
35	Mesozoic
36	Tertiary
37	Pleistocene
37	Seals
37	Traps
37	Source rocks
38	Source rock maturation
38	Timing of hydrocarbon generation
40	Hydrocarbon shows
41	Petroleum assessment

41	Petroleum plays
41	Georgia Cretaceous structural gas play
41	Georgia Tertiary structural gas play
41	Georgia Pleistocene stratigraphic gas play
45	Discussion of assessment results
45	Tofino assessment region
45	Exploration history and regional studies
47	Regional geology
47	Geological setting and tectonic evolution
47	Stratigraphy and structure
49	Petroleum geology
49	Reservoirs
49	Tertiary
49	Pleistocene
50	Seals
50	Traps
50	Source rocks
50	Source rock maturation
51	Timing of hydrocarbon generation
51	Hydrocarbon shows
51	Petroleum assessment
51	Petroleum play
51	Tofino Tertiary structural gas play
51	Discussion of assessment results
53	Basin comparisons
54	Conclusions
55	References
63	Appendix A

## Figures

6	1. Geographic setting of the west coast region of Canada
7	2. Regional setting and basin outlines of the west coast region of Canada
8	3. Detail map of Georgia Basin in Vancouver area, the Fraser River lowlands and delta, southeastern Vancouver Island and Gulf Islands of British Columbia and northwestern Washington
12	4. Tertiary and Cretaceous onshore sedimentary basins, west coast region of Canada
14	5. Seismic reflection profile, Dixon Entrance, northern Queen Charlotte Basin
14	6. Seismic reflection profile, Hecate Strait, central Queen Charlotte Basin
15	7. Seismic reflection profile, Queen Charlotte Sound, southern Queen Charlotte Basin
16	8. Simplified stratigraphic column for Queen Charlotte, Georgia, Tofino and Juan de Fuca sub-basin regions
18	9. Geological cross-sections across Hecate Strait and Queen Charlotte Sound
21	10. Maturation–depth profile of strata in the Sockeye B-10 location, Queen Charlotte Basin
22	11. Maturation–depth profile of strata in the Sockeye E-66 location, Queen Charlotte Basin
23	12. Subsidence and hydrocarbon generation model for the area of Sockeye B-10 well, central Hecate Strait, Queen Charlotte Basin
23	13. Subsidence and hydrocarbon generation model for the area of Sockeye E-66 well, central Hecate Strait, Queen Charlotte Basin
23	14. Heat-flow model for central Hecate Strait, Queen Charlotte Basin
25	15. Queen Charlotte Cretaceous oil and gas play area
26	16. Estimate of in-place oil potential of the Cretaceous play in Queen Charlotte Basin
26	17. Estimate of in-place gas potential of the Cretaceous play in Queen Charlotte Basin
28	18. Queen Charlotte Miocene oil and gas play area
29	19. Estimate of in-place oil potential of the Miocene play in Queen Charlotte Basin
29	20. Estimate of in-place gas potential of the Miocene play in Queen Charlotte Basin
30	21. Queen Charlotte Pliocene oil and gas play area
31	22. Estimate of in-place oil potential of the Pliocene play in Queen Charlotte Basin

- 31 23. Estimate of in-place gas potential of the Pliocene play in Queen Charlotte Basin
- 31 24. Estimate of total oil potential for the Queen Charlotte Basin region
- 31 25. Estimate of total gas potential for the Queen Charlotte Basin region
- 33 26. Geological cross-sections across the Strait of Georgia, and running parallel to the mainland coast from Burrard Inlet to Bellingham
- 39 27. Maturation–depth profile of strata in the Yellow Point d-84-C location, Georgia Basin
- 39 28. Maturation–depth profile of strata in the Richfield Pure Point Roberts 6-3-5 location, Georgia Basin
- 40 29. Subsidence and hydrocarbon generation model for the Point Roberts area, Tertiary Georgia Basin
- 40 30. Subsidence and hydrocarbon generation model for Yellow Point area, Cretaceous Georgia Basin
- 42 31. Georgia Cretaceous structural gas play area
- 43 32. Estimate of in-place gas potential of the Cretaceous structural play in Georgia Basin
- 44 33. Georgia Tertiary structural gas play area
- 45 34. Estimate of in-place gas potential of the Tertiary structural play in Georgia Basin
- 46 35. Georgia Pleistocene stratigraphic gas play area
- 47 36. Estimate of in-place gas potential of the Pleistocene stratigraphic play in Georgia Basin
- 47 37. Estimate of total gas potential for the Georgia Basin region
- 48 38. Geological cross-section in Tofino Basin
- 48 39. Seismic reflection profile, Pacific Ocean offshore, southern Vancouver Island
- 49 40. Seismic reflection profile, Pacific Ocean offshore, northern Vancouver Island
- 50 41. Maturation–depth profile of strata in the Shell Anglo Pluto I-87 well location, Tofino Basin
- 51 42. Subsidence and hydrocarbon generation model for area of Pluto I-87 well, Tofino Basin
- 52 43. Tofino Basin Tertiary structural gas play area
- 53 44. Estimate of in-place gas potential of the Tertiary structural play in Tofino Basin region
- 55 45. Estimate of total gas potential for west coast basins

**Tables**

- 9 1. Well reference numbers in Queen Charlotte, Tofino, and Juan de Fuca basins
- 10 2. Well reference numbers in Georgia Basin
- 26 3. Petroleum trap types in west coast basins
- 27 4. Oil and gas potential in Queen Charlotte, Georgia, and Tofino regions



# PETROLEUM RESOURCE POTENTIAL OF SEDIMENTARY BASINS ON THE PACIFIC MARGIN OF CANADA

## *Abstract*

There are three major sedimentary basins on the Pacific margin of Canada: the Tofino Basin, which overlies the present convergent plate margin west of Vancouver Island; the Queen Charlotte Basin, a transtensional basin lying adjacent to the transform boundary between the North American and Pacific plates; and the Georgia Basin, a sedimentary basin with foreland affinities in Late Cretaceous to Eocene time, which in the southeast is overlain by the Fraser River delta. The location of these sedimentary basins at complex plate margins has resulted in a variety of depositional settings that have affected source rock potential and reservoir development. The plate interactions have resulted in a highly variable and complicated tectonic history, which has influenced trap formation, maturation history and petroleum migration. Limited gas production from Pleistocene sediments in the Fraser delta, a good gas show in a well on the Olympic Peninsula and numerous seepages and other indications of hydrocarbons at surface, in wells and on seismic sections throughout the area, all point to an unrealized hydrocarbon potential. The probabilistic assessment of oil and gas potential for the sedimentary basins on the Pacific margin indicates that the total median estimate of in-place hydrocarbon volumes for all west coast basins is in the order of  $1560 \times 10^6 \text{ m}^3$  ( $9.8 \times 10^9$  bbl) of oil and  $1228 \times 10^9 \text{ m}^3$  (43.4 TCF) of gas. There are no discovered reserves in these west coast plays, but some 97 gas pools larger than  $3 \times 10^9 \text{ m}^3$ , and two oil pools larger than  $160 \times 10^6 \text{ m}^3$ , are expected to be present. Several of the conceptual plays have significant potential with respect to undiscovered gas volume, particularly the Pliocene and Miocene gas plays in Queen Charlotte Basin and the Tertiary structural play in the Tofino region. Estimates for oil potential are less optimistic.

## *Résumé*

Au Canada, la marge continentale du Pacifique renferme trois importants bassins sédimentaires : le bassin de Tofino, qui s'étend sur l'actuelle marge continentale active à l'ouest de l'île de Vancouver; le bassin de Reine-Charlotte, un bassin de transtension attenant à la marge transformante entre la plaque nord-américaine et la plaque pacifique; et le bassin sédimentaire de Georgia, étroitement apparenté à un avant-pays au Crétacé tardif-Éocène, qui comprend au sud-est le delta du fleuve Fraser. L'emplacement de ces bassins sédimentaires aux marges de plaques complexes a donné naissance à divers milieux de sédimentation qui ont eu une incidence sur le potentiel des roches mères et sur la formation des réservoirs. Les interactions des plaques sont à l'origine d'une histoire tectonique extrêmement complexe et variable qui a influé sur la formation des pièges et sur l'histoire de maturation et la migration du pétrole. La production limitée de gaz naturel dans les sédiments pléistocènes du delta du fleuve Fraser, un bon indice de gaz dans un puits situé sur la presqu'île Olympic et de nombreux suintements et autres indices en surface d'hydrocarbures dans des puits et des profils sismiques à l'échelle de la région, sont autant d'indicateurs d'un potentiel latent en hydrocarbures. L'évaluation probabiliste du potentiel en pétrole et en gaz des bassins sédimentaires de la marge du Pacifique montre que l'estimation médiane globale des volumes d'hydrocarbures en place est de l'ordre de  $1560 \times 10^6 \text{ m}^3$  ( $9,8 \times 10^9$  barils) de pétrole et de  $1228 \times 10^9 \text{ m}^3$  ( $43,4 \times 10^{12} \text{ pi}^3$ ) de gaz. Ces zones pétrolifères et gazéifères de la côte ouest ne recèlent aucune réserve découverte. Cependant, on estime qu'elles comptent environ 97 gisements de gaz naturel de plus de  $3 \times 10^9 \text{ m}^3$  et deux gisements de pétrole de plus de  $160 \times 10^6 \text{ m}^3$ . Un certain nombre de ces zones possibles montrent un important potentiel en regard du volume de gaz non encore découvert, en particulier, les zones gazéifères du Pliocène et du Miocène présentes dans le bassin de Reine-Charlotte et la zone d'origine structurale du Tertiaire dans la région de Tofino. Les estimations du potentiel en pétrole sont moins optimistes.

## *Summary*

The oil and gas resource potential in Cenozoic and Mesozoic strata of the sedimentary basins on the Pacific margin of Canada are described in this report. The overall appraisal of the hydrocarbon potential of these west coast basins constitutes one of a series of reports from an ongoing comprehensive update of the total petroleum resource for all the sedimentary basins of Canada. The previous Canada-wide assessment was



described in a 1983 Geological Survey of Canada paper (Procter et al., 1983). These subsequent “updated” reports contain major revisions and new estimates of petroleum potential for most parts of Canada. Several additional reports on other sedimentary basins, similar to this volume, are planned.

A petroleum play is defined as a group of prospects forming a common geological population linked by one or more factors such as stratigraphy, structure, reservoir type, or source-rock type. For the west coast basins of Canada (the Queen Charlotte, Georgia, Tofino and Winona basins), ten conceptual plays were defined on the basis of various geological controls, with most plays identified from stratigraphic considerations (e.g., Tertiary and Cretaceous plays). Once defined, the ten plays were each statistically analyzed to estimate their petroleum resource potential. Resource numbers (total potential and field sizes) quoted in this report are all median value estimates of in-place hydrocarbon volumes.

The oil and natural gas potential of conceptual plays is calculated using a subjective assessment technique termed *conceptual play analysis*. Conceptual plays are defined as those plays that do not yet have discoveries or established reserves, but which may exist according to geological analyses. Conceptual play analysis assumes that the individual sizes of “pools” in a properly defined play form a natural geological population and that the distribution of pool sizes within that population is lognormal. Judging by previous studies using the *discovery process model* for mature play analysis, lognormal distributions adequately represent geological populations in most cases. Also, if lognormal distributions of individual reservoir parameters are entered into the standard “pool-size” equation, a lognormal distribution of pool or prospect size is derived. The distribution of pool sizes is then combined with an additional distribution describing the number of prospects and the marginal probabilities of risk factors to calculate an estimate of both play potential and individual undiscovered pool sizes. In conceptual plays, where detailed engineering studies of petroleum reserves are missing, it is proper to identify these accumulations as fields rather than pools. Fields in this context are characterized as one or more oil or gas pools in a single structure or trap.

The estimated undiscovered potential for all plays in the west coast basins of Canada is  $1560 \times 10^6 \text{m}^3$  ( $9.8 \times 10^9$  bbl) of in-place oil and  $1228 \times 10^9 \text{m}^3$  (43.4 TCF) of in-place gas. No reserves have been established in the area; 100 per cent of the resource quoted remains to be discovered.

Results of the conceptual play analysis indicate that three plays have high potential for containing significant amounts of gas. These are 1) the Pliocene gas play in the Queen Charlotte Basin region, 2) the Queen Charlotte Miocene gas play, and 3) the Tofino structural gas play.

Total estimated in-place oil and gas potential for the six Queen Charlotte conceptual plays (i.e., oil and gas components in three defined plays) is  $1560 \times 10^6 \text{m}^3$  ( $9.8 \times 10^9$  bbl) and  $734 \times 10^9 \text{m}^3$  (25.9 TCF), respectively. Estimated oil resource in three of the Queen Charlotte plays is slightly less abundant than gas in terms of energy-equivalent volumes. The combined presence of abundant reservoir strata, good petroleum source rocks, numerous and diverse structural traps, and the common occurrence of oil and gas shows reflect the significant potential for petroleum accumulations. The most prospective areas occur in Neogene strata within the Queen Charlotte Basin, beneath eastern Graham Island and in the offshore shelf areas of Dixon Entrance, Hecate Strait and Queen Charlotte Sound. Forty-one undiscovered gas fields with in-place gas volumes of  $3 \times 10^9 \text{m}^3$  (~ 100 BCF) or larger are predicted to be present in these three gas plays. Also, two fields greater than  $160 \times 10^6 \text{m}^3$  ( $10^9$  bbl) of original in-place oil are anticipated for the three oil plays. Considerable potential is recognized in at least two gas plays in the Queen Charlotte area, making renewed exploration attractive. The current assessment indicates a substantially greater petroleum resource potential for the Queen Charlotte Basin region than previously thought (Procter et al., 1983).

The total estimated potential for three Georgia Basin conceptual plays is  $185 \times 10^9 \text{m}^3$  (6.5 TCF) of in-place gas. Available geochemical information indicates there is little oil potential in the Georgia Basin or Tofino regions and as such no oil play assessments were prepared for these areas. Eighteen undiscovered gas fields with in-place gas volumes of greater than  $3 \times 10^9 \text{m}^3$  are predicted to occur in two gas plays. Significant potential is predicted for the Georgia Cretaceous structural play even with substantial risk assigned to adequacy of reservoir and source rock. The presence of large closed structures and large play area increases play potential.

The Tofino Basin region has a single defined play. The potential for the play is  $266 \times 10^9 \text{ m}^3$  (9.4 TCF) of in-place gas. Thirty-eight fields with more than  $3 \times 10^9 \text{ m}^3$  gas volume are predicted to occur, indicating considerable gas potential in this single play. Substantial risk was assigned to reservoir facies and source rock in this play, but the abundance of large structures across a vast area provide the framework for a moderately favourable resource estimate.

## *Sommaire*

Le présent document décrit les ressources pétrolières et gazières potentielles que renferment les strates cénozoïques et mésozoïques des bassins sédimentaires de la marge pacifique du Canada. L'évaluation globale du potentiel en hydrocarbures des bassins de la côte ouest s'insère dans une série de rapports dont le but est de présenter une mise à jour exhaustive en cours des ressources globales en pétrole de tous les bassins sédimentaires du Canada. La dernière évaluation faite à l'échelle du Canada a été publiée par la Commission géologique du Canada (CGC) en 1983 (Procter et al., 1983). Les prochains rapports de «mise à jour» feront état d'importantes révisions et présenteront de nouvelles estimations sur le potentiel en pétrole de la plupart des régions du Canada. Plusieurs rapports semblables au présent document, mais portant sur d'autres bassins sédimentaires, sont prévus.

Une zone pétrolifère comporte un groupe de sites d'intérêt formant une population géologique qui partage une ou plusieurs caractéristiques, telles que la stratigraphie, la structure, le type de réservoir ou le type de roche mère. Dix zones possibles ont été définies dans les bassins de la côte ouest du Canada (bassins de Reine-Charlotte, de Georgia, de Tofino et de Winona) en fonction de divers facteurs géologiques; la plupart des zones ont été identifiées au moyen de critères stratigraphiques (p. ex. les zones du Tertiaire et du Crétacé). Puis, les dix zones ont fait l'objet d'une analyse statistique destinée à l'évaluation des ressources pétrolières potentielles. Les chiffres sur les ressources (potentiel global et dimension des champs) fournis dans le présent rapport correspondent tous à des estimations des valeurs médianes des volumes d'hydrocarbures en place.

Le potentiel en pétrole et en gaz naturel des zones possibles est calculé à l'aide d'une méthode d'évaluation subjective appelée *analyse des zones possibles*. Les zones possibles sont des zones dans lesquelles aucune découverte n'a été effectuée, ou aucune réserve n'a été établie, mais pour lesquelles l'analyse géologique en indique l'existence possible. L'analyse des zones possibles part du principe que les volumes de chaque «gisement» dans une zone adéquatement définie forment une population géologique naturelle et que la distribution de ces volumes est lognormale. Si on en juge par des études antérieures qui ont appliqué le *modèle du processus de découverte* à l'analyse de zones bien explorées, dans la plupart des cas, les distributions lognormales représentent adéquatement les populations géologiques. De même, si les distributions lognormales des paramètres de chaque réservoir entrent dans l'équation standard volume du «gisement», on peut en dériver une distribution lognormale des volumes des gisements ou des sites d'intérêt. La distribution des volumes des gisements est alors combinée à une autre distribution décrivant le nombre de sites d'intérêt et les probabilités marginales des facteurs de risque afin d'évaluer le potentiel des zones et les volumes de chaque gisement non encore découvert. Dans les zones possibles, où on n'a pas effectué d'études techniques détaillées des réserves pétrolières, il est indiqué de définir ces accumulations comme des champs et non comme des gisements. Dans ce contexte, les champs sont définis comme correspondant à un ou plusieurs gisements de pétrole ou de gaz contenus dans un piège ou une structure unique.

Dans les bassins de la côte ouest du Canada, le potentiel non découvert de l'ensemble des zones pétrolifères et gazéifères est estimé à  $1\,560 \times 10^6 \text{ m}^3$  ( $9,8 \times 10^9$  barils) de pétrole en place et à  $1\,228 \times 10^9 \text{ m}^3$  ( $43,4 \times 10^{12} \text{ pi}^3$ ) de gaz en place. Aucune réserve n'a été prouvée dans la région; les ressources dont il est fait mention restent à être découvertes dans leur intégralité.

Les résultats de l'analyse des zones possibles révèlent que trois zones recèlent un potentiel élevé susceptible de contenir d'importantes quantités de gaz naturel, soit 1) la zone gazéifère du Pliocène dans la région du bassin de Reine-Charlotte, 2) la zone gazéifère du Miocène de Reine-Charlotte et 3) la zone gazéifère d'origine structurale de Tofino.

Au total, le potentiel en pétrole et en gaz en place des six zones possibles du bassin de Reine-Charlotte (c.-à-d. les composantes pétrole et gaz des trois zones définies) est estimé respectivement à  $1\,560 \times 10^6 \text{ m}^3$  (9,8

x  $10^9$  barils) et à  $734 \times 10^9 \text{ m}^3$  ( $25,9 \times 10^{12} \text{ pi}^3$ ). Les ressources estimées en pétrole de trois zones du bassin de Reine-Charlotte sont légèrement inférieures aux ressources en gaz en termes de volumes énergétiques équivalents. La présence simultanée d'abondantes strates réservoirs, de bonnes roches mères de pétrole, de nombreux pièges structuraux de nature diverse et de nombreuses occurrences d'indices de pétrole et de gaz est indicatrice d'un important potentiel en accumulations de pétrole. Les zones les plus prometteuses se trouvent dans des strates néogènes à l'intérieur du bassin de Reine-Charlotte, sous la partie orientale de l'île Graham, et dans les régions de plate-forme extracôtières de l'entrée Dixon et des détroits d'Hecate et de la Reine-Charlotte. Selon nos prévisions, ces trois zones gazéifères contiendraient 41 champs de gaz non découverts renfermant des volumes de gaz en place de  $3 \times 10^9 \text{ m}^3$  ( $\sim 10^{11} \text{ pi}^3$ ) ou plus. En outre, on s'attend à ce que les trois zones pétrolifères contiennent deux champs renfermant plus de  $160 \times 10^6 \text{ m}^3$  ( $10^9$  barils) de pétrole initial en place. Un potentiel considérable a été reconnu dans au moins deux zones gazéifères présentes dans la région de Reine-Charlotte, ce qui rend la reprise des travaux d'exploration intéressante. L'évaluation en cours indique que la région du bassin de Reine-Charlotte renferme des ressources pétrolifères potentielles considérablement plus élevées que ce que l'on pensait (Procter et al., 1983).

Au total, le potentiel des trois zones possibles du bassin de Georgia est estimé à  $185 \times 10^9 \text{ m}^3$  ( $6,5 \times 10^{12} \text{ pi}^3$ ) de gaz en place. Les données géochimiques disponibles révèlent un potentiel en pétrole de peu d'importance dans les régions du bassin de Georgia ou de Tofino. Par conséquent, leurs zones pétrolifères n'ont pas été évaluées. On estime que deux zones gazéifères comptent 18 champs de gaz recelant des volumes en place supérieurs à  $3 \times 10^9 \text{ m}^3$  et que la zone structurale de Georgia du Crétacé renferme un potentiel considérable même si des risques substantiels lui sont attribués en raison des caractéristiques des réservoirs et de la roche mère. La présence de grandes structures fermées et la grande superficie de la zone en augmentent le potentiel.

Une seule zone a été définie dans la région du bassin de Tofino. Son potentiel s'élève à  $266 \times 10^9 \text{ m}^3$  ( $9,4 \times 10^{12} \text{ pi}^3$ ) de gaz en place. Selon les estimations, elle renferme 38 champs contenant un volume de plus de  $3 \times 10^9 \text{ m}^3$  de gaz, ce qui représente un potentiel considérable en gaz pour cette seule zone. Des risques importants liés aux faciès réservoirs et à la roche mère lui sont attribués, mais l'abondance de grandes structures dans l'ensemble d'une vaste région offre un cadre permettant d'effectuer une évaluation des ressources modérément favorable.

## INTRODUCTION

### Scope

Regional petroleum resource assessments have been prepared periodically for various sedimentary basins in Canada by the Geological Survey of Canada. These studies incorporate systematic basin analysis and statistical resource evaluations (Podruski et al., 1988; Wade et al., 1989; Sinclair et al., 1992; Reinson et al., 1993; Bird et al., 1994; Dixon et al., 1994). This report summarizes the assessment of oil and gas potential in the Queen Charlotte Basin and environs, as well as the Tofino, Winona, Juan de Fuca and Georgia basins surrounding Vancouver Island. These basins constitute the principal hydrocarbon-prospective sedimentary accumulations along the Pacific margin of Canada. It is important to note that while both oil and gas resources were evaluated in the Queen Charlotte Basin region, only gas plays were estimated in the remainder of the Pacific coast area. Geochemical data indicate the Georgia and Tofino basins are probably gas-prone, with little if any oil potential.

Based on geographic and tectonic considerations, the basins and sub-basins along the west coast of Canada were grouped into three general assessment regions: Queen Charlotte, Georgia, and Tofino. The Queen Charlotte assessment region includes the Queen Charlotte and Hecate basins and surrounding smaller sub-basins in Dixon Entrance and near Banks Island in onshore and continental shelf areas, and the deep-water Queen Charlotte Terrace west of the Queen Charlotte Islands (Fig. 1, 2). The Georgia assessment region includes the onshore-offshore Bellingham, Nanaimo, Comox and Suquash sub-basins and surrounding unnamed smaller sub-basins. These sub-basins encompass the Fraser River lowlands and delta, the Strait of Georgia, the Gulf Islands, and the eastern shore of Vancouver Island in southwestern British Columbia and Whatcom County in northwestern Washington State (Fig. 1, 2, 3). The Tofino assessment region includes the Tofino Basin and Juan de Fuca sub-basin beneath the continental shelf, and the deep-water Winona Basin and accretionary wedge seaward of the continental shelf edge (Fig. 1, 2). Some of the basins and sub-basins within the assessment regions extend into the United States (Washington State in the south, Alaska in the north). Relevant geological information from American parts of the basins were evaluated during the assessment studies, and total resource estimate numbers apply to the combined Canadian and American areas. Estimates for Canadian areas only are given later in the report.

### Purpose

The objective of this report is to provide an overview of the petroleum geology of Canada's west coast basins and to

present quantitative estimates of the oil and gas resources contained therein. This geological and resource framework will assist government agencies in evaluating land-use and moratorium issues, and petroleum industry companies in pursuing future exploration opportunities.

### Terminology

The terminology and procedures used in this report follow those outlined in Reinson et al. (1993) and are summarized below.


*Oil* is defined as any naturally occurring liquid that, at the conditions under which it is measured or estimated, is primarily composed of hydrocarbon molecules and is readily producible from a borehole.

*Natural gas* is defined as any gas (at standard pressure and temperature, 101.33 kPa and 15°C) of natural origin comprising mostly hydrocarbon molecules producible from a borehole (Potential Gas Committee, 1990). Natural gas may contain significant amounts of non-hydrocarbon gas such as H<sub>2</sub>S, CO<sub>2</sub> or He. In this study, non-hydrocarbon gas was not considered because of a lack of information on gas compositions in these basins.


*Raw gas* is unprocessed natural gas, containing methane, inert and acid gases, impurities and other hydrocarbons, some of which can be recovered as liquids. *Sales gas* or *marketable gas* is natural gas that meets specifications for end use. This usually requires processing that removes acid gases, impurities and hydrocarbon liquids. *Nonassociated gas* is natural gas that is not in contact with oil in a reservoir. *Associated gas* is natural gas that occurs in oil reservoirs as free gas. *Solution gas* is natural gas that is dissolved in crude oil in reservoirs. In this report, insufficient information is available to differentiate nonassociated, associated, and solution gas. All gas figures reported represent initial raw gas volumes.

*Resource* indicates all hydrocarbon accumulations known or inferred to exist. *Resource*, *resource endowment* and *endowment* are synonymous and can be used interchangeably. *Reserves* are that portion of the resource that have been discovered, whereas *potential* represents the portion of the resource not yet discovered but inferred to exist. The terms *potential* and *undiscovered resources* are synonymous and may be used interchangeably. Since no oil or gas pools or fields have yet been discovered in any west coast basin, all of the hydrocarbon volumes presented in this report represent potential or undiscovered resource figures.


*Gas-in-place* indicates the gas volume found in the ground, regardless of what portion is recoverable. *Initial in-place volume* is the gross volume of raw gas, before production. *Recoverable in-place volume* represents the



**Figure 1.** Geographic setting of the west coast region of Canada. Place names and geographic features mentioned in text are shown.



**Figure 2.** Regional setting and basin outlines of the west coast region of Canada. Assessment regions are shaded. Eastern edge of Hecate Basin modified from Haggart (1993). Well names are listed in Table 1 and cross-sections A-A', B-B' and E-E' are discussed in text.



**Figure 3.** Detail map of Georgia Basin in Vancouver area, the Fraser River lowlands and delta, southeastern Vancouver Island and Gulf Islands of British Columbia and northwestern Washington. Well names are listed in Table 2 and cross-sections C-C' and D-D' are discussed in text. Place names and geographic features mentioned in text are shown.

volume expected to be recovered with current technology and costs. These definitions can be applied to oil volumes as well.

A *prospect* is defined as an untested exploration target within a single stratigraphic interval; it may or may not contain hydrocarbons. A prospect is not synonymous with an undiscovered pool. An undiscovered pool is a prospect that contains hydrocarbons, but has not been tested as yet. A *pool* is defined as a discovered accumulation of oil or gas, typically within a single stratigraphic interval, that is separate, hydrodynamically or otherwise, from another hydrocarbon accumulation. A *field* consists of one or more oil and/or gas pools within a single structure or trap. Similar to most frontier regions, the assessment of west coast petroleum resources is based on estimates of field rather than pool sizes. A *play* is defined as a family of pools and/or prospects that share a common history of hydrocarbon

generation, migration, reservoir development and trap configuration.

Plays are grouped into two categories; *established* and *conceptual*. *Established plays* are demonstrated to exist by the discovery of pools with established reserves. *Conceptual plays* are those that have no discoveries or reserves, but which may exist, according to geological analyses. Established plays are categorized further into *mature* and *immature* plays depending on the adequacy of play data for statistical analysis. Mature plays are those plays that have sufficient numbers of discoveries within the discovery sequence so that the *discovery process model* of the PETRIMES assessment procedure is of practical use (Lee and Tzeng, 1989; Lee and Wang, 1990; Lee, 1993). Immature plays do not have a sufficient number of discoveries with established reserves to properly apply the model. Conceptual play analysis was applied exclusively in

**Table 1**  
**Well reference numbers in Queen Charlotte,**  
**Tofino, and Juan de Fuca basins**

<b>QUEEN CHARLOTTE BASIN (Fig. 2)</b>	
1	Bow Valley et al. Naden Harbour b-A27-J
2	British Columbia Coal Co. Tian Bay
3	Union Port Louis c-28-L
4	Queen Charlotte No. 1
5	Richfield-Mic Mac-Homestead Tow Hill d-93-C
6	Richfield-Mic Mac-Homestead Masset c-10-l
7	Richfield-Mic Mac-Homestead Nadu River b-69-A
8	Richfield-Mic Mac-Homestead Cape Ball d-41-L
9	Richfield-Mic Mac-Homestead Gold Creek c-56-H
10	Richfield-Mic Mac-Homestead Tiell c-56-D
11	Shell Anglo South Coho I-74
12	Shell Anglo Tyee N-39
13	Shell Anglo Sockeye B-10
14	Shell Anglo Sockeye E-66
15	Shell Anglo Murrelet L-15
16	Shell Anglo Auklet G-41
17	Shell Anglo Harlequin D-86
18	Shell Anglo Osprey D-36
<b>TOFINO BASIN</b>	
19	Shell Anglo Apollo J-14
20	Shell Anglo Zeus I-65
21	Shell Anglo Zeus D-14
22	Shell Anglo Pluto I-87
23	Shell Anglo Prometheus H-68
24	Shell Anglo Cygnet J-100
<b>JUAN DE FUCA SUB-BASIN</b>	
25	Merrill-Ring No. 1
26	R. D. Merrill No. 1
27	Merrill & Ring No. 25-1
28	Twin River State No. 30-1

this study because of the lack of any discovered pools with established reserves.

### Method and content

This report incorporates two essential components: geological basin analysis and statistical assessment. Basin analysis fundamentally describes and characterizes the exploration play. Fields and prospects in a play form a natural geological population that can be delimited areally. Once a play is defined, a numerical and statistical resource assessment is undertaken using field or prospect data from that specific play.

The analysis of oil and gas potential in the Queen Charlotte, Georgia, Tofino, Winona and Juan de Fuca basins entailed the delineation and the systematic evaluation of 10 conceptual petroleum plays. These plays are summarized with respect to play definition, geology, exploration history and estimated resource potential. This study is based on reviews of published and unpublished data and reports, interpretations and mapping from marine seismic reflection data, evaluation of well history records and logs, modelling of thermal maturation histories, and probabilistic analyses of the plays.

### Previous assessments

Based on early drilling results and initial accounts of the regions's geological setting (Sutherland Brown, 1968; Shouldice, 1971), a quantitative assessment of petroleum potential in the West Coast region was prepared by the Geological Survey of Canada (Haimala and Procter, 1982). That assessment produced estimates of recoverable petroleum resources of  $38.5 \times 10^6 \text{m}^3$  ( $241 \times 10^6$  bbl) of oil and  $265 \times 10^9 \text{m}^3$  (9.4 TCF) of gas (mean values). Slightly modifying the 1982 estimates, Procter et al. (1983) presented west coast petroleum potential estimates (average expectations) of  $50 \times 10^6 \text{m}^3$  ( $315 \times 10^6$  bbl) and  $270 \times 10^9 \text{m}^3$  (9.5 TCF) of recoverable oil and gas, respectively. By frontier basin standards, these estimates suggest a small resource potential.

Since publication of the early 1980s assessments, numerous studies have been undertaken in the region and considerable amounts of new geological and geophysical data collected. Qualitative assessments of petroleum potential in the Queen Charlotte Basin region have been presented in a number of papers incorporating results from the Geological Survey of Canada's Frontier Geoscience Program. Yorath (1987) stated that the Queen Charlotte Basin contains the highest potential for hydrocarbon accumulation; Georgia and Tofino basins have less potential as a result of porosity and source rock risks in the sedimentary succession. Gordy (1988) concluded that Georgia Basin is gas-prone with a high likelihood for the presence of closed hydrocarbon-bearing structures. Thompson et al. (1991) presented a generally positive view of petroleum potential by comparing the geology of the Queen Charlotte Islands area to the petroleum-producing Cook Inlet Basin of Alaska. In an assessment of the southern Queen Charlotte Islands and adjacent shelf areas, Dietrich et al. (1992) identified tracts of varying petroleum potential, including low potential for onshore and nearshore areas and moderate-to-high potential of offshore areas in Hecate Strait. Lyatsky and Haggart (1993) concluded that the Queen Charlotte Sound area has high petroleum potential, based on interpretations of regional distributions of reservoir and source rocks.

### Tectonic setting

Prior to general acceptance of the plate tectonic theory in the early 1970s, the Canadian Cordillera was divided into two belts using the classical geosynclinal hypothesis (Daly, 1912; Kay, 1951; White, 1959; King, 1969). An eastern belt consisting of sedimentary rocks with lesser volcanic and intrusive rocks was classified as a miogeosyncline, while the western belt, containing abundant volcanic, plutonic and metamorphic rocks, comprised the eugeosyncline. Knowledge of plate tectonic processes was used by Monger



**Table 2**  
**Well reference numbers in Georgia Basin**

<b>GEORGIA BASIN (Fig. 3)</b>			
<b>CANADA</b>			
29	BP Laurel Harmac c-36-F	43	Noble Francis No. 1
30	Novacorp Cedar b-29-F	44	Conoco Dynamic Mud Bay
31	BP Yellow Point d-84-C	45	Boundary Bay No. 3
32	Charter et al. Saturna No. 1	46	Richfield Pure Sunnyside 16-13
33	Gulf Ridge No. 1	47	Royal Can-Van Tor Big Horn Kuhn No. 1
34	Royal City No. 1	48	Outwest d-89-A
35	Fritts (Steveston well)	49	Conoco Dynamic Murray Creek
36	Great Basins No. 1	50	Port Haney
37	Richfield Pure Point Roberts 6-3-5	51	Siloam No. 1
38	Smith Savage No. 1	52	Empire No. 3
39	Spartan No. 2	53	Conoco Dynamic Stateside Campbell River
40	Spartan No. 1	54	Hercon Key Evans No. 1
41	Allenbee South Brazeau No. 1	55	Richfield Pure Abbotsford 16-17-16
42	Surrey Dome No. 1		
<b>UNITED STATES</b>			
56	Ridgeway-Heppner No. 1	104	Lingbloom No. 3
57	Ridgeway-Heppner No. 1A	105	Lingbloom No. 1 (Chamber of Commerce No. 1)
58	Ridgeway-Heppner No. 2	106	Peoples No. 1 (P.G.O. Lingbloom No. 1)
59	Kris Whatcom No. 1	107	Lingbloom No. 2 (Chamber of Commerce No. 2)
60	Stremler No. 2	108	Lingbloom No. 4 (Chamber of Commerce No. 4)
61	Lynden (Stremler) No. 1	109	Beyers No. 1
62	Ives No. 1	110	Bettsinger No. 2
63	Ives No. 2	111	Harden No. 1 (Hunter No. 1)
64	Thom No. 1	112	Bettsinger No. 1
65	Selien No. 1	113	Hunter No. 3
66	International No. 6	114	Harden No. 2 (Hunter No. 2)
67	AHEL Birch Bay No. 1	115	Chamber of Commerce No. 5
68	Heinrich No. 1	116	King No. 1 (Hale No. 1)
69	Johnson No. 1	117	Peoples No. 3
70	Seline No. 1	118	Peoples No. 4
71	Dahle No. 2	119	Peoples No. 6
72	Dahle No. 1	120	Peoples No. 5
73	Hillje No. 2	121	Shale Oil & Gas No. 1
74	Hillje No. 1	122	Hanson No. 1
75	Anderson	123	Holman Water Well No. 2
76	Home Birch Bay No. 1	124	Holman Water Well No. 1
77	Mills No. 1	125	Holman No. 3
78	Home No. 1	126	Water Well
79	Standard Ferndale (Community)	127	International No. 3
80	Hart No. 1	128	Jepson Water Well
81	Acme No. 1	129	Ridge No. 1
82	Acme No. 2	130	Ridge No. 2
83	Soderberg No. 1	131	Water Well
84	AHEL Terrell No. 1	132	Erickson Water Well
85	Sherman No. 1	133	Bellingham Natural Gas Company No. 1
86	Enterprise No. 1	134	Bellingham Natural Gas Company No. 2
87	Enterprise No. 2	135	Bellingham Natural Gas Company No. 3
88	Greenacres Water Well	136	Green Water Well
89	Sinnes Water Well	137	Water Well
90	International No. 5 (Laurel)	138	Barnhart Water Well
91	Hillebrecht No. 1	139	Water Well
92	International No. 4	140	Diamond Drill Hole
93	Russler No. 1	141	Molin No. 1
94	AHEL Ferndale No. 1	142	Ross No. 1
95	Lange No. 1	143	El Paso Ross No. 1
96	Lange No. 2	144	Can Am Squalicum No. 1
97	Lange Coal Test	145	Jensen No. 1
98	Livermore No. 1	146	Stewart-Hamilton
99	Cowden No. 1	147	Luce Water Well
100	Whatcom No. 2	148	Pelican Dome No. 1
101	Whatcom No. 1 (Lange No. 3)	149	Clark Water Well
102	North Coast No. 1	150	Happy Valley (Fairhaven)
103	Peoples No. 6		

et al. (1972) to identify five major morphogeological belts in the Cordillera. The five belts, from east to west, are Foreland, Omineca, Intermontane, Coast and Insular. Pertinent to this study are the Insular Belt and, to a lesser extent, the Coast Belt. Individual belts may encompass

numerous tectonic settings, such as ancient volcanic arcs, plutonic–metamorphic complexes, oceanic crust, shelf and slope deposits and successor basins. Coney et al. (1980) identified a series of fault-bounded lithotectonic terranes within the Cordillera that may be “suspect” with respect to

North American paleogeography. These exotic terranes are believed to have been accreted to the western continental margin of ancestral North America during the Mesozoic (Monger et al., 1982). Sometimes, these exotic terranes were amalgamated outboard of the continent to form superterranes prior to attachment to the continent. Subsequent to accretion, these terranes became disrupted along major dextral transcurrent faults. Tipper et al. (1981) divided the Canadian Cordillera into tectonic assemblages. These assemblages bounded by regional faults reflect a specific depositional or tectonic event. Wheeler and McFeely (1991) published a tectonic assemblage map of the Canadian Cordillera that incorporates the current interpretation of the regional tectonic setting. Basins containing thick sedimentary successions that formed during or subsequent to terrane accretion are important elements with respect to petroleum geology in the Cordillera.

The continental–oceanic plate boundary along the west coast of southern Vancouver Island is marked by the Cascadia subduction zone. North of Brooks Peninsula, on the northwest coast of Vancouver Island, the boundary is defined by a dextral transform fault displaying oblique convergence located on the continental shelf edge west of Queen Charlotte Sound and the Queen Charlotte Islands (Fig. 2). The plate boundary is located at or near the present-day shelf edge (approximated by the 500 m isobath, Fig. 1).

The Queen Charlotte Basin region encompasses the Insular Belt within the North American Plate and part of the Pacific Plate. The Queen Charlotte Terrace overlies Pacific Plate oceanic crust. The Georgia Basin region occurs entirely within the North American Plate, encompassing parts of the Insular Belt and adjacent Coast Belt. The Tofino Basin region encompasses the outermost parts of the North American Plate (including the Pacific Rim and Crescent terranes and other fossil accretionary wedges) and parts of the Explorer and Juan de Fuca plates. The Winona Basin and accretionary wedge overlie oceanic crust of the Explorer and Juan de Fuca plates, respectively.

The Insular Belt comprises several terranes, the largest of which are the Wrangellia and Alexander terranes. Alexander Terrane extends from the St. Elias Mountains in southwestern Yukon Territory to the Coast Mountains north and east of Hecate Strait. Wrangellia constitutes most of the remainder of the belt; that is, the remaining portion of the Queen Charlotte Island region and most of Vancouver Island. Wrangellia also exists in parts of the southwestern Coast Belt. Beneath western and southern Vancouver Island, the Pacific Rim and Crescent terranes have been emplaced beneath Wrangellia along major northwesterly and westerly thrust faults (Fig. 4). The Pacific Rim Terrane, comprising melanges and volcanic rocks, was emplaced against and beneath western and southern Wrangellia in the latest Cretaceous or earliest Tertiary. Subsequently, the ophiolitic

Crescent Terrane was accreted beneath the Pacific Rim Terrane commencing in the Late Eocene (Yorath, 1991). According to multichannel seismic profiles (Campbell et al., 1991), even younger terranes have been accreted to the base of Wrangellia. This emplacement and underplating of terranes along with subsequent and current accretion of the modern subduction complex has resulted in both uplift of western Vancouver Island and subsidence of Georgia Basin to the east (Campbell et al., 1991). Timing constraints imposed by apatite fission-track dating of the Cowichan fold and thrust system of southern Vancouver Island imply a temporal linkage between the formation of the fold and thrust system during the Middle Eocene and continued accretion of the Pacific Rim and Crescent terranes against Wrangellia at about 45 Ma (England et al., 1997).


## **RESOURCE ASSESSMENT PROCEDURE**

### **Geological play definition**

The definition of play type and play area are essential objectives of the geological basin analysis that precedes any numerical resource evaluation. A properly defined play will possess a single population of pools and/or prospects that satisfies the assumption that geological parameters within a play can be approximated by a family of lognormal distributions. A mixed population from an improperly defined play adds uncertainty to the resource estimate. Pools and/or prospects in a specific play form a natural geological population characterized by one or more of the following: age, depositional model, structural style, trapping mechanism, geometry, and diagenesis. Prospects or areas within a basin or region can be assigned to specific plays on the basis of a commonality of some or all of these geological elements.

### **Compilation of play data**

Since conceptual plays have no defined pools or discoveries, probability distributions of reservoir parameters such as prospect area, reservoir thickness, porosity, trap fill, and hydrocarbon fraction are needed. Prospect size can then be calculated using the standard “pool”-size equation. Seismic, well, and outcrop data prove particularly useful in identifying the limits for sizes of prospect area and reservoir thickness as well as porosity limits. Geochemical data are useful in identifying prospective areas as well as the composition of the hydrocarbon accumulations, that is, oil vs. gas proneness. Research in similar hydrocarbon-bearing basins is also important in order to provide reasonable constraints on reservoir parameters as well as contributing further information on other aspects of petroleum geology that may prove useful for the study.



**Figure 4.** Tertiary and Cretaceous onshore sedimentary basins, west coast region of Canada. Regional basements with respect to basins are illustrated. Example seismic lines are indicated illustrating structural styles in certain areas.

## Conceptual play analysis

There are several methods for estimating the quantity of hydrocarbons that may exist in a play, region or basin (White and Gehman, 1979; Masters, 1984; Rice, 1986; Lee, 1993). Petroleum assessments undertaken by the Geological Survey of Canada are currently based on probabilistic methods (Lee and Wang, 1990), developed in the Petroleum Exploration and Resource Evaluation System, PETRIMES (Lee and Tzeng, 1989). The conceptual hydrocarbon plays defined in the West Coast region were analyzed by applying a subjective probability approach to the reservoir parameters. The lognormal option in PETRIMES was utilized since experience indicates that geological populations of pool parameters can be adequately represented by lognormal distributions.

Conceptual resource assessments in the frontier regions use field-size estimates rather than pool-size predictions as derived from mature and immature play analysis. A field consists of one or more oil or gas pools or prospects in a single structure or trap. Probability distributions of oil and gas field sizes are computed by combining probability distributions of reservoir parameters, including prospect area, reservoir thickness, porosity, trap fill, hydrocarbon fraction, oil shrinkage, and gas expansion.

Probability distributions of oil and gas field sizes were combined with estimates of numbers of prospects (from seismic and play area mapping) and exploration risks, to calculate play potential and to estimate sizes of undiscovered fields.

Exploration risks at a play or prospect level are determined on the basis of the presence or adequacy of geological factors necessary for the formation of petroleum accumulations. Essential factors are reservoir, seal, source rock, timing of hydrocarbon generation, trap closure and preservation. Appropriate marginal probabilities are assigned to each geological parameter to obtain risk factors. All of the Queen Charlotte and Vancouver Island plays have a high probability of existing (low risk). Within each play, certain prospect-level risks are high and these are assigned appropriate risk factors. Exploration risk is an estimate, incorporating all risk factors, of the percentage of prospects within a play expected to contain hydrocarbon accumulations.


Because of the nature of conceptual assessment results and since no discovered pool sizes can be used to constrain sizes of undiscovered accumulations, the uncertainty of oil and gas play potential and pool size estimates for a given range of probabilities is necessarily greater than the limits derived by discovery process analysis used in assessing mature plays.

## QUEEN CHARLOTTE ASSESSMENT REGION


### Exploration history and regional studies

The major phase of offshore exploration in the region was undertaken by Shell Canada Ltd. from 1965 to 1969. Several thousand line-kilometres of marine seismic reflection data were acquired. Shouldice (1971) combined this seismic data with that from aeromagnetic surveys and studies of outcrop geology on the shoreline margins of the Queen Charlotte Basin to analyse and describe the region's geological setting. The overall basin geometry, the gentle folds within the Tertiary sedimentary rocks and the onlap of Tertiary sedimentary rocks onto basement were first interpreted from these data. Additional marine seismic surveys were carried out by Chevron Canada Ltd. in 1971. Stacey (1975) used gravity data obtained between 1963 and 1967 by the Canadian government to identify and interpret the geometry and relationships of Upper Tertiary sedimentary rocks to underlying Mesozoic volcanic and sedimentary rocks in the offshore Queen Charlotte Basin. One thousand kilometres of marine reflection data were recorded in 1988 by the Geological Survey of Canada in the Queen Charlotte Basin (*see* Fig. 4 for location of example lines; Fig. 5, 6, 7) (Rohr and Dietrich, 1990, 1991). Interpretation of these data inferred that the Tertiary sedimentary succession is highly varied in thickness and the basin fill and underlying basement are extensively faulted to form a complex pattern of sub-basins and half-grabens (Rohr and Dietrich, 1992) (Fig. 5, 6, 7). These structural features imply a transtensional tectonic setting. A seismic refraction survey (Spence et al., 1991), was also carried out in 1988 by the Geological Survey of Canada. These data were used to estimate crustal thickness in Queen Charlotte Sound. Lyatsky (1991) combined regional magnetic and gravity surveys with physiographic lineaments to construe that the lateral crustal movements accompanying the formation of Tertiary Queen Charlotte Basin were probably small and consequently, no rift-related structures were formed. Lyatsky concluded that a fault-block tectonic pattern likely dominated during the Mesozoic and Cenozoic in the area. An integrated geophysical approach was utilized in the Queen Charlotte Basin by Lowe and Dehler (1995) to calculate crustal thickness. They considered both the deep marine seismic reflection and refraction data collected by the Geological Survey of Canada in 1988 and integrated it with gravity and bathymetry data.


The first petroleum exploration well in the Queen Charlotte Basin region (Tian Bay) was drilled in 1913 on the west coast of Graham Island (*see* Fig. 2 for well locations and Table 1 for well listings). Eight more wells were drilled onshore (Graham Island) between 1949 and 1971 (Fig. 2). Richfield Oil Corporation completed six of the wells. The Royalite Oil Company and Union Oil Company each drilled



**Figure 5.** Seismic reflection profile, Dixon Entrance, northern Queen Charlotte Basin (data from Rohr and Dietrich, 1990). Neogene half-grabens and tilted fault blocks are associated with down-to-east normal faults. Neogene sedimentary strata overlie a variable Mesozoic complex of sedimentary, volcanic and plutonic rocks.



**Figure 6.** Seismic reflection profile, Hecate Strait, central Queen Charlotte Basin (data from Rohr and Dietrich, 1990). Structural features include Mesozoic half-grabens and rift blocks [and associated normal faults (1)], a Late Miocene unconformity [apparent at profile's east end (2)], and Pliocene reverse faults and inversion folds (3). Solid circle indicates stratigraphic position of oil show (in Miocene sandstones) encountered by the Sockeye B-10 well. Star indicates position of interpreted hydrocarbon indicators.



**Figure 7.** Seismic reflection profile, Queen Charlotte Sound, southern Queen Charlotte Basin (data from Rohr and Dietrich, 1990). Miocene half-grabens and tilted fault blocks are overlain by relatively undeformed Pliocene strata. The Harlequin well encountered numerous thick sandstones within the Neogene section, many of excellent reservoir quality.

one well (Table 1). Eight offshore wells were completed by Shell Canada Ltd. between 1965 and 1969 (Table 1). Six were drilled in Hecate Strait (Coho, Tyee, Sockeye B-10 and E-66, Murrelet and Auklet) (Shell Canada Ltd., 1968a, b, c, e, 1969b, c) and two in Queen Charlotte Sound (Harlequin and Osprey) (Fig. 2) (Shell Canada Ltd., 1968d, 1969a). In 1972, the Canadian federal government imposed an indefinite moratorium on petroleum exploration of federal lands in the Pacific offshore in response to environmental concerns. Since that time, petroleum exploration has been limited in the area, with only one onshore well completed on Graham Island (Bow Valley Industries, 1984, Naden well).

## Regional geology

### *Geological setting and tectonic evolution*

The Queen Charlotte and Hecate basins were previously interpreted as overlying both Wrangellia and Alexander terranes (Yorath and Chase, 1981). The Paleozoic to Lower Mesozoic Wrangellia and Alexander terranes accreted to the North American Plate margin in mid-Jurassic time (van der Heyden, 1992). The location of the boundary between the two terranes has important implications for petroleum potential because Wrangellia Terrane contains the region's principal oil source rocks. More recent studies addressing the terrane boundary question indicate that Wrangellian rocks probably occur across most or all of the assessment region (Woodsworth, 1988; Thompson et al., 1991; Wheeler et al., 1991). Hecate Basin is a plate-margin parallel, Cretaceous forearc basin (Dietrich, 1995), that developed in response to post-terrace accretion convergence and

orthogonal subduction of the Pacific Plate beneath the continental margin. Basin-fill sediments were derived from uplifted coast mountains to the east. Hecate Basin underlies part of the Queen Charlotte Basin and all of Queen Charlotte Islands (Fig. 2). The Queen Charlotte Basin is an upper Tertiary strike-slip basin that developed in response to transtensional and transpressional Pacific Plate–North American Plate interactions. Queen Charlotte basin-fill sediments were derived from variable source areas and directions. The Queen Charlotte Terrace is a Plio-Pleistocene sedimentary prism that developed in response to oblique transpression of the Pacific Plate against the Queen Charlotte Islands crustal block (Prims et al., 1997).


The Neogene Queen Charlotte Basin is the largest basin on the west coast, encompassing some 40,000 km<sup>2</sup>. The basin underlies eastern Graham Island and large portions of the continental shelf of Dixon Entrance, Hecate Strait and Queen Charlotte Sound. Northernmost parts of the basin extend into offshore areas of southeast Alaska (Risley et al., 1992). The basin is underlain by Mesozoic and Tertiary volcanic, plutonic and sedimentary rocks, with a geological history linked to evolution of the Pacific continental margin and associated convergent, transcurrent, and possible extensional plate interactions (Lewis et al., 1991; Rohr and Dietrich, 1992; Rohr and Currie, 1997). In detail, the Queen Charlotte Basin consists of a series of separate or partly coalesced strike-slip sub-basins, which developed across a 150 km wide, plate-margin-parallel shear zone (Rohr and Dietrich, 1992; Dietrich, 1995). The sub-basins are commonly half grabens bound by northwest- or north-trending normal or oblique-slip faults (Fig. 5, 6, 7). The sub-basins contain syn-rift clastic sediments, and local volcanic

flows, characterized by considerable local variability in lithology and thickness. Overlying these deformed sedimentary rocks is a Pliocene sequence that is relatively undisturbed. These rocks drape the deeper structures and are more laterally continuous. Relatively recent compression and shortening in Hecate Strait is reflected by evidence of reactivation of the block faults and deformation of the sedimentary rocks. Some erosional truncation of folds at the seafloor suggests very recent deformation (Rohr and Dietrich, 1991).

### Stratigraphy and structure

Volcanic rocks of the Wrangellian Triassic Karmutsen Formation are several thousand metres thick and represent “basement” for petroleum exploration in the Queen

Charlotte Basin region (Fig. 8). On Queen Charlotte Islands, the Karmutsen Formation is conformably overlain by up to 1000 m of Upper Triassic and Lower Jurassic limestones, sandstones, and shales of the Kunga and Maude groups (Fig. 8). The Kunga Group consists of 200 m of massive carbonate overlain by thin-bedded fossiliferous limestone (Monger et al., 1991). In the uppermost Kunga, the limestone is overlain by Lower Jurassic argillite with lesser interbedded sandstone and tuff. This unit is up to 400 m thick and is similar to the coeval Harbledown Formation on Vancouver Island. The Lower and Middle Jurassic Maude Group consists of 400 m of shale, shaly limestone, coquinooid sandstone and minor tuff and tuffaceous siltstone. Upper Kunga limestone and argillite and lower Maude shale contain oil source rocks. These rocks are deposited in a stable shelf setting. Principal structures in Lower Mesozoic rocks are Middle Jurassic northwest-trending folds and



**Figure 8.** Simplified stratigraphic column for Queen Charlotte, Georgia, Tofino and Juan de Fuca sub-basin regions (modified from Niem and Snavelly, 1991; Haggart, 1992, 1993; Mustard and Rouse, 1994). Potential reservoir strata occur in the Skonun Formation and at the base of the Queen Charlotte succession in Queen Charlotte Basin, in coarse clastics of the Huntingdon Formation and Nanaimo Group in Georgia Basin, and in thin coarse clastic intervals in Tofino and Juan de Fuca basins. Petroleum source rocks occur in upper Kunga–lower Maude, lower Skonun, and (?) Upper Queen Charlotte Group strata in Queen Charlotte Basin. Considerable variability exists in the distribution and thickness of all stratigraphic units.

thrust faults (Thompson et al., 1991; Lewis et al., 1991). Karmutsen, Kunga and Maude rocks accumulated in intra-oceanic and island arc settings, as part of the exotic terrane, Wrangellia.

The Wrangellian (Lower Mesozoic) succession is unconformably overlain by several hundred metres of volcanic and volcanoclastic rocks of the Middle Jurassic Yakoun and Moresby groups (Fig. 8). The Yakoun Group comprises 480 m of volcanic breccia, lapilli tuff, and agglomerate with minor lenticular sandstone and siltstone (Monger et al., 1991). The overlying siliciclastic sedimentary rocks of the Middle Jurassic Moresby Group consist of siltstone, shale, and sandstone with minor pebble conglomerate. The succession varies in thickness from 45 to 200 m. Middle Jurassic and older rocks are locally intruded by late Middle to Late Jurassic plutons of intermediate to felsic compositions. These plutonic bodies are aligned on northwesterly and northerly trends (Sutherland Brown, 1968). They range in composition from hornblende diorite to quartz diorite and dip steeply eastward. The enclosing host rocks, specifically Karmutsen Formation, are metamorphosed to fine-grained amphibolites in restricted contact aureoles (Yorath et al., 1991). Unconformably overlying Moresby–Yakoun rocks are up to 2500 m of Upper Jurassic–Cretaceous sandstone, shale and conglomerate of the White Point beds and Queen Charlotte Group. The White Point beds, also known as the ‘Longarm Formation’ (Haggart, 1993), have both a proximal and a distal facies. Proximal facies-type rocks present in southeastern Moresby Island consist of boulder, pebble and granule conglomerate as well as coarse-grained sandstone. The unit is about 180 m thick. There is a pronounced angular unconformity between the White Point beds and underlying Kunga Group rocks in the area. Andesitic clasts similar in composition to the Yakoun Group are found in the conglomerate. Distal facies of the 500 m thick succession include calcareous siltstone, fine- to medium-grained greywacke and argillite. The greywacke and argillite represent deep-water turbidites. There are also minor andesitic agglomerates and porphyritic flows within the succession. Overlying the White Point beds, probably unconformably, is the mid- to Upper Cretaceous Queen Charlotte Group consisting of shale, sandstone, conglomerate, and minor coaly fragments. The basal part of the Queen Charlotte Group consists of a time-transgressive succession (up to 200 m thick) of shallow-marine sandstone and granule conglomerate (Haggart, 1991). Some sandstone and conglomerate in this unit have good reservoir characteristics and are potential petroleum reservoirs. The coarse-grained deposits are overlain by several hundred metres of siltstone and carbonaceous shale. Upper parts of the Queen Charlotte Group locally contain volcanic rocks and conglomeratic beds. Structures in Jurassic and Cretaceous strata include northwest-trending contractional folds of early Tertiary age (Lewis et al., 1991).

The Mesozoic succession is locally intruded by Late Eocene to Oligocene plutons and overlain by Upper Eocene to Pliocene clastic sedimentary strata and volcanic rocks (Fig. 8). Unnamed Eocene–Oligocene sedimentary and volcanic rocks form a minor component of the lower Tertiary succession, attaining a maximum thickness of a few hundred metres. Sandstone, shale, argillite, and coal with laterally equivalent volcanic rocks are present in this unit.

Neogene volcanic and sedimentary rocks of the Masset and Skonun formations unconformably overlie the Paleogene and older rocks in the Queen Charlotte Basin region and comprise the bulk of the upper Tertiary succession. Neogene volcanic rocks range up to 2000 m in thickness. Basalt and rhyolite flows, pyroclastics and related intrusions constitute the Masset Formation. The Masset Formation is both overlain by and locally interbedded with the Neogene Skonun Formation. The Skonun Formation consists of interbedded sandstone, shale, conglomerate and coal, and is up to 6000 m in thickness in some offshore locations. These rocks were deposited in both marine and nonmarine settings throughout the basin. Structural features within the Queen Charlotte Basin developed in association with Miocene transtensional and Plio-Pleistocene transpressional tectonics (Rohr and Dietrich, 1992). Miocene structures include north- and northwest-aligned normal and oblique-slip faults (Fig. 5, 6, 7, 9; see Fig. 2 for cross-section locations and Fig. 4 for seismic line locations). Pliocene structures in basin-fill strata include reverse faults (commonly developed as inversions of Miocene normal faults), contractional folds and combination fault–fold flower structures (Fig. 6, 9). Pleistocene structural features in the Queen Charlotte Basin and Queen Charlotte Terrace include local folds and tilted, truncated Neogene strata. Pleistocene folding of strata in the Queen Charlotte Terrace occurred in association with transpression of the Pacific Plate against the Queen Charlotte Islands.


## **Petroleum geology**

### ***Reservoirs***

#### ***Mesozoic***

The Upper Jurassic–Cretaceous White Point beds and Queen Charlotte Group (Fig. 8) contain thick sections of sandstone and conglomerate, portions of which have reservoir potential. The best reservoir qualities occur in shallow-marine sandstones and granule conglomerates within the basal part of the Queen Charlotte Group. This time-transgressive unit, referred to as “basal transgressive lithofacies” (Haggart, 1991), contains texturally mature, arkosic sandstone (compositional data in Sutherland Brown, 1968; Fogarassy and Barnes, 1991). The basal transgressive





**Figure 9.** Geological cross-sections (derived from seismic and well data) across Hecate Strait (A-A') and Queen Charlotte Sound (B-B'; locations in Fig. 2). Numbers (1 to 5) indicate possible petroleum trap types within and beneath the Neogene Queen Charlotte Basin (from Dietrich, 1995).

strata were deposited along northwest–southeast aligned paleoshorelines in the Queen Charlotte Islands area and probably the western parts of Dixon Entrance, Hecate Strait and Queen Charlotte Sound (Haggart, 1991; Lyatsky and Haggart, 1993). Fogarassy and Barnes (1991) described the reservoir characteristics of outcrop sections of the basal lithofacies on the Queen Charlotte Islands. Porosity averages 5 to 10%, with values locally exceeding 15%, in the 30 to 190 m thick unit. The observed porosity is a combination of preserved intergranular porosity and appreciable secondary porosity due to calcite dissolution. Permeability is considered fair to good as a result of well-rounded framework grains and lack of clay cements.

Other marine sandstone and conglomerate units within the Upper Jurassic–Cretaceous succession, although of considerable thickness locally, are characterized by more lithic (commonly volcanic) compositions and have generally poor reservoir potential (Fogarassy and Barnes, 1991). Cretaceous nonmarine strata occur in the subsurface along the east side of Hecate Strait (Fig. 1). Three offshore wells in Hecate Strait (Tye and Sockeye B-10 and E-66; Fig. 2, 6, 9) penetrated probable Upper Cretaceous coal-bearing nonmarine strata (Haggart, 1991; J. M. White, pers. comm., 1992). Sandstones are abundant in the nonmarine Cretaceous sections penetrated in offshore wells, but in all cases well logs indicate low porosity and permeability (Shell Canada Ltd., 1968a, b, c). Although known surface and subsurface occurrences of Cretaceous sandstone (apart from basal transgressive units) are characterized by poor reservoir qualities, Cretaceous strata in parts of Hecate Basin may have improved reservoir potential associated with local developments of secondary porosity (Fogarassy and Barnes, 1991).

### *Paleogene*

Paleogene sedimentary strata in the western Queen Charlotte Basin region may have limited reservoir potential. The Port Louis well on western Graham Island (Fig. 2) penetrated a 575 m thick section of Eocene–Oligocene volcanoclastic sandstone, conglomerate, and shale beneath Masset Formation volcanic rocks (White, 1991). Similar sedimentary strata of probable Paleogene age occur in outcrop on Hippa Island, southwest of the Port Louis well location (not shown on Fig. 1; Higgs, 1989; Lewis et al., 1991). Sandstones and conglomerates in both areas are characterized by poor reservoir qualities, as a result of occlusion of pores by decomposition products of volcanic and feldspar grains (Higgs, 1989). Three drillstem tests of low porosity sandstone in the Port Louis well had insignificant recoveries (Union Oil Company of Canada Ltd., 1971). Paleogene sedimentary strata undoubtedly occur beneath Masset Formation volcanic rocks in other parts of Graham Island and perhaps elsewhere in the region. These

rocks could have locally improved reservoir potential if sandstone compositions differ from known sections (Higgs, 1989). Until additional subsurface control is available, however, assessments of Paleogene reservoir potential will remain equivocal.

### *Neogene*

The Neogene Skonun Formation (Fig. 8) contains large volumes of sandstone and conglomerate that were deposited in a wide range of nonmarine and marine depositional environments, including alluvial fan, fan delta, delta plain, shelf, and marine slope settings (Higgs, 1991; Dietrich et al., 1993). Most or all Neogene sub-basins contain a mix of nonmarine and marine strata, with northern sub-basins beneath Graham Island and Hecate Strait containing larger percentages of nonmarine deposits. The wide geographic distribution, large volume and commonly good reservoir characteristics of Skonun sandstones and conglomerates make them the principal petroleum exploration targets in the region. Skonun sandstone and conglomerate beds are up to tens of metres thick, comprise 25 to 75 per cent of the total Neogene sedimentary section, and attain cumulative thicknesses of up to 2000 m in some sub-basins. In the 14 exploration wells that penetrated Skonun strata, only five drillstem tests were completed, all in onshore wells, with the best flow rates recorded from the Tow Hill well (450 barrels of water/day; Richfield Oil Corporation, 1958).

Skonun sandstone is predominantly arkose and lithic arkose and generally has the highest mineralogical maturity of all arenaceous rocks in the region (Shouldice, 1971). At depths of less than 2000 m, Skonun sandstone has generally good reservoir qualities, characterized by very high porosity (25 to 35%) and fair to very good permeability (10 to 1500 md) (Dietrich, 1995, fig. 5, 6). At depths from 2000 to 3000 m, Skonun sandstones have high porosity (20 to 30%) and fair permeability (10 to 100 md) (Dietrich, 1995, fig. 5, 7). Skonun reservoir potential is limited below depths of about 3000 m, because of low permeability.

The wide range of permeability in Skonun sandstone and limited correlation between permeability and porosity (Dietrich, 1995, fig. 5) reflect composition-related diagenetic variations. Less permeable sandstone contains greater amounts of pore-bridging clays, products of feldspar grain decomposition (Shouldice, 1971; Fogarassy and Barnes, 1988). Tidal-shelf and storm-dominated shelf sandstone is commonly the most permeable Skonun reservoirs, as a result of advanced textural and compositional maturity (Higgs, 1991). Shelf sandstone occurs throughout the Queen Charlotte Basin and is abundant in the southern Queen Charlotte Sound half of the basin and in post-rift sections in the northern half of the basin.

In some areas, Neogene sections contain volcanoclastic sandstone and conglomerate characterized by poor reservoir qualities, even at shallow depths. Low porosity and permeability in volcanoclastic strata are due to advanced cementation associated with diagenesis of volcanic rock fragments (Galloway, 1974). Volcanic and volcanoclastic rocks occur in some (but not all) sub-basins and, where present, usually occur in the oldest syn-rift sections adjacent to basin-bounding faults (Fig. 9). From a regional perspective, volcanoclastic strata comprise a relatively small portion of the basin's total sediment volume.

### Seals

Cretaceous and Tertiary reservoir strata are interstratified with shale, siltstone, and volcanic rocks that may provide effective seals for petroleum accumulations. Basal Cretaceous sandstone units are commonly overlain by thick Upper Cretaceous shale sections. In the Graham Island area and parts of the offshore, Neogene volcanic rocks may form seals for Cretaceous or Paleogene reservoirs. The Queen Charlotte Basin contains thick successions (commonly between 1000 and 3000 m depth) of interbedded Neogene reservoir and seal rocks, providing potential for entrapment of oil or gas at many stratigraphic intervals. Impermeable strata are least common in shallow Neogene sections (above 1000 m depth) in the northern half of the basin. Fault-related seals for Tertiary or Cretaceous reservoirs may be associated with impermeable fault zones or crossfault juxtaposition of permeable and impermeable strata (Fig. 6, 9).

### Traps

Table 3 lists the various petroleum trap types expected to occur within the west coast basins according to stratigraphic succession. A variety of potential structural and stratigraphic petroleum traps occurs in Cretaceous and Tertiary strata within and beneath the Queen Charlotte Basin (Fig. 6; 1 to 5 in Fig. 9). Traps involving Cretaceous strata include Late Cretaceous folds and extensional fault blocks (4 and 5 in Fig. 9). The fault-block traps may be associated with Cretaceous subcrop patterns at the sub-Neogene unconformity (example in Fig. 6, at 2.5s). Neogene reservoir strata occur in Miocene to Pleistocene structural and stratigraphic traps. Miocene structures include tilted fault blocks, fault-related rollover and drag closures and low-relief drape anticlines (1 in Fig. 6; 2 in Fig. 9). Stratigraphic traps may occur in Miocene strata that onlap or subcrop below local unconformities, or within cone-shaped alluvial fan or fan delta deposits banked against fault scarps or pinching out updip within half-grabens (2 in Fig. 6; 3 in Fig. 9). Pliocene structural traps, restricted to the northern half of the basin (Hecate Strait and Dixon Entrance), include abundant large-amplitude folds, commonly cut by steep-dipping reverse faults (3 in Fig. 6; 1 in Fig. 9). Most of the Late Pliocene folds in Neogene strata (including the Sockeye anticline, Fig. 6), are structurally detached from underlying Mesozoic rocks. Structural decoupling results in lateral separation of closures in Neogene and Cretaceous strata. Stratigraphic traps may also be locally present in shallow parts of the Queen Charlotte Basin where tilted Neogene strata are unconformably overlain by Quaternary mudstone.

**Table 3**  
**Petroleum trap types in west coast basins**

Assessment region	Petroleum trap types
Queen Charlotte	
Cretaceous	simple compressional anticline, structural complex subthrust fault (rift blocks)
Miocene	simple compressional anticline, faulted anticline (fault-related rollover), thrust fault (drag closure), tilted fault block, subcrop unconformity, onlap unconformity, alluvial fan against fault scarp, porosity/permeability pinchout
Pliocene	simple compressional anticline, faulted anticline (reverse fault), subcrop unconformity
Georgia	
Cretaceous	simple compressional anticline, faulted anticline (fault-related rollover), normal fault, thrust fault (drag closure), porosity/permeability pinchout, subcrop unconformity
Tertiary	simple compressional anticline, faulted anticline (fault-related rollover), normal fault, thrust fault (drag closure), porosity/permeability pinchout, subcrop unconformity
Pleistocene	porosity/permeability pinchout
Tofino	
Tertiary	simple compressional anticline, faulted anticline (fault-related rollover), normal fault, thrust fault (drag closure), shale diapir, porosity/permeability pinchout

Based on seismic mapping and outcrop information and related extrapolations into limited-data areas, the number of structural traps alone within the assessment region is estimated to be in the hundreds. The largest structural closures involve Neogene strata within Pliocene folds, some of which exceed 50 km<sup>2</sup> in area.

### Source rocks

Good to excellent hydrocarbon source rocks have been identified in Upper Triassic–Lower Jurassic Kunga and Maude strata (Fig. 8), from analyses of outcrop and shallow onshore corehole samples (Macauley, 1983; Vellutini and Bustin, 1991a). Upper Kunga limestone and thin-bedded argillite and lower Maude shale contain oil-prone Type I and oil-and gas-prone Type II organic matter, with total organic content (TOC) averaging 1 to 4% in sections up to several hundred metres thick. Organic-rich shale with 5 to 10% TOC occurs in beds up to 10 m thick. Measured hydrocarbon yields from organic-rich beds from the central Queen Charlotte Islands area are up to 50 to 100 mg HC/g rock, indicating excellent oil source potential (Macauley, 1983). The subsurface distribution of Kunga–Maude strata is largely unknown, but is expected to be highly irregular, because of varying effects of episodic Middle Jurassic to Tertiary erosion. Cretaceous uplift and erosion was probably widespread in areas close to or landward (east) of the Hecate Basin margin (Fig. 2). Kunga–Maude strata are most likely preserved in greatest abundance in the southwestern half of the region, beneath Graham Island and western parts of Dixon Entrance, Hecate Strait and Queen Charlotte Sound (Thompson et al., 1991; Lyatsky and Haggart, 1993).


Sedimentary strata within the Upper Jurassic–Cretaceous succession contain Type III (gas-prone) organic matter, with generally poor hydrocarbon source potential (TOC less than 1%; Vellutini and Bustin, 1991a). In offshore areas, carbonaceous beds and coal seams in nonmarine Upper Cretaceous strata may have some gas potential (Fig. 8).

The Neogene Skonun Formation and unnamed upper Paleogene strata (Fig. 8) contain coal beds and dispersed Type III organic matter, with good gas and fair to good oil source potential (Bustin et al., 1990; Vellutini and Bustin, 1991a). Organic content in Skonun strata averages 0.5 to 1.5% TOC, with higher TOC values (5 to 25%) occurring in coal-bearing zones. Coal beds are abundant in the northern half of the Queen Charlotte Basin, where nonmarine deposits are thick and widespread. Neogene coal and carbonaceous beds locally contain resinite (fossil tree sap), a potential source of oil and condensate in otherwise gas-prone strata (Snowdon et al., 1988). Skonun shale and siltstone locally contain Type II organic matter, with up to 2.5% TOC, with good oil and gas source potential (Vellutini and Bustin,

1991a). Overall, Tertiary strata are lower in source rock quality than Kunga–Maude rocks, but occur in greater volume and distribution.


### Source-rock maturation

Present-day thermal maturation conditions of Mesozoic and Tertiary strata are known from pyrolysis TMAX data (Bustin et al., 1990), vitrinite reflectance measurements (Vellutini and Bustin, 1991b) and colour conodont alteration index values (Orchard and Forster, 1991) from outcrop and well samples. For the following discussion (and Fig. 10, 11), oil and gas windows are defined by vitrinite reflectance levels of 0.5 to 1.3% Ro and 0.5 to 2.6% Ro, respectively. References to mature source rocks imply oil window maturation levels for Kunga–Maude strata (Types I–II organic matter) and gas window maturation levels for Cretaceous and Tertiary strata (Type III organic matter).



**Figure 10.** Maturation–depth profile of strata in the Sockeye B-10 location, Queen Charlotte Basin (see Fig. 2 and Table 1 for well location). Mesozoic stratigraphy below well TD has been inferred from seismic data and onshore geology. Measured maturation values from Yorath and Hyndman (1983) and Bustin et al. (1990), the latter as vitrinite reflectance equivalent of pyrolysis TMAX data. Model-predicted maturation profile based on illustrated stratigraphy and heat-flow model depicted in Figure 14. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

Kunga–Maude rocks are overmature on the southwestern Queen Charlotte Islands, because of proximity to Jurassic and Tertiary plutons and dyke swarms (Orchard and Forster, 1991). In contrast, Kunga–Maude strata are marginally mature to mature on the central and northern Queen Charlotte Islands, reflecting reduced effects of magmatic heating (Vellutini and Bustin, 1991b). Maturation levels of Kunga–Maude strata in offshore areas are unknown, but are expected to vary from mature to overmature. As examples, measured and model-predicted maturation profiles for two offshore wells (Sockeye B-10 and E-66; Fig. 10, 11) indicate that Kunga–Maude strata, if present below well drill-depths, will be overmature at the B-10 location (Fig. 10) and mature to overmature at the E-66 location (Fig. 11). Model-predicted maturation differences in Mesozoic strata in the two Sockeye wells reflect differences in burial depths (Fig. 10, 11). For the Sockeye area and probably many parts of central Hecate Strait, these maturation trends indicate Kunga–Maude sections will be within the oil window at depths above 3000 m. For other offshore areas, including Queen Charlotte Sound, subsurface well data are insufficient to estimate maturation levels for Kunga–Maude strata.




**Figure 11.** Maturation–depth profile of strata in the Sockeye E-66 location, Queen Charlotte Basin (see Fig. 2 and Table 1 for well location). Mesozoic stratigraphy below well TD has been inferred from seismic data and onshore geology. Measured maturation values are vitrinite reflectance equivalent of pyrolysis TMAX data. Model-predicted maturation profile based on illustrated stratigraphy and heat-flow model depicted in Figure 14. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

Neogene strata within the Queen Charlotte Basin are immature to mature, with an estimated 30 to 40 per cent of the total basin fill occurring at maturation levels within oil or gas generation windows. The depth to the top of the hydrocarbon generation window (0.5% Ro) in Neogene strata occurs typically at 2000 to 2500 m. In the Sockeye B-10 well, coal-bearing Skonun strata are mature from depths of about 2000 m to the base of the formation at 4300 m; the base of the oil window occurs at about 3700 m (Fig. 10). Skonun strata are marginally mature at the base of the Sockeye E-66 well (Fig. 11). Other offshore wells exhibit similar depth-maturation trends in Neogene Skonun strata (R.M. Bustin, pers. comm., 1993).


### Timing of hydrocarbon generation

Maturation of source rock within the Queen Charlotte Basin region was controlled to varying degrees by heat flow associated with Jurassic and Tertiary magmatism, and Late Tertiary rifting and subsidence. Temporal and spatial variations in heat flow resulted in substantial variability in timing of source rock maturation and hydrocarbon generation across the region. Time-temperature modelling of stratigraphic sections from different parts of the Queen Charlotte Islands (Vellutini and Bustin, 1991b) indicates Kunga–Maude strata entered the oil generation window at various times from Late Jurassic to Late Miocene, with a general geographic trend to younger times for oil generation onset in the central and the northeastern parts of the archipelago. This reflects greater distances from plutonic and volcanic centres. Paleogene sedimentary strata entered the oil window in the Pliocene in the western Graham Island area (wells 1, 2, and 3, Fig. 2). With the exception of the deepest part of the Tow Hill well (well 5 of Fig. 2, Table 1), Neogene strata penetrated by onshore wells along eastern Graham Island (wells 4, 6–10) have yet to enter the oil window.

Similar variations in maturation history and timing of hydrocarbon generation occurred offshore. As part of this study, subsidence and hydrocarbon generation models were calculated for a number of different offshore locations, using known or interpreted stratigraphy from well and seismic data. Two of the models (Fig. 12, 13) serve to illustrate how timing of hydrocarbon generation from Kunga–Maude and Skonun source rock units may have varied within a local area. The first model (Fig. 12) is based on the Sockeye B-10 well location, where 4500 m of Miocene strata unconformably overlie Mesozoic rocks (Fig. 10). The second model is based on a seismically defined location near the Sockeye E-66 well (Fig. 13) where 2000 m of Neogene strata unconformably overlie the flank of a Mesozoic fault block. The heat-flow history (Fig. 14) used for hydrocarbon generation modelling was derived by iteratively matching kinetic-model-calculated and observed maturation



**Figure 12.** Subsidence and hydrocarbon generation model for the area of Sockeye B-10 well, central Hecate Strait, Queen Charlotte Basin. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995). Hydrocarbon generation models based on standard (BasinMod) kinetic parameters for organic matter types (Types I-II in Kunga-Maude strata and Type III in Skonun strata). See Figure 14 for heat-flow input of this model. See text for discussion.




**Figure 13.** Subsidence and hydrocarbon generation model for the area of Sockeye E-66 well, central Hecate Strait, Queen Charlotte Basin. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995). Hydrocarbon generation models based on standard (BasinMod) kinetic parameters for organic matter types (Types I-II in Kunga-Maude strata and Type III in Skonun strata). See Figure 14 for heat-flow input of this model. See text for discussion.

conditions in the Sockeye B-10 well, the deepest well in the region (Fig. 10). The best-fit heat-flow model was temporally variable, with a critical thermal control on maturation being associated with a Late Tertiary period of rifting and high heat flow (Fig. 14). The Sockeye area hydrocarbon generation models (Fig. 12, 13) derived from this heat-flow model illustrate dramatically different results. The Sockeye B-10 model indicates hydrocarbon generation occurred from middle Jurassic to earliest Miocene time in Kunga-Maude strata and from early Miocene to Recent time in lower Skonun strata. In contrast, the Sockeye E-66 model (10 km from Sockeye model B-10) indicates Kunga-Maude hydrocarbon generation from early Miocene to Recent time and no Skonun hydrocarbon generation. These hydrocarbon generation models, and others not illustrated here, indicate the potential for a variable and complex hydrocarbon charge history, even within local areas.

### Hydrocarbon shows

The potential for significant petroleum accumulations in the Queen Charlotte Basin region is perhaps best demonstrated by the common occurrence of oil and gas shows. Over 50



**Figure 14.** Heat-flow model for central Hecate Strait, Queen Charlotte Basin (input for hydrocarbon generation models of Fig. 12, 13). Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

sites of oil, tar, or natural gas seeps have been identified on the Queen Charlotte Islands (Hamilton and Cameron, 1989). Most of the surface seeps occur in Cretaceous and Tertiary volcanic and sedimentary rocks, many of which are in the Masset Formation. Geological and geochemical studies indicate the seeps are migrated conventional oils, sourced from both Jurassic (Kunga–Maude) and Tertiary sedimentary strata (Fowler et al., 1987; Hamilton and Cameron, 1989). One of the more areally extensive surface oil seeps occurs in Masset volcanic rocks and fractured Cretaceous shale at Lawn Hill on the southeast coast of Graham Island (Fig. 1). Geological and geochemical data from the Lawn Hill seeps indicate that the hydrocarbons were probably sourced from underlying or subjacent Jurassic rocks, with migration into the host rocks in late Neogene time (Snowdon et al., 1988; Hamilton and Cameron, 1989). The Lawn Hill area is part of a high-standing block within the westernmost Queen Charlotte Basin. Cretaceous or Neogene reservoir strata in surrounding or basinward areas may be highly prospective for conventional accumulations of similar oils.

Subsurface hydrocarbon shows were encountered in several petroleum and mineral exploration wells, including gas flows from the Tian Bay well (Hamilton and Cameron, 1989), oil staining in Tertiary volcanic rocks in the Port Louis and Naden wells (Union Oil Company, 1971; Bow Valley Industries, 1984), oil staining in Cretaceous sandstone in the Queen Charlotte well (Royalite Oil, 1949), and oil staining in Neogene sandstone in the Tow Hill and Sockeye B-10 wells (Richfield Oil Corporation, 1958; Shell Canada Ltd., 1968b). The best subsurface hydrocarbon show was encountered in the Sockeye B-10 well, which penetrated 40 m of live-oil-stained Miocene sandstone. Geochemical analysis (gas chromatography–mass spectrometry) of a saturate fraction from the Sockeye oil show indicated the presence of a biomarker compound diagnostic of Jurassic Kunga Group rocks (M. Fowler, pers. comm., 1991). The same biomarker compound is indicated in similar analyses of Kunga outcrop samples from the Queen Charlotte Islands. Other geochemical characteristics of the Sockeye oil show indicate a probable derivation from carbonate rocks. The recognition of a probable Jurassic Kunga source for the Sockeye oil show is an important finding that links the region's principal source and reservoir rocks. The modelled timing of Kunga hydrocarbon generation for the Sockeye area (Fig. 12, 13) suggests the oil may have migrated laterally into the Sockeye structure, a Late Pliocene inversion anticline, (Fig. 6) from an adjacent Mesozoic fault block. Kunga source rocks are overmature beneath the B-10 well location, but may be mature in the surrounding area (Fig. 10, 11). The Sockeye B-10 well also encountered numerous shows of gas-cut mud in coal-bearing zones in Skonun and Cretaceous strata below 3000 m depth (Fig. 10).

Offshore subsurface gas accumulations are inferred to be directly indicated in marine seismic reflection data. Shallow gas in upper Neogene and Quaternary strata have been inferred in many locations beneath Hecate Strait, from acoustic anomalies on high resolution seismic profiles (Barrie, 1988). Indications of possible deep gas accumulations in Neogene strata have been identified on conventional seismic profiles, in several offshore locations (Dietrich, 1995). One such example (star in Fig. 6), illustrates a possible direct hydrocarbon indicator as a subhorizontal, low-frequency reflection at the crest of a fault-bound structure. The direct hydrocarbon indicators occur at a stratigraphic level similar to the Sockeye B-10 well show (Dietrich, 1995).

## **Petroleum assessment**


The West Coast petroleum assessment was undertaken in order to provide quantitative estimates of total oil and gas potential and possible sizes of undiscovered fields in the region. Petroleum assessments of basins or regions are usually based on analyses of a number of exploration plays. The Queen Charlotte assessment involved analysis of three, regional-scale, conceptual plays. Based on considerations of source rock types and hydrocarbon shows, all of the Queen Charlotte plays were considered to have both oil and gas resource components. Appendix A lists all input data used for statistical analysis for each play. Probability distributions of reservoir parameters and marginal probabilities for prospect and play level risks are tabulated.

## **Petroleum plays**

### ***Queen Charlotte Cretaceous oil and gas play***


*Play definition.* This oil and gas play involves all structures and prospects within Cretaceous strata beneath and adjacent to Queen Charlotte Basin. The Cretaceous play area encompasses most of the Queen Charlotte Islands and adjacent shelf areas, extending as far east as the Hecate Basin margin (Fig. 2, 15).

*Geology.* Potential hydrocarbon traps involve Cretaceous sandstone, principally within the basal units of the Queen Charlotte Group, in fault block or anticlinal structures (Table 3). Onshore areas have been mapped where Cretaceous reservoir strata directly overlie Kunga–Maude source rocks (Thompson et al., 1991), an optimum stratigraphic relationship that probably occurs in some fault blocks in the subsurface. The most prospective part of the play area occurs in a southeast-trending fairway from central Graham Island to southwestern Queen Charlotte Sound, an




**Figure 15.** Queen Charlotte Cretaceous oil and gas play area. Major structural elements are illustrated. Area of moderate to high prospectivity is depicted and discussed in text.





**Figure 16.** Estimate of in-place oil potential of the Cretaceous play in Queen Charlotte Basin. Median value of probabilistic assessment is  $392 \times 10^6 \text{m}^3$  of in-place oil distributed in 62 fields.



**Figure 17.** Estimate of in-place gas potential of the Cretaceous play in Queen Charlotte Basin. Median value of probabilistic assessment is  $75 \times 10^9 \text{m}^3$  of in-place gas distributed in 50 fields.

area where the main reservoir facies was deposited (Fig. 15). The play includes areas (such as Graham Island) where potential reservoir strata underlie thick Tertiary volcanic rocks. This play is characterized by relatively small structures and single-reservoir zones.

**Exploration risks.** All of the Queen Charlotte plays are believed to have a high probability of existing risk (i.e. low play risk). However, within each play, risks associated with individual prospects are considered high. A major prospect-level risk in all Queen Charlotte plays is the possible local absence of or inadequate maturation conditions for source rocks (Appendix A, Tables A-1b–A-6b). In addition, a significant risk factor associated with Cretaceous prospects, in particular, is the possible absence of adequate reservoir facies (Appendix A, Tables A-1b, A-2b). The subsurface distribution and reservoir characteristics of Cretaceous strata are expected to be erratic, with little subsurface control currently available. Prospects involving Cretaceous strata may be difficult to map seismically in areas (such as Graham Island) where overlying Masset volcanic rocks are thick and extensive. Assigned exploration risk for Cretaceous plays are 0.11 for oil and 0.09 for gas, with a significant component of the risk associated with the presence of reservoir facies. These exploration risks imply that oil and

gas accumulations will occur in about 11% or 9% (respectively) of all prospects within the play.

**Play potential.** The Cretaceous play is characterized by numerous, small, structurally complex prospects and reservoir zones. This play has an estimated in-place median oil potential of  $392 \times 10^6 \text{m}^3$  (Fig. 16). The mean value of the number of predicted fields is 62. The largest undiscovered field is expected to contain  $96 \times 10^6 \text{m}^3$  of oil (median value). Potential for the Cretaceous gas play is  $75 \times 10^9 \text{m}^3$  (median in-place value) (Fig. 17). The estimate assumes a total field population of 50 (mean value), with the largest undiscovered field having an initial in-place volume of  $21 \times 10^9 \text{m}^3$  of natural gas (see Table 4 for listing of plays, mean and median of potential, and median of the largest pool size).

### **Queen Charlotte Miocene oil and gas play**

**Play definition.** Neogene strata within the Queen Charlotte Basin were assessed in two plays, each play with an oil and gas component, differentiated on the basis of trap type and timing of trap formation (Miocene and Pliocene plays). The Miocene oil and gas play occurs basinwide in an area of about 40 000  $\text{km}^2$  and involves all extensional structure and

**Table 4**  
**Oil and gas potential in Queen Charlotte, Georgia, and Tofino regions (O=oil; G=gas)**

Play name	Expected no. of fields (mean)	Median play potential (in-place) (million m <sup>3</sup> )	Mean play potential (in-place) (million m <sup>3</sup> )	Median of largest field size (in-place) (million m <sup>3</sup> )
Queen Charlotte Region				
Cretaceous oil	62	392	478	96
Cretaceous gas	50	75 435	94 336	20 675
Miocene oil	28	574	668	165
Miocene gas	40	285 710	317 080	71 190
Pliocene oil	13	398	652	233
Pliocene gas	30	321 750	389 710	95 774
Total	103 (O); 120 (G)	1559.8 (O); 733,760 (G)		
<b>Georgia Region (gas only)</b>				
Georgia Pleistocene stratigraphic	92	207	217	10
Georgia Tertiary structural	93	59 329	65 483	9803
Georgia Cretaceous structural	45	118 500	146 780	31 977
Total	230	185,150		
Tofino Region				
Tofino Tertiary structural	41	266 003	266 590	25 982
Total	41	266 003		
Total West Coast basins	103(O); 391(G)	1559.8 (O); 1 228 300 (G)		

stratigraphic traps found within Neogene strata that developed during the transtensional phase of basin development (Fig. 18).

*Geology.* Prospects involve Miocene sandstone and conglomerate and structural or stratigraphic traps. Structural traps include tilted fault blocks, fault-related rollover and drag features and drape anticlines (Table 3). Potential stratigraphic traps are associated with intra-Tertiary unconformities (onlap or subcrop), and updip pinchouts within half-grabens and against fault scarps (Table 3). The Tertiary plays incorporate areas or structures where reservoir strata are in direct stratigraphic or structural contact with Mesozoic rocks, providing favourable conditions for local hydrocarbon charging from Kunga–Maude source rocks.


*Exploration risks.* Along with the significant risk attached to all Queen Charlotte plays concerning inadequate maturation conditions of source rocks, there is a significant risk in the Miocene play associated with the inadequacy of preservation (Appendix A, Tables A-3b, A-4b). Neogene structures are commonly faulted and shallow parts of many northern sub-basins contain high percentages of permeable sandstone. The

Miocene play is assigned an exploration risk of 0.10 for oil and 0.15 for gas, with most of the risk associated with source rock, presence of closure, and preservation (Appendix A, Tables A-3b, A-4b).


*Play potential.* Estimates of the potential for the Miocene oil play show a median in-place volume of  $574 \times 10^6 \text{m}^3$  distributed in 28 fields (mean value) (Fig. 19, Table 4). The largest undiscovered oil field is predicted to contain  $165 \times 10^6 \text{m}^3$  (median value). The Miocene gas play predicts a mean value of 40 fields with a median in-place potential of  $286 \times 10^9 \text{m}^3$  (Fig. 20, Table 4). The largest estimated gas field is  $71 \times 10^9 \text{m}^3$  (median in-place volume).

#### ***Queen Charlotte Pliocene oil and gas play***


*Play definition.* This petroleum play includes all structural traps in Neogene reservoirs formed during contractional deformation associated with late Pliocene transpression. The play area is restricted to the northern half of the Queen Charlotte Basin (Fig. 21).



**Figure 18.** Queen Charlotte Miocene oil and gas play area. Major structural elements are illustrated.



**Figure 19.** Estimate of in-place oil potential of the Miocene play in Queen Charlotte Basin. Median value of probabilistic assessment is  $574 \times 10^6 \text{m}^3$  of in-place oil distributed in 28 fields.



**Figure 20.** Estimate of in-place gas potential of the Miocene play in Queen Charlotte Basin. Median value of probabilistic assessment is  $286 \times 10^9 \text{m}^3$  of in-place gas distributed in 40 fields.

**Geology.** Similarly to the Miocene play, Skonun Formation sandstone and conglomerate constitute the principal reservoir in the Pliocene petroleum play. The Pliocene play is differentiated from the Miocene play on the basis of structural style and timing of trap development. Pliocene structures include large-amplitude folds and faulted anticlines (flower structure) (Rohr and Dietrich, 1992) (Table 3). Many of the Pliocene antiforms are structurally detached from underlying Mesozoic rocks. In comparison to Miocene prospects, the Pliocene prospects are generally larger in area and involve thicker (multi-zone) reservoir sections. Like the Miocene play, the Pliocene play includes many prospects where reservoir strata are in direct structural contact with Mesozoic rocks, which locally include Kunga–Maude source units.


**Exploration risks.** The major prospect-level risk in the Pliocene play is associated with source rock (Appendix A, Tables A-5, A-6b). A source rock timing risk of particular significance to Pliocene prospects is the possibility that Kunga–Maude oil generation and migration occurred prior to late Pliocene trap formation. Another significant assigned risk factor is the inadequacy of seal. Many Pliocene prospects are extensively faulted (hydrocarbon leakage possibility) or involve stratigraphic sections with high percentages of permeable sandstone.

The Pliocene play has an exploration risk of 0.07 for oil and 0.16 for gas, with the most significant risk components associated with adequacy of source rocks and seals. The difference in oil and gas risk factors reflects the greater uncertainty for the presence of oil source rocks.


**Play potential.** The Pliocene play has a median oil potential of  $398 \times 10^6 \text{m}^3$  distributed in 13 fields (in-place volume) (Fig. 22, Table 4). The largest undiscovered field has a predicted volume of  $233 \times 10^6 \text{m}^3$  of oil (median value). The predicted median gas resource within the Pliocene play is  $322 \times 10^9 \text{m}^3$ , in 30 fields (Fig. 23, Table 4). A median value of  $96 \times 10^9 \text{m}^3$  of gas is predicted for the largest field.

### Discussion of assessment results


**Resource potential.** Median estimates of total petroleum potential for the Queen Charlotte Basin region (from all plays) are  $1.56 \times 10^9 \text{m}^3$  ( $9.8 \times 10^9$  bbl) of in-place oil and  $734 \times 10^9 \text{m}^3$  (25.9 TCF) of in-place gas (Table 4; Fig. 24, 25) (Note that the total median estimates for assessment regions are not derived arithmetically by adding together the median hydrocarbon potentials of individual plays. These numbers are summed using statistical techniques). High confidence (90% probability) and speculative




**Figure 21.** Queen Charlotte Pliocene oil and gas play area. Major structural elements are illustrated.




**Figure 22.** Estimate of in-place oil potential of the Pliocene play in Queen Charlotte Basin. Median value of probabilistic assessment is  $398 \times 10^6 \text{m}^3$  of in-place oil distributed in 13 fields.



**Figure 23.** Estimate of in-place gas potential of the Pliocene play in Queen Charlotte Basin. Median value of probabilistic assessment is  $322 \times 10^9 \text{m}^3$  of in-place gas distributed in 30 fields.



**Figure 24.** Estimate of total oil potential for the Queen Charlotte Basin region. Median value of probabilistic assessment is  $1.6 \times 10^9 \text{m}^3$  of in-place oil.



**Figure 25.** Estimate of total gas potential for the Queen Charlotte Basin region. Median value of probabilistic assessment is  $734 \times 10^9 \text{m}^3$  of in-place gas.

(10% probability) estimates of total oil potential are  $657 \times 10^6 \text{m}^3$  and  $3088 \times 10^6 \text{m}^3$  ( $6.3$  and  $19.4 \times 10^9$  bbl), respectively. High confidence and speculative estimates of gas potential are  $338$  and  $1351 \times 10^9 \text{m}^3$  (12 TCF and 48 TCF), respectively (Fig. 24, 25). Individual field-size estimates display similar probability-dependent variations. The wide range of estimates of total potential and field sizes are typical of frontier region assessments and reflect the geological uncertainties in quantifying lightly explored or conceptual exploration plays.

*Resource distributions.* The highest oil potential (volume) occurs in the Miocene play and highest gas potential in the Pliocene play (Table 4). The largest individual oil and gas fields are predicted to occur in the Pliocene play, with median size estimates of  $233 \times 10^6 \text{m}^3$  ( $1466 \times 10^6$  bbl) of in-place oil and  $96 \times 10^9 \text{m}^3$  (3.3 TCF) of in-place gas. Field size rankings for all plays suggest that about 60% of the region's total petroleum resource is expected to occur in the five largest oil and gas fields. This resource distribution indicates a moderately concentrated hydrocarbon habitat, typical of large convergent and transform plate margin basins (Klemme, 1984).

The assessment results indicate that the Neogene Queen Charlotte Basin is expected to contain about 80% of the region's total petroleum resource volume and nine of the ten largest fields, a concentration reflecting the greater abundance and quality of reservoirs within the Neogene Skonun Formation. In terms of general geographic areas within the basin, southern Hecate Strait is considered the most prospective, followed, in order, by the Queen Charlotte Sound, eastern Graham Island, northern Hecate Strait, and Dixon Entrance areas (Fig. 1, 2). The high potential for the southern Hecate Strait area reflects the optimum combination of abundant Neogene reservoir strata, numerous large structures, and presence of Neogene and, at least locally, Jurassic source rocks. Outside the Queen Charlotte Basin margins, the western Graham Island and adjacent shelf areas have some potential in Cretaceous and to a lesser extent, Paleogene prospects. Very little or no petroleum potential is expected in the onshore and inter-island areas of the southern Queen Charlotte Islands and adjacent Pacific continental shelf, because of limited distribution of reservoir strata and the overmature source rocks (Dietrich et al., 1992).

*Assessment results and exploration history.* The exploration risks estimated in the assessment suggest success rates for exploratory drilling in the region should average about one in nine. The absence of discoveries in the 18 wells drilled during early phases of petroleum exploration in the Queen Charlotte Basin region may indicate a higher exploration risk than estimated here. However, many of the previous wells are considered inadequate or only partly diagnostic

exploration tests. Nine of the ten onshore wells were drilled to depths of less than 2000 m, three penetrating entirely volcanic sections. Of the eight offshore wells, three were drilled as purely stratigraphic tests (Osprey, Harlequin and Auklet), three were drilled on the flanks of structures (Murrelet, Sockeye B-10 and E-66), and two tested only parts of fault-segmented anticlines (Coho and Tye). Seismic reflection surveys used to support early exploration programs were characterized by fair or poor data quality compared to current standards. Some wells were undoubtedly drilled in less than optimum locations. Even without these considerations, 18 wells represent evaluation of only a small fraction of the total prospective area within the Queen Charlotte Basin region. Historically, the first significant hydrocarbon discovery in a frontier region is often preceded by many unsuccessful exploration wells. As examples, 36 wells were drilled in the Grand Banks region of offshore Newfoundland before the first field (Hibernia) was discovered (Sinclair et al., 1992), and 16 wells were drilled in the Cook Inlet Basin of southern Alaska before the first field (Swanson River) was discovered (Magoon and Kirschner, 1990). In the case of Cook Inlet Basin, about 15% of subsequent exploration wells led to additional discoveries, an exploration success rate slightly higher than predicted for the Queen Charlotte Basin region.

## GEORGIA ASSESSMENT REGION

### Exploration history and regional studies


The earliest public-domain geophysical survey that encompassed the Georgia Basin region was a 1955 regional aeromagnetic study by the Geological Survey of Canada. In 1959, a gravity survey was carried out by Petcal that encompassed most of the Fraser River Valley west of Abbotsford (Petcal Company Ltd., 1959).

The first extensive seismic reflection survey was acquired by Richfield Oil Corporation in 1959. It extended from Abbotsford to the Strait of Georgia and between the Fraser River and the United States border (Richfield Oil Corporation, 1959a). Richfield also conducted a short seismic program in American waters off Point Roberts (Richfield Oil Corporation, 1959b). BC Hydro Gas Operations was interested in the potential for underground gas storage in the Lower Mainland and conducted a 322 km seismic program in 1977. Conoco conducted a petroleum exploration program in the Lower Mainland in the early 1990s. About 380 km of seismic work was completed, resulting in numerous indications of structural closures and prospects. Both four-way closures and reverse fault traps were identified. Conoco also participated in a survey in the Strait of Georgia, as a result of which more prospects were identified.

South of the border, in Whatcom County, Washington State, seismic reflection work was first conducted in the early 1940s by Chevron and later in 1985 by CGG (Companie Général Géophysique). In 1987, Canadian Hunter performed an extensive 160 km survey in the area, identifying the structure at Birch Bay.

More than 2700 km of marine seismic data were acquired by the petroleum industry in the Strait of Georgia area in the 1960s. Canadian Superior Oil recorded approximately 245 km of gas exploder seismic data in the Strait of Georgia in 1962 (Canadian Superior Oil Ltd., 1962). British American Oil Company conducted a 1150 km-long gas exploder marine seismic survey (British American Oil Company Ltd., 1965). Texaco Exploration Canada

conducted an extensive marine seismic program in the Strait of Georgia in 1968 and 1969 (Texaco Exploration Canada Ltd., 1968, 1969) when about 300 km of marine seismic data were recorded. Two structural highs were delineated: one near Robert's Bank with an area of substantial closure, the other in central Strait of Georgia west of Vancouver. A major normal fault, paralleling the folding and down-dropped to the northeast with a throw of about 1200 m is interpreted from a reflection profile running east of Mayne Island towards Point Roberts (see Fig. 26, cross-section C-C', same fault interpreted just east of Valdes Island). On the east side of Vancouver Island, British Petroleum conducted seismic surveys totalling about 160 km. From these surveys, two wells were drilled on seismically defined structures (British Petroleum Resources Canada Ltd., 1987a, b).



**Figure 26.** Geological cross-sections (derived from seismic and well data) across the Strait of Georgia (C-C'), and running parallel to the mainland coast from Burrard Inlet to Bellingham (D-D'; locations in Fig. 3) (modified from Gordy, 1988 and Mustard and Rouse, 1994) showing main Tertiary and Upper Cretaceous units of Georgia Basin in the area. Main faults are either projected from the surface or known from petroleum exploration seismic lines.



The first petroleum exploration wells were drilled in 1901 in Whatcom County, Washington, and in 1906 in the Fraser River valley in Canada (Johnston, 1923; McFarland, 1983). Since 1901, a total of 118 wells have been drilled for oil and gas within the Bellingham sub-basin (*see* Fig. 3 for well locations and Table 2 for well listings). The Bellingham sub-basin encompasses the Fraser River lowland and Whatcom County (Fig. 2). Twenty-three of these wells were completed in the lower Fraser River valley; the remaining 95 are located in Whatcom County. The vast majority of the wells are shallow (less than 300 m) and were probably originally drilled as water wells.

Most of Georgia Basin, that is, eastern Vancouver Island, Strait of Georgia, and the western portion of the Lower Mainland, is underlain by a Cretaceous sedimentary succession. Seven wells penetrated the Cretaceous succession in the area (Pacific Petroleum Ltd., 1959; Richfield Oil Corporation, 1962b, 1963; British Petroleum Resources Canada Ltd., 1987a, b; Hurst, 1991). The three wells drilled on the mainland penetrated Cretaceous strata underlying a thick Tertiary cover. On Vancouver Island and Saturna Island, four wells were collared in Cretaceous strata (Fig. 3, Table 2).

## **Regional geology**

### ***Geological setting and tectonic evolution***

There has been much debate concerning the tectonic setting of Georgia Basin. Mustard (1994) speculated that the Upper Cretaceous–Tertiary Georgia Basin is a hybrid foreland and strike-slip basin that developed in response to (post-Wrangellian) terrane accretion and subduction along the Vancouver Island continental margin. The Cretaceous basin developed within a foreland tectonic setting, with significant portions of basin-fill likely derived from uplift areas to the east and southeast (Mustard, 1991, 1994). Southern parts of the basin subsequently evolved in (Tertiary) strike-slip or pull-apart tectonic regimes, with basin-fill sediments derived from multiple source areas (Johnson, 1982; Mustard and Rouse, 1994). England and Bustin (1998) designate Georgia Basin as a broad-ridged forearc basin analogous to the Great Valley forearc basin of California (Dickinson, 1976; Dickinson and Seely, 1979).

Georgia Basin occupies 14 000 km<sup>2</sup> in southwestern British Columbia and northwestern Washington State (Fig. 2, 3). The basin includes the Lower Fraser River area and Whatcom County as well as the Strait of Georgia and eastern Vancouver Island. There are four sub-basins within Georgia Basin. Nanaimo sub-basin encompasses the southeastern coast of Vancouver Island and adjacent Strait of Georgia as well as the Gulf Islands. Comox sub-basin is located farther north along the east-central coast of

Vancouver Island and adjacent Strait of Georgia. Suquash sub-basin is located in the vicinity of Port Hardy on the northeastern shore of Vancouver Island. The Bellingham or Whatcom sub-basin includes the Fraser Delta and northwestern Washington. Upper Cretaceous to Recent sedimentary strata comprise the fill of the Georgia Basin (Fig. 8, 26).

The Upper Cretaceous and (?)lowest Tertiary Nanaimo clastic deposits in Georgia Basin represent an overlap assemblage that was eroded from rocks uplifted during terrane accretion. Monger (1991a) postulated the basin developed in a foreland setting with associated west-verging thrust faults, crustal thickening, and regional uplift of the Coast Mountains. England and Calon (1991) suggested that Nanaimo sedimentation occurred in a forearc setting related to Late Cretaceous plutonism in the Coast Mountains. Mustard (1994) interpreted a foreland basin model for the Cretaceous Georgia Basin based on recognition of Late Cretaceous thrust systems in the southern Coast Belt and northwest Cascades and syntectonic Nanaimo Group sedimentation. Recognition of multiple source areas for Cretaceous sedimentary rocks, including areas to the west of the basin, suggest a foreland rather than a forearc basin setting (Mustard, 1994). England and Bustin (1998) on the other hand, interpreted Georgia Basin as a forearc basin that developed between a Late Cretaceous arc in the eastern Coast Mountains and a trench or transform fault thought to be present at the time, west of Vancouver Island.

Seismic refraction profiles in the Strait of Georgia interpreted by White and Clowes (1984), indicate thickening of Cretaceous sedimentary rocks to the west and no major discontinuity separating the Coast Plutonic Complex from the Insular Belt. These profiles also suggest that the Coast Range intrusive rocks extend beneath the Cretaceous sedimentary layer in the Strait of Georgia.

### ***Stratigraphy and structure***

Basement rocks in Georgia Basin consist of complexly deformed assemblages of Devonian to Lower Cretaceous volcanic, plutonic and sedimentary rocks, occurring in parts of several terranes within the Insular, Coast and, in Washington, Cascade tectonic belts (Monger and Journeay, 1994; Mustard, 1994). On Vancouver Island, the Wrangellian Karmutsen Formation is up to 6000 m thick and comprises pillow basalts, breccias and massive flows (Muller, 1977). The upper part of the Karmutsen Formation is locally interbedded with carbonate rocks of Late Triassic age (Fig. 8). The Karmutsen Formation is conformably overlain by the Quatsino Formation, which consists of 300 m of shallow-water and fossiliferous carbonate rocks (Carlisle and Susuki, 1974). A gradational contact separates the Quatsino Formation from overlying dark siliceous and

carbonaceous shales of the Parson Bay Formation. Calcareous sandstone and fine-grained sandstone are lesser components of the 300 m thick Parson Bay Formation. This formation is in turn gradationally overlain by the Jurassic Harbledown Formation, a unit of thinly bedded shale, siltstone and tuff about 300 m thick. The Harbledown Formation grades westward into Lower Jurassic Bonanza Group rocks, a calc-alkaline, dominantly pyroclastic volcanic assemblage with interbedded shale and siltstone. The Bonanza Group varies in thickness up to 2000 m. Comagmatic with the Bonanza Group are the Early and Middle Jurassic Island Intrusions of Vancouver Island (Muller, 1977; Isachsen et al., 1985). These intrusions are unconformably overlain by Upper Cretaceous sedimentary rocks of the Nanaimo Group. A complex and composite boundary separates basement modified by tectonic, magmatic and metamorphic processes from relatively unaffected basin fill. Structures within the basement rocks include broad, northwest-aligned antiforms and northwest- and northeast-aligned thrust and strike-slip faults.

Eastern Georgia Basin strata overlie mid-Jurassic to mid-Cretaceous intrusive complexes of the Coast Belt (Monger and Journeay, 1994). Lower Cretaceous Gambier Group volcanogenic sedimentary and volcanoclastic rocks unconformably overlie the intrusive 'basement' (Fig. 8) (Lynch, 1991, 1992; Monger and Journeay, 1994). Gambier Group strata consist of massively bedded intermediate to felsic volcanoclastic rocks, local flows and interbedded, fine grained tuff and argillite (Monger, 1991b). Recent palynological studies on well cuttings from the Richfield Sunnyside and Point Roberts wells indicate an Albian to Cenomanian sedimentary succession below the Nanaimo Group (Mustard and Rouse, 1991). These rocks are equivalent in age to Gambier Group rocks exposed farther north (Mustard and Rouse, 1994). In the western Georgia Basin region, the Early Jurassic Bonanza Group of volcanic rocks and its laterally equivalent shale unit (Harbledown Formation), are overlain unconformably by the Upper Cretaceous Nanaimo Group (Fig. 8). In eastern Georgia Basin, the Nanaimo Group unconformably overlies the volcanogenic Gambier succession. The Nanaimo Group consists of up to 4000 m of siliciclastic strata (Mustard, 1991). The succession consists of alternating coarse- and fine-grained sedimentary rocks and is characterized by an overall upward progression from nonmarine to marine deposits (Mustard, 1991, 1994). Coal beds are common in the lower alluvial/fluvial facies. The largest portion of the succession is represented by deep-marine fan deposits. Fluviodeltaic and turbidite sandstone and conglomerate are potential petroleum reservoirs (England, 1991).

Paleogene strata in the Georgia Basin are represented by the Huntingdon Formation in Canada and the laterally equivalent Chuckanut Formation in Washington State (Vance, 1975; Johnson, 1984; England and Hiscott, 1992)

(Fig. 8). The succession consists of nonmarine conglomerate and sandstone, with minor amounts of mudstone and coal. Paleogene strata are up to 2500 m thick and 6000 m thick in Canadian and American parts of the basin, respectively (Johnson, 1984, 1991; Mustard and Rouse, 1994). In the subsurface, the Huntingdon Formation is interpreted as a thick fluvial sequence with laterally migrating meandering channels in a sand-dominated floodplain (Mustard and Rouse, 1994). There is no evidence of a marine component. In the Vancouver area, Georgia Basin Paleogene and older sedimentary rocks are locally intruded by Oligocene igneous dykes, sills and flows (Hamilton and Dostal, 1994; Mustard and Rouse, 1994).

The Miocene–Pliocene Boundary Bay Formation (Fig. 8) underlies the Fraser Delta area of the Georgia Basin. The Boundary Bay Formation is up to 1200 m thick and consists of interbedded fluvial sandstone and mudstone, and lesser amounts of conglomerate and coal (Mustard and Rouse, 1994). Channel and overbank, floodplain and crevasse splay deposits are recognized (Mustard and Rouse, 1994).

Pleistocene glacial and interglacial sediments blanket the Fraser Delta and Whatcom County areas. In the Bellingham sub-basin, Pleistocene sediments are up to 700 m thick (Hamilton and Ricketts, 1994). The succession consists of glacial till, stratified fluvio-glacial sand and gravel, and peat accumulations (Clague, 1994). The Holocene Fraser River delta beneath the Strait of Georgia consists of fluviodeltaic sand and mud, in places exceeding 200 m in thickness (Johnston, 1921; Clague et al., 1983; Luternauer et al., 1994).

The principal structures deforming Cretaceous and Paleogene strata in Georgia Basin are northwest-aligned folds and thrust faults. The contractional structures are part of a regional Eocene fold and thrust belt that developed in association with terrane accretion and crustal underplating beneath western Vancouver Island (England and Calon, 1991; England et al., 1997). In parts of the Georgia Basin, strata are also disrupted by Late Eocene and younger northwest- and northeast-trending normal and strike-slip faults (Fig. 26) (Johnson, 1984; Monger and Journeay, 1994; Mustard and Rouse, 1994).

## **Petroleum geology**

### ***Reservoirs***

#### ***Mesozoic***

The oldest strata inferred to have significant reservoir potential in the Georgia Basin are clastic rocks of the Upper Cretaceous Nanaimo Group. These outcrop along the east coast of Vancouver Island on the Gulf Islands in the southern

Strait of Georgia. Isolated outcrops occur in the Suquash sub-basin on northern Vancouver Island, the Alberni valley, and in the Lang Bay area directly south of Powell River on the mainland (Fig. 1, 2, 4). Additional isolated outliers of Nanaimo Group are located on Texada Island, Lasqueti Island and on Blue Mountain near Vancouver (Fig. 1, 2, 4). The strata on Blue Mountain were previously considered to be Tertiary (Mustard and Rouse, 1991). Rocks formerly mapped as Tertiary Burrard Formation at the margins and north of Burrard Inlet in the Vancouver area are now considered to be correlative with Nanaimo Group sedimentary rocks (Mustard and Rouse, 1994). The Nanaimo Group subcrops beneath Tertiary sedimentary rocks in the Strait of Georgia, western Fraser Valley, and Birch Bay area of Washington State (Fig. 3, 26).

The Nanaimo Group has been subdivided into eleven mappable units (not shown on Fig. 8). The formations (in upward stratigraphic order) are Comox, Haslam, Extension, Pender, Protection, Cedar District, De Courcy, Northumberland, Geoffrey, Spray, and Gabriola. These formations encompass variably alternating sequences of shallow- and deep-marine, marginal-marine and nonmarine clastic deposits. Sandstone, siltstone, shale, conglomerate, and coal seams are present throughout the succession, which varies in thickness between 300 and 4000 m. Neritic to bathyal marine depositional environments are represented by outer-shelf deep-marine turbidites, submarine fans, and slope facies. Shallow-marine and littoral facies associations represent marginal-marine deposition. Terrestrial deposition includes fluvial-alluvial deltaic and lagoonal deposits. Coal measures are important features of the Extension, Pender, and Protection formations.

Sandstone and conglomerate are potential reservoir units within the Nanaimo Group. Coarse clastics constitute about 50 per cent of the Nanaimo Group succession, of which five to eight per cent are considered reservoir quality. Six wells penetrate the Nanaimo Group in Georgia Basin. Most coarse clastic beds are tight in these wells. There are, however, minor thin sandstone or conglomerate beds with slight to moderate porosity. An 8 per cent porosity was used as a cutoff for marking reservoir-quality facies. Decomposition of feldspar and volcanic grains within the coarse-grained arkosic and lithic sediments resulted in an increased clay content and a subsequent occlusion of pores and reduction in porosity and permeability. Sediments derived from the Coast Plutonic Belt to the east are feldspar-rich. Sand and conglomerate derived from Vancouver Island to the west also produce reservoirs of poor to intermediate quality as a result of poor sorting and feldspar content. England (1990) made porosity measurements on surface outcrops of the Nanaimo Group on eastern Vancouver Island. His measurements varied from 0 to 11.5 per cent with 8 out of 78 samples exhibiting greater than 8 per cent porosity. Porosities are generally low in Nanaimo Group surface

samples (below 5%; Yorath, 1987). Secondary fracture porosity has not been reported with respect to Nanaimo Group rocks. Permeabilities are very low in these rocks in the subsurface (0.01 to 0.06 md). Total reservoir thickness within the 4000 m Cretaceous succession varies from 10 to 200 m, averaging 60 m.

### *Tertiary*

Tertiary rocks underlie the western Fraser Valley, northwestern Washington, and southern Georgia Strait (*see* Fig. 26). Miocene Boundary Bay clastic rocks unconformably overlie Paleogene Huntingdon and Chuckanut formation strata in the Fraser Valley and northwestern Washington State (Fig. 8). These rocks are mainly exposed in scattered outcrops along the lower Fraser River valley and east and northeast of Bellingham in northwestern Washington. Tertiary rocks also outcrop on Tumbo, Sucia, Lummi, and Lasqueti islands in the Strait of Georgia. The succession is estimated to be about 2500 m thick in the Fraser Lowland and thickens to 6000 m near Bellingham. The thickness of the prospective succession (interval containing reservoir-quality material), however, seems to be restricted to the upper 2000 m. Sandstones below 2000 m depth tend to have lower porosities and lack permeability.

Porous sandstone units are generally thin, the majority varying in thickness between 0.6 and 5 m. There are local 10 m thick reservoir-quality sandstone units and very rare 30 m porous sandstone units. Potential reservoir sands represent about seven per cent of the total succession. Within individual structures, a succession of stacked reservoirs is present. Gordy (1988) indicated that prospective sandstones in southwestern British Columbia vary in porosity from 8 to 34 per cent, with an average of 15 per cent. In Washington State, porous sandstone has an average porosity of 12 to 15 per cent. There is evidence of secondary fracture porosity due to significant water and gas flows below 2000 m depth, where primary matrix porosity is negligible. Significant permeability measurements have been obtained within the Tertiary succession. In the AHSL Birch Bay No. 1 well, permeability is reduced below 1370 m. However, good quality reservoir rock is present above this depth (>100 md/ft.) (Hurst, 1991).

The Paleogene Chuckanut and equivalent Huntingdon formations are nonmarine fluvial and alluvial-type clastic deposits (Johnson, 1984). Medium- to coarse-grained arkosic and lithic sandstone and conglomerate are the principal rock types, with lesser shale, mudstone, siltstone, and lignite (Richfield Oil Corporation, 1962a). Potential reservoir facies include coarse clastic deposits preserved in braided stream channels and alluvial fans. Feldspars and lithic fragments in Paleogene sandstone are less degraded,

contain little silica cement, and show less compaction compared to Nanaimo Group sedimentary rocks. Reservoir-quality rocks are more likely to occur in Tertiary sedimentary rocks than in the Nanaimo Group.

Core analysis of Miocene sedimentary rocks of southwestern British Columbia reveal that there is a general fining-upward character of quartz and feldspar grains in a cement matrix. Microporosity within the cements (rather than intergranular porosity) characterizes these rocks. The percentage of clay in Miocene sedimentary rocks generally decreases with depth, resulting in some improvement in reservoir quality in deeper parts of the succession. Porosities vary from 8 to 21% and permeabilities increase with enlargement of pore size. Reservoir quality is defined as poor to good in Miocene rocks.

### *Pleistocene*

Biogenic gas has been produced from the Pleistocene sands and gravels on the Fraser River delta of British Columbia and the Bellingham sub-basin in northwestern Washington State. Highly porous (15-20% estimate) lenticular gravels and sands within the tills constitute potential reservoir facies within the succession. Gas is likely trapped stratigraphically against impermeable clays. Estimated fraction of reservoir-grade gravels and sands compared to total thickness of Pleistocene deposits is 30%.

### *Seals*

In general, adequate lateral and top seals for Cretaceous and Tertiary reservoir strata are provided by numerous interbedded and overlying shale units in Georgia Basin. Structure-related seals may be present where sandstone and shale units are in fault contact. Seal potential may be reduced for Paleogene strata as a result of an overall high sand content for the units (England, 1991). As mentioned previously, potential gas in unconsolidated gravels and sands in Pleistocene glacial material on the mainland may be trapped laterally and/or vertically by clay-rich boulder clays and tills. However, a greater prospect-level risk for seal was assigned to the Pleistocene interval because of nondeposition of sealing material in some prospective areas.

### *Traps*

Potential petroleum traps found in Upper Cretaceous Nanaimo Group rocks are simple compressional anticlines, faulted anticlines (e.g. Birch Bay; Hurst, 1991), normal faults, drag closures, sandstone pinch-outs, and unconformity truncations (Table 3). The structural traps were formed during mid- to Late Eocene time. Seismic

sections and/or seismic structural contour maps reveal a minimum of approximately 60 identifiable structures or prospects. The largest closure area is 50 km<sup>2</sup>.

Traps within the Tertiary succession of Georgia Basin are simple compressional folds, reverse faults, normal faults, sandstone pinch-outs, and unconformity truncations, singly or in combination (Table 3). An older trend of northwest-aligned compressional folds and minor faults of probable Late Eocene age are cut by a younger set of northeast- to east-trending high-angle normal and reverse faults with apparent dip-slip offsets. These younger structures are interpreted as having formed in mid-Tertiary and older strata (Mustard and Rouse, 1994). The Sumas Mountain and Vedder faults bound a graben structure filled by Quaternary and Recent sediments (Fig. 26). At least 60 structures or prospects in Tertiary strata have been identified from seismic mapping in the area. The identifiable prospects are either four-way closures or traps formed against reverse faults. The largest closure observed covers an area of 50 km<sup>2</sup>. Structures occur at various levels in the Tertiary sequence.

Thin, lenticular, sand and gravel body pinch-outs against glacial till constitute the stratigraphic trap type in Pleistocene deposits of the Georgia Basin (Table 3). Numerous traps are preserved as a result of the complex stratification of glaciofluvial sands within impermeable boulder clays and tills.

### *Source rocks*

Bustin and England (1991) examined numerous samples of Nanaimo Group material for hydrocarbon potential. Generally, the strata are characterized by low TOC content (< 1%). However, elevated TOC contents do occur in some samples from the Comox, Extension, Pender, and Protection formations as a result of the presence of coal seams and coaly material (Fig. 8; Bustin and England, 1991). Humic terrestrial deposits of coal and carbonaceous mudstone represent excellent gas source material. Therefore, coal measures present in these formations have good potential for gas and limited capacity for producing liquid hydrocarbons (England et al., 1989). However, significant gas-generating source material (> 5% TOC), is restricted to the northwestern part of the basin (England, 1991). These rocks have both biogenic and thermogenic gas potential. Bustin and England (1991) also measured hydrogen and oxygen indices in Nanaimo Group rocks. Moderate to low average hydrogen and oxygen indices indicate that the strata are mainly composed of Type II (oil and gas prone) and Type III (gas prone) organic matter. The low average TOC content and moderate to low Hydrogen Index values indicate poor potential for generation of liquid petroleum and greater potential for gas (Bustin and England, 1991). Liquid hydrocarbons derived from these source rocks are more

likely to be condensate rather than oil (England et al., 1989). There may be better source rock potential, possibly oil-generating source material, in the central undrilled area of the basin (Georgia Strait) as a result of euxinic conditions during deposition of marine shale (England, 1991). It is also possible that source rock exists in Triassic basement of Wrangellia. According to England (1991), gas and condensate seeps are present on the western margin of Georgia Basin.

Source rocks have been identified in the Tertiary sedimentary succession. Coal seams within Eocene sedimentary rocks in northwest Washington and organic-rich shale found throughout the basin are important source material for light hydrocarbons (gas and condensate). Good oil source rocks have not been identified as yet in Tertiary Georgia Basin. Even though abundant liquid-hydrocarbon-prone Type IIB kerogens have been distinguished in Eocene Chuckanut Formation rocks, oil generation has probably been retarded by low maturity resulting from the shallow burial of Eocene strata. The Eocene Chuckanut Formation rocks also have TOC contents up to 6% (Hurst, 1991). In Birch Bay No. 1 well (Fig. 3; Table 2), high concentrations of Type IIB kerogens are present with numerous intervals of elevated TOC content (Hurst, 1991). Rocks below 1830 m depth in the Birch Bay well contain humic Type III kerogen (gas prone) with a low TOC content. Organic shale and coal seams in Miocene sedimentary rocks retain Type III kerogens with TOC varying from 1 to 17%. Gas seeps have been reported throughout the lower Fraser Valley and in northwestern Washington (Johnston, 1923; McFarland, 1983; Hurst, 1991; Lingley and von der Dick, 1991).

It has been determined that reported oil shows, especially within eastern Georgia Basin, represent industrial contaminants rather than natural occurrences (Johnston, 1923; Moen, 1969; McFarland, 1983). Geochemical data indicate the area is not likely oil-prone (Yorath, 1987; Gordy, 1988; England, 1991; Bustin and England, 1991).

In summary, Georgia Basin contains good gas-source rocks in both Cretaceous and Tertiary sections, but no mature oil-source rocks have yet been identified. Thus, Georgia Basin is currently considered to be exclusively a gas/condensate hydrocarbon province.

### ***Source-rock maturation***


Generally, the Nanaimo Group sedimentary succession is mature with respect to hydrocarbon generation (England, 1990). Nanaimo Group rocks are overmature (Ro as high as 4.6%) adjacent to Tertiary plutons or in areas of overthrusting (England, 1991; Mustard, 1994). In subthrust positions in the Cowichan Fold and Thrust Belt, vitrinite reflectance values are well into the gas window (Ro > 1.3%)

(England, 1991). A depth/maturity plot for the Yellow Point well drilled entirely in Nanaimo Group strata (Fig. 27; see Fig. 3 and Table 2 for well location) reveals all the succession is in the gas generation zone (0.7 to 1.3% Ro). These rocks also occupy the oil window from surface to total depth, but lack of a suitable source rock for oil precludes oil generation. Bustin and England (1991) reported vitrinite reflectance values of Nanaimo Group rocks varying from 0.4 to 5% Ro (immature to overmature) from their sampling program on Vancouver Island. The sampling revealed that throughout the Georgia Basin proper, most of the strata have maturation levels within the oil window while in the Suquash sub-basin, the rocks are mainly immature. According to the predicted model, the Cretaceous Nanaimo and Gambier-equivalent sedimentary succession at Point Roberts is mature with respect to hydrocarbon generation where it is buried under 3000 m of Tertiary and Quaternary rocks (Fig. 28, see Fig. 3 and Table 2 for well location). Maturation levels offshore in Georgia Strait are unknown, but they are expected to vary from mature to overmature. Possible oil generation may occur in the more deeply buried marine shale found in the Nanaimo Group in Georgia Strait.

According to numerous studies of the Tertiary succession of Georgia Basin, levels of organic maturation range from immature to marginally mature, regardless of depth (Bustin, 1990; England, 1991; Hurst, 1991; Lingley and von der Dick, 1991; Mustard and Rouse, 1991). Vitrinite reflectance measurements from the Point Roberts well, for example, vary from 0.4 to 0.6% Ro from surface to 4400 m total depth. There is effectively no increase of reflectance with depth of burial (Bustin, 1990; Fig. 28). The ambiguity of this data prevents rigorous interpretation. The only conclusion is that Tertiary strata are immature to marginally mature with respect to hydrocarbon generation. Thermal alteration index values obtained from the Richfield Sunnyside well near by indicate marginally mature strata (Mustard and Rouse, 1991). Surface and subsurface samples from the Point Roberts area also reveal the TAI is in the marginal mature range. England (1991) demonstrated that surface samples of Paleogene strata at southeastern Vancouver Island and Gulf Islands are mainly immature to early mature (0.29 to 0.66% Ro). Thus, all available maturation data indicate Tertiary strata in Georgia Basin are probably too immature for the generation of large volumes of thermogenic gas. In several wells in the Bellingham sub-basin in Washington State, pyrolysis and vitrinite reflectance data indicate Tertiary strata are marginally mature, with little apparent increase in maturation levels to depths of up to 2000 m (Hurst, 1991; Lingley and von der Dick, 1991).


### ***Timing of hydrocarbon generation***

Major compressional structures formed during mid to late Eocene time. Most of the basin fill was deposited during Late



**Figure 27.** Maturation–depth profile of strata in the Yellow Point d-84-C location, Georgia Basin (see Fig. 3 and Table 2 for well location). Single surface vitrinite reflectance value obtained from England and Calon (1991). Measured subsurface TMAX values are from Bustin and England (1991). Model-predicted maturation profile based on illustrated stratigraphy and heat-flow model where heat flow is reduced from 40 mW/m<sup>2</sup> to 35 mW/m<sup>2</sup> at 50 Ma. Subduction of oceanic crust commencing in mid-Eocene beneath Vancouver Island produces a large overlying wedge of cool crust (Lewis et al., 1992). Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

Cretaceous to early Eocene. England (1991) suggested the timing of hydrocarbon generation with respect to structure formation due to compressional deformation poses a potential risk for the trapping of hydrocarbons in Georgia Basin. England's basin model suggests that the initial gas charge produced by normal burial metamorphism with prevailing low geothermal gradients predates trap formation. However, trap formation is inferred to postdate hydrocarbon generation in other hydrocarbon-bearing settings such as the Foothills of the Canadian Cordillera. This study considers that this timing problem does not significantly detract from the potential in the Georgia Basin, although it does increase the play risk. Secondary hydrocarbon generation can be achieved by continuous burial of Cretaceous and Paleogene strata by thick accumulations of Neogene material in eastern Georgia Basin and by tectonic burial in subthrust positions in



**Figure 28.** Maturation–depth profile of strata in the Richfield Pure Point Roberts 6-3-5 location, Georgia Basin (see Fig. 3 and Table 2 for well location). Measured subsurface vitrinite reflectance values obtained from Bustin (1990). Model-predicted maturation profile based on illustrated stratigraphy and constant heat flow of 35 mW/m<sup>2</sup> (heat flow derived from Lewis et al., 1992). Note ambiguous results for measured vitrinite reflectance. No apparent increase in vitrinite reflectance as a function of depth, expected with normal burial metamorphism, occurs in the subsurface. The only conclusion possible is that all strata are immature to mature with respect to hydrocarbon generation. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

the Cowichan Fold and Thrust Belt (England, 1991; Mustard, 1994). Traps resulting from extensional deformation, such as normal fault traps, postdate the major hydrocarbon charge brought on by normal burial metamorphism. Deep-seated faults may provide pathways for hydrocarbon migration upsection or updip into potential traps.

In eastern Georgia Basin, modelled burial history curves indicate hydrocarbon generation occurring in the interval from late Miocene to Recent time in Gambier-‘equivalent’ sedimentary rocks. This postdates mid-Eocene folding and faulting (Point Roberts well; Fig. 29). This model suggests that hydrocarbons will encounter numerous compressional and extensional traps. Timing is not inferred to be a problem in this area. This model assumed a constant heat flow

throughout the burial history ( $35 \text{ mW/m}^2$ ). Published heat-flow data for the southern Canadian Cordillera gives an average of  $35 \text{ mW/m}^2$  for the Georgia Basin region (Lewis et al., 1992).


Modelling of the Nanaimo Group sedimentary succession at the Yellow Point well on southeastern Vancouver Island reveals that hydrocarbon generation commences in latest Cretaceous time and continues to Recent (Fig. 30). The mid-Eocene deformation episode provides opportunities for trapping generated hydrocarbons. Timing of trap formation with respect to hydrocarbon generation is not inferred to be a significant problem in this part of the Georgia Basin. This model assumes a reduction in heat flow at 50 Ma (mid-Eocene) ( $40 \text{ mW/m}^2$  to  $35 \text{ mW/m}^2$ ) reflecting the commencement of subduction of oceanic crust beneath Wrangellia, which in turn cools the overlying crustal wedge.

### Hydrocarbon shows


Numerous gas shows and limited gas production have been described in drillers' logs, well history reports, geological reports, and published papers (Key Oil and Gas (1955) Ltd., 1958; Pacific Petroleum Ltd., 1959; Moen, 1969; British

Petroleum Resources Canada, 1987a; Gordy, 1988; Hurst, 1991). Gas production from glacial sands and gravels at shallow depths has been reported in the Bellingham sub-basin. Six wells near Birch Bay are all former domestic gas producers (Fig. 3). Production rates and pressures were reported for five of these wells ( $21\,237$  to  $141\,585 \text{ m}^3/\text{day}$  at  $186$  to  $482 \text{ kPa}$ ) (Glover, 1935; Moen, 1969). Gas analyses indicate major constituents are methane and nitrogen suggesting a dry biogenic gas source, probably coal seams (Moen, 1969). The gas derived from underlying coal seams migrated upward and accumulated in the porous glacial sands and gravels in stratigraphic traps. Gas shows have also been reported from Pleistocene sediments encountered in eight wells in Bellingham sub-basin.

Numerous gas shows have been reported from Tertiary strata in Bellingham sub-basin. More than 98 hydrocarbon shows were encountered in 84 wells that penetrated the Tertiary succession. Many old drilling reports refer to "several" or "a few" hydrocarbon shows. Miocene Boundary Bay sedimentary rocks have more than 20 hydrocarbon shows of which 17 were coalbed methane 'kicks' encountered in one well. Seventy-eight shows occur in Eocene-Lower Oligocene Huntingdon and Chuckanut formations. Most shows are gas, but minor oil occurrences



**Figure 29.** Subsidence and hydrocarbon generation model for the Point Roberts area, Tertiary Georgia Basin. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995). Hydrocarbon generation models based on standard (BasinMod) kinetic parameters for organic matter types (Types II–III in Boundary Bay and Huntingdon strata and Type III in Nanaimo and Gambier-equivalent strata). See text for discussion.



**Figure 30.** Subsidence and hydrocarbon generation model for Yellow Point area, Cretaceous Georgia Basin. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995). Hydrocarbon generation models based on standard (BasinMod) kinetic parameters for organic matter types (Types II–III in Protection, Pender, Extension and Comox strata and Type III in De Courcy, Cedar District and Haslam strata). See text for discussion.

have been reported. It has been suggested, however, that these oil occurrences are probably contaminants or refined oil spills rather than naturally occurring accumulations (Moen, 1969; Gordy, 1988).

In the Cretaceous interval, six gas shows from six wells have been reported; most are methane gas kicks from coal seams. Hurst (1991) reported that American Hunter's well at Birch Bay encountered gas shows below 6000 feet (1830 m) depth in Cretaceous rocks. Analysis of gas seeps from old well casings revealed a thermogenic origin with some biogenic mixing (Hurst, 1991). The presence of numerous thermogenic gas shows both in Cretaceous and Tertiary sediments indicates potential for commercial quantities of gas in Georgia Basin.

## **Petroleum assessment**

Two conceptual and one immature play are recognized in Georgia Basin. Assessment computations were performed for natural gas only in the Georgia Basin area. Oil assessments were not done for these plays because geochemical data suggest the area is not likely oil-prone.

## **Petroleum plays**

### ***Georgia Cretaceous structural gas play***

*Play definition.* The Georgia Cretaceous gas play includes structural and structural–stratigraphic traps in Cretaceous strata in Georgia Basin (Fig. 2, 3, 31). The play area includes the western Fraser Delta, Strait of Georgia, southern Gulf Islands and eastern coast of Vancouver Island.

*Geology.* Prospects involve single or stacked Cretaceous sandstone units in anticlines or thrust fault blocks. Stratigraphic trap components include pinch-outs and unconformity truncations.

*Exploration risks.* The exploration risk for the conceptual Georgia Cretaceous play is estimated at 0.10, with most of the risk associated with adequacy of reservoir facies. Pacht (1984) identified lithic arkose and volcanic lithic arenite as the predominant sandstone units on eastern Vancouver Island. Porosity in these rocks is probably low because diagenetic activity causes breakdown of abundant feldspars and volcanic clasts into clay minerals that plug pores in sandstone. Although structure, source rock, and maturity characteristics are favourable for hydrocarbon formation and accumulation, the lack of porosity development will establish a higher risk for exploration success in the play (Appendix A, Table A-7b).

*Play potential.* The total median play potential is  $118.5 \times 10^9 \text{m}^3$  of gas (Fig. 32, Table 4). The estimated median of the largest field size is  $32 \times 10^9 \text{m}^3$ . The number of fields expected in the play is 45. Compared to the other plays in the Georgia Basin, this play possesses the greatest potential. However, the Cretaceous play carries a higher exploration risk for adequate conditions for hydrocarbon accumulation.

### ***Georgia Tertiary structural gas play***

*Play definition.* The Georgia Tertiary gas play includes structural and structural–stratigraphic traps in Tertiary strata in Georgia Basin. The play area encompasses the onshore Lower Mainland in the Vancouver area and offshore Strait of Georgia (Fig. 3, 33).

*Geology.* Prospects include Tertiary sandstones (commonly in stacked successions) in anticlines, thrust and normal fault traps, and stratigraphic pinch-outs and unconformity truncations.

*Exploration risks.* The Georgia Tertiary play was assigned an exploration risk of 0.56, with most of the risk associated with the presence of closure and the adequacy of seal (Appendix A, Table A-8b).

*Play potential.* The estimated median resource potential for the play is  $59 \times 10^9 \text{m}^3$  of gas in 93 fields (Fig. 34, Table 4). The median estimate of the largest field size is  $9.8 \times 10^9 \text{m}^3$  of gas.


### ***Georgia Pleistocene stratigraphic gas play***

*Play definition.* The Georgia Pleistocene stratigraphic play encompasses all stratigraphically trapped gas within Quaternary unconsolidated sediments in Bellingham sub-basin. The play area encompasses the Fraser delta and Fraser lowlands (Fig. 2, 3, 35).


*Geology.* The play consists of stratigraphic traps involving lenticular sands and gravels encased in impermeable mudstone. These traps reflect the complex stratification of glaciofluvial sands within impermeable boulder clays and tills. Coal seams found in underlying Tertiary sedimentary rocks are potential sources for the dry, biogenically generated gas for the play.

*Exploration risks.* The exploration risk for the Georgia Pleistocene play was estimated at 0.40, with most of the risk associated with the adequacy of seal (reflecting the shallow burial of reservoir units and erratic distribution of mudstone and clay) (Appendix A, Table A-9b).





**Figure 31.** Georgia Cretaceous structural gas play area. Major structural elements are illustrated.



**Figure 32.** Estimate of in-place gas potential of the Cretaceous structural play in Georgia Basin. Median value of probabilistic assessment is  $118.5 \times 10^9 \text{ m}^3$  of in-place gas distributed in 45 fields.

**Play potential.** The median potential gas resource of the play is  $207 \times 10^6 \text{ m}^3$  of gas in 92 expected fields (mean value) (Fig. 36, Table 4). The largest field in the play is predicted (at median value) to contain  $10.4 \times 10^6 \text{ m}^3$  of in-place gas.

### Discussion of assessment results

**Resource potential.** Total gas potential for all three plays in Georgia Basin is  $185 \times 10^9 \text{ m}^3$  (6.5 TCF) (in-place volume) distributed in 230 predicted fields (Table 4; Fig. 37). The high confidence (90% probability) and speculative (10% probability) estimates of total gas potential in Georgia Basin are  $43.3 \times 10^9 \text{ m}^3$  and  $419.9 \times 10^9 \text{ m}^3$  (1.5 and 15 TCF), respectively. Total estimated gas resource for the Georgia Basin constitutes about one quarter of the predicted potential for the Queen Charlotte Basin area. Although the Georgia Basin assessment region is predicted to contain a larger number of gas fields than the Queen Charlotte Basin assessment region (Table 4), the sizes of the fields are predicted to be smaller in the former area.


**Resource distributions.** The greatest gas potential occurs in the Cretaceous structural play, principally because of its larger play area. The largest individual gas field is predicted to occur in the Cretaceous play as well, with a median size

estimate of  $32 \times 10^9 \text{ m}^3$  (1.1 TCF). Field-size rankings of all plays suggest about 40 per cent of the basin's total petroleum resource is contained in the five largest expected gas fields. This is comparable to average distributions of hydrocarbons in most basins worldwide (Klemme, 1984).


For Georgia Basin as a whole, the Cretaceous play contains about 70 per cent of the region's expected total gas volume and 8 of the 10 largest fields. Geographically, the most prospective area in Georgia Basin is the Bellingham sub-basin onshore including the Fraser River valley and delta in British Columbia and Whatcom County in Washington (Fig. 2, 3). The optimum combination of reservoir-quality sandstone, large traps, and abundant source material is present in the area. Less prospective areas in the Basin, in descending order, are the Strait of Georgia, the Nanaimo sub-basin of eastern Vancouver Island and Gulf Islands, the Suquash sub-basin, and the Comox sub-basin. Very little petroleum potential is expected on eastern Vancouver Island because of the paucity of reservoir-quality sandstone and source rock, particularly in the Comox sub-basin.

**Assessment results and exploration history.** Average exploration risks estimated in the assessment for the Georgia Basin region (for all three plays) suggest success rates for exploration drilling in the region should average about one in three, that is, the probability of finding accumulations of gas in Georgia Basin in all three plays is approximately 0.33. The lower success rate of historical drilling (that is, 15 former domestic gas-producing wells out of 122 wells drilled) may indicate a higher exploration risk than estimated here. However, a vast majority of these boreholes were originally drilled as water wells and are probably inadequate or only partly diagnostic petroleum tests. Many of the wells (81) were drilled before 1960 and presumably no geophysical surveys or mechanical well logs were used for locating many of the potential drillsites. Stratigraphy and structure deduced from limited outcrop exposure and lithological descriptions from earlier drilling provided the basis for petroleum exploration previous to 1960. The majority of seismic reflection survey work acquired in the 1960s characteristically exhibits poor to fair data quality compared to current techniques. No doubt many wells placed by these surveys were drilled in less than optimum locations. Only five wells have been completed since 1980, following development and refinement of modern seismic survey techniques.

**Distribution of resources in Canada.** Hydrocarbon plays in the Georgia region occupy areas on both sides of the International Border. If it can be assumed that the hydrocarbon resources are evenly distributed throughout the play area, the proportion of resource residing in Canada can be estimated by comparing play areas between the two countries. Although the location of the largest field cannot be



**Figure 33.** Georgia Tertiary structural gas play area. Major structural elements are illustrated.



**Figure 34.** Estimate of in-place gas potential of the Tertiary structural play in Georgia Basin. Median value of probabilistic assessment is  $59 \times 10^9 m^3$  of in-place gas distributed in 93 fields.

determined, there is no reason that all or part of the largest predicted field is not present in Canada. About two thirds of the Georgia Pleistocene stratigraphic play area is located in Canada, so the amount of estimated gas resource potential is  $138 \times 10^6 m^3$  (median value) in Canada. Similarly, two thirds of the play area of the Georgia Tertiary structural play is found in Canada, so the median gas potential for Canada is  $39.5 \times 10^9 m^3$ . In the Georgia Cretaceous structural play, 85 per cent of the play area is located in Canada. Assuming even distribution of the gas resource, the proportion of gas potential present in Canada is  $101 \times 10^9 m^3$  (median value).


## TOFINO ASSESSMENT REGION

### Exploration history and regional studies


Aeromagnetic and reflection and refraction seismic surveys were conducted in the 1960s by Shell Canada in the Tofino Basin. The aeromagnetic survey delineates long and linear total intensity magnetic anomalies parallel to the west coast of Vancouver Island (Shouldice, 1971).

Numerous academic and government seismic surveys and studies were also performed in the area. Continuous seismic profiles combined with bathymetry illustrate basin topography and geometry of the western Canadian continental margin (Chase et al., 1975). Side-scan sonar, 3.5 kHz profiles and submersible traverses carried out by the Geological Survey of Canada (GSC) and multichannel seismic lines acquired by Shell were used to delineate and investigate the Apollo structure in the Tofino Basin (see Fig. 38 for schematic geological section through the Apollo structure partly derived from seismic data) (Yorath, 1980). Numerous seismic reflection profiles acquired by Chevron, the University of British Columbia and the Geological Survey of Canada were used to provide a unified interpretation of the tectonics and structure of the Winona Basin (Davis and Riddihough, 1982). Four multichannel deep-crustal seismic lines were recorded onshore Vancouver Island as part of the LITHOPROBE programme. These lines depict the deep crustal structure and geometry of the subduction zone beneath the Island (Green et al., 1985; Yorath et al., 1985a, b; Sutherland Brown and Yorath, 1985; Green et al., 1986; Clowes et al., 1987). Multichannel seismic reflection lines in Tofino Basin collected in 1985 by the GSC along with acoustic imagery and bathymetry data were used to illustrate the surface morphology, internal geometry and deformation style of the accretionary prism (Fig. 4, 39) (Davis and Hyndman, 1989; Davis et al., 1990; Hyndman et al., 1990). A 1988 deep seismic reflection survey of the Queen Charlotte Basin region included one line that crossed the Winona Basin (Fig. 4, 40) (Rohr and Dietrich, 1990, 1991). This line illustrates the wedge-shaped sediment package overlying oceanic crust. Another marine multichannel seismic survey (722 km) acquired by the GSC in 1989 in Tofino Basin clarifies the regional structure and stratigraphy, examines the physical properties related to the bottom-simulating reflector interpreted as a methane hydrate layer, and studies the nature of the detachment that could produce large earthquakes (Spence et al., 1985; Singh et al., 1990). Dehler and Clowes (1992) and Clowes et al. (1987) used integrated geophysical studies to develop structural models across the continental margin west of Vancouver Island. Gravity and magnetic data, as well as seismic profiles, were interpreted to constrain the offshore positions of various accretionary terranes.


Shell Canada Ltd. drilled six offshore wells on seismically defined structures in the late 1960s in Tofino Basin (Fig. 2, Table 1) (Shell Canada Ltd., 1968f, g, h, i, 1969d, e). Four onshore wells were also drilled on the northern coast of Olympic Peninsula in the Juan de Fuca Basin (Fig. 2, Table 1) (Niem and Snively, 1991).



**Figure 35.** Georgia Pleistocene stratigraphic gas play area. Major structural elements are illustrated.



**Figure 36.** Estimate of in-place gas potential of the Pleistocene stratigraphic play in Georgia Basin. Median value of probabilistic assessment is  $207 \times 10^6 \text{m}^3$  of in-place gas distributed in 92 fields.



**Figure 37.** Estimate of total gas potential for the Georgia Basin region. Median value of probabilistic assessment is  $185 \times 10^9 \text{m}^3$  of in-place gas.

## Regional geology

### Geological setting and tectonic evolution

The Tofino Basin and adjacent accretionary wedge are upper Tertiary forearc and accretionary basins that developed in response to convergence and orthogonal subduction of the Juan de Fuca Plate beneath the continental margin. It is speculated that Winona Basin is a Plio-Pleistocene deep-water basin that developed by flexural bending in response to oblique convergence or transpression of the Explorer oceanic plate against the North American Plate (Davis and Riddihough, 1982; Yorath and Hyndman, 1983).


The Tofino Basin underlies some  $15\,000 \text{ km}^2$  of the continental shelf offshore Vancouver Island. The Juan de Fuca sub-basin represents a southern extension of the basin beneath Juan de Fuca Strait and adjacent coastal areas of Vancouver Island and Olympic Peninsula (Fig. 2).

The deep-water Winona Basin and accretionary wedge encompass a total area of about  $25\,000 \text{ km}^2$  seaward of the continental shelf edge. The Winona Basin is a Plio-Pleistocene basin at the foot of the continental slope. The accretionary wedge is a deformed Tertiary sedimentary


prism overlying oceanic crust of the Juan de Fuca Plate. Along the outer continental shelf, the accretionary wedge is underthrust beneath the Tofino Basin (Hyndman et al., 1990).

### Stratigraphy and structure

The Tofino Basin contains up to 6000 m of upper Paleogene–Neogene sedimentary strata. The basin-fill includes deep-marine mudstone and sandstone and minor conglomerate of the upper Paleogene Carmanah Formation and an overlying unnamed Neogene succession of marine sandstone, siltstone, and mudstone (Fig. 8). In the eastern Tofino Basin, ‘basement’ for petroleum exploration consists of Cretaceous metasedimentary rocks of the Pacific Rim Terrane and Eocene volcanic rocks of the Crescent Terrane (Fig. 39). In outer shelf areas where the basin overlies Paleogene accretionary wedge sedimentary rocks, there is no sharply defined ‘basement’. The accretionary wedge, seaward of the shelf edge, consists of up to 6000 m of Tertiary strata. The Winona Basin contains up to 8000 m of Pleistocene strata. Deposits in both areas consist exclusively of deep-marine mudstone and turbidite sandstone. On the northern shore of Olympic Peninsula, 6000 m of middle



**Figure 38.** Geological cross-section (derived from seismic and well data) in Tofino Basin (E-E'; location in Fig. 2) (modified from Yorath, 1980). Main faults are interpreted from seismic sections.




**Figure 39.** Seismic reflection profile, Pacific Ocean offshore, southern Vancouver Island (data from Spence et al., 1985; interpretation modified from Hyndman et al., 1990). The upper Tertiary–Quaternary Tofino Basin overlies deformed (accretionary wedge) Tertiary sedimentary strata and Mesozoic volcanic and metasedimentary rocks of the Pacific Rim and Crescent terranes. Structural features include compressional folds and thrust faults. The Prometheus well encountered gas shows in shallow Pleistocene sandstone.

Eocene to early Miocene sedimentary rocks unconformably overlie the lower Eocene Crescent Formation. These sediments consist of lithic turbidite sandstone, deep-marine mudstone, and minor polymict conglomerate and sedimentary breccia (Niem and Snavelly, 1991).

The most abundant structures in Tofino Basin are northwest-aligned curvilinear folds. One such fold, the Apollo anticline, occurs above an intra-Neogene detachment (Fig. 38). The folds developed in response to Plio-Pleistocene subduction of the Juan de Fuca Plate and accretionary wedge beneath the basin. In eastern and

southern Tofino Basin, Tertiary strata are variably disrupted by Neogene thrust, strike-slip and normal faults that developed in association with terrane underplating. Seismic surveys recorded by Shell Canada in the 1960s manifest the structural style in the deep-water offshore area. Generally, structures are large and dips are moderate so deformation is not extreme. Some of the structures involve volcanic 'basement'; others do not. Some of the compressional folds are faulted and some indicate an extremely complex structural history with small, episodic periods of growth over a long period of time.



**Figure 40.** Seismic reflection profile, Pacific Ocean offshore, northern Vancouver Island (data from Rohr and Dietrich, 1990). The wedge-shaped Winona Basin at base of the continental slope (1500–2200 m water depths) contains up to 8 km of Plio-Pleistocene sedimentary strata, overlying oceanic crust. Interbasin structural features include compressional folds and thrust faults.

## Petroleum geology

### Reservoirs

#### Tertiary

Deep-marine outer shelf to bathyal mudstone and siltstone units dominate the Tertiary sedimentary succession within the Tofino Basin (Fig. 8). There are rare nearshore marine and submarine channel sandstone and conglomerate. Porous and permeable sandstone and conglomerate units occur in the upper 2000 m of the sedimentary succession. These units occur as thin interbeds in the thick mudstone sequence. Although Shouldice (1971) and Yorath (1987) reported low porosities as a result of clay-mineral plugging, sidewall core samples retrieved from the six offshore wells drilled to date in the Tofino Basin and outcrop samples on the Olympic Peninsula of Washington in the Juan de Fuca sub-basin indicate fair to good porosity and permeability (Shell Canada Ltd., 1968f, g, h, i, 1969d, e; Snavely, 1987). Sidewall cores exhibit porosity values ranging from 20 to 46 per cent. Secondary porosity due to fracturing is probably minor in the basins. About 3 per cent of the total Tofino Basin succession consists of reservoir-quality strata.

In the Juan de Fuca sub-basin, turbiditic, coarse-grained, clastic material is more widespread than in the Tofino Basin. Niem and Snavely (1991) stated that, in general, these matrix-rich rocks have low to moderate reservoir potential as a result of diagenetic and detrital clays and siliceous cements clogging the pores. Outcrops of Eocene–Oligocene lithic sandstone are typically characterized by moderate porosity and low permeability, averaging around 20 per cent and 5 md., respectively. Rare, cleaner micaceous sandstone has improved reservoir quality, with measured values of up to 25 per cent porosity and 657 md. permeability. However Tertiary sections intersected in drillholes onshore indicate tight siltstone, mudstone, and minor turbiditic sandstone. Minor secondary fracture porosity was observed in these holes (Niem and Snavely, 1991).

#### Pleistocene

Pleistocene turbidite sandstone units likely occur in the Winona Basin and some may have reservoir potential. However, with no wells yet drilled in the basin, evaluation of reservoir characteristics remains equivocal.



## Seals

Impermeable shale and siltstone dominate the Tertiary clastic succession in these basins. Thin reservoir lenticular sandstone units abut against and are overlain by these fine-grained rocks. Therefore, more than adequate seal is present throughout the succession in the basins. In the Juan de Fuca sub-basin, thick impermeable mudstone and shale units are interbedded with thin reservoir-quality sandstone. However, Niem and Snively (1991) stated that most of the north-dipping reservoir units of the homocline in the sub-basin have been breached by erosion, which increases the prospect-level risk for seal in this particular part of the basin.

## Traps

Petroleum trap types in the Tofino Basin include simple anticlinal folds, faulted anticlines, normal and thrust faults and shale diapirs (Fig. 38, Table 3). Anomalously high geopressure gradients have been measured in the Tofino Basin (Shouldice, 1971). Injection of incompetent shale into overlying sedimentary rocks as a result of pressure buildups produced the diapiric structures that provide potential sites for petroleum accumulation on the crest or along the flanks of the diapir. Episodic compressional folding and/or fault-related deformation occurred from mid-Miocene to Pleistocene time. Shale diapirism postdated the folding episodes. Very large structures are observed on available seismic lines. Closure areas vary from approximately 12 to 145 km<sup>2</sup>, with an average closure of 25 km<sup>2</sup>. Vertical closure on traps ranges from 20 to 1100 m. Structures occur throughout the stratigraphic succession. In the southern part of Tofino Basin, the Neogene sedimentary rocks are essentially undeformed except at the shelf edge, where broad anticlines occur (Shouldice, 1971). In central Tofino Basin, curvilinear folds occur landward of the continental shelf (Fig. 38; Apollo anticline, Yorath, 1980). These anticlines are interpreted as having formed as a result of gravitational sliding along shallow detachment surfaces in Neogene and Quaternary sediments (Yorath, 1980).

The presence of isolated, lenticular sandstone within thick mudstone sections provides potential for stratigraphic traps within Tofino Basin. This trapping relationship has been documented by Niem and Snively (1991) in the Juan de Fuca sub-basin.


## Source rocks

Tertiary strata encountered in the offshore Tofino Basin have poor source-rock potential. Strata penetrated by exploration wells are characterized by terrestrial Type III organic matter, with low organic content (average 0.8% TOC) and low hydrogen index (Bustin, 1995). Strata with somewhat

different source rock characteristics have been identified along the margins of the Juan de Fuca sub-basin (Niem and Snively, 1991). Tertiary mudstone and accretionary ('melange') sedimentary rocks in this area, although still characterized by low organic content (<1% TOC), locally contain both Type III and (oil-prone) Type II organic material. A possibility exists that similar or richer source rocks may be present in parts of the accretionary wedge beneath the offshore Tofino Basin. The source-rock characteristics of Winona Basin strata are unknown.

## Source-rock maturation

Organic maturation data (R<sub>max</sub> and vitrinite reflectance) from Tofino Basin wells indicate that Tertiary strata are immature to marginally mature, with the depth to the top of the oil window occurring at about 2000 m (Bustin, 1995; Fig. 41). In onshore areas of Juan de Fuca sub-basin, Tertiary strata vary from immature to mature, with highest



**Figure 41.** Maturation–depth profile of strata in the Shell Anglo Pluto I-87 well location, Tofino Basin (see Fig. 2 and Table 1 for well location). Measured subsurface Ro and TMAX values obtained from Bustin (1995). Model-predicted maturation profile based on illustrated stratigraphy and a constant geothermal gradient of 25 C/km. Uplift and erosion of 1500 m subsequent to deposition of Middle Miocene mudstone and siltstone is estimated by extrapolating measured maturation gradients. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995).

maturation levels (vitrinite reflectance values of up to 0.75%) occurring in Eocene accretionary wedge sediments (Snively, 1987; Niem and Snively, 1991).

### Timing of hydrocarbon generation

Maturation models indicate hydrocarbon generation in Tofino Basin probably occurred in the late Tertiary, both during and after late stages of basin development and trap formation, that is, folding (e.g. Fig. 42). As such, there should be relatively little risk associated with timing of hydrocarbon generation in most parts of Tofino Basin.

Considered together, the known and expected variations in timing of hydrocarbon generation, trap development and source rock distributions indicate the complex geographic distribution of oil and gas accumulations within the region. Large hydrocarbon accumulations will probably be restricted to a small number of local areas or structures where optimum timing occurred between trap development and source-rock maturation.

### Hydrocarbon shows

Of the six offshore wells drilled in Tofino Basin, two encountered gas shows in shallow Neogene sandstone (Shouldice, 1971). One gas show was discovered in Tertiary

sandstone among the four wells drilled onshore Olympic Peninsula (Juan de Fuca sub-basin). In addition, three surface gas seeps have been reported from the onshore Juan de Fuca sub-basin (Niem and Snively, 1991). These gas seeps are believed to be thermogenic, with a source in underlying Tertiary melange. Surface exposures of melange mudstones contain local indications of oil and gas.

### Petroleum assessment

One all-encompassing conceptual gas play was identified for the western Vancouver Island offshore basins, including the Tofino and the Winona basins, and the Juan de Fuca sub-basin. Assessment computations were performed for natural gas only in the Tofino region of the West Coast. Oil assessments were not undertaken for these plays because geochemical data suggest the area is likely not oil-prone.

### Petroleum plays

#### Tofino Tertiary structural gas play

*Play definition.* The Tofino Tertiary gas play includes structural and stratigraphic traps in Tertiary and Quaternary strata in the Tofino and Winona basins and Juan de Fuca sub-basin. The play area covers the continental shelf offshore Vancouver Island and parts of the adjacent deep-water slope and Pacific basin (Fig. 2, 43). Western Juan de Fuca Strait and northwestern Olympic Peninsula are also included in the play.


*Geology.* Prospects include Tertiary or Quaternary sandstone units in large compressional folds, shale-diapir-cored anticlines and reverse-fault structural traps. Isolated sandstone encased in thick mudstone sections provides opportunities for stratigraphic trap potential.

*Exploration risks.* The exploration risk for the Tofino Tertiary play was estimated at 0.09, with most of the risk associated with the presence of source rocks and the adequacy of migration pathways (Appendix A, Table A-10b).


*Play potential.* The median estimate of play potential is  $266 \times 10^9 \text{ m}^3$  of in-place gas and the mean value estimate of number of fields is 41 (Fig. 44, Table 4). The largest field in the play is estimated (at median value) to contain  $25.9 \times 10^9 \text{ m}^3$  of in-place gas.

### Discussion of assessment results

*Resource distributions.* One exploration play has been assessed for the combined Tofino–Winona–Juan de Fuca basin area. The gas volume predicted is  $266 \times 10^9 \text{ m}^3$  (9.4



**Figure 42.** Subsidence and hydrocarbon generation model for area of Pluto I-87 well, Tofino Basin. Model derived from commercial basin modelling system, BasinMod 1-D (Platte River Associates, 1995). Hydrocarbon generation models based on standard (BasinMod) kinetic parameters for organic matter types (Type III for all Tertiary strata). See text for discussion.




**Figure 43.** Tofino Basin Tertiary structural gas play area. Major structural elements are illustrated.

TCF) distributed in 41 fields (Fig. 44). The largest individual field is estimated to contain  $26 \times 10^9 \text{ m}^3$  (0.9 TCF) of gas. Approximately 33 per cent of the region's total petroleum resource is concentrated in the five largest fields. This distribution indicates a moderate to low concentration of gas resource consistent with collisional convergent margin basins (Klemme, 1984).

The Juan de Fuca Strait area is considered the most prospective because of the presence of source rock, greater volumes of coarse clastic sediments (potential reservoir facies), and known gas seeps. One well on the Olympic Peninsula reported a gas flow ( $1416 \text{ m}^3/\text{day}$ ). The Tofino Basin offshore Vancouver Island may be somewhat less prospective as a result of greater uncertainties regarding source and reservoir rocks. The Winona Basin and accretionary wedge are considered the least prospective parts of the Tofino assessment region.

*Assessment results and exploration history.* The exploration risk estimated suggests success rates for exploratory drilling in the region should average 1 in 11. The absence of commercial discoveries in 10 wells drilled to date does not preclude the presence of economically viable fields in the region. Because of an exploration moratorium imposed in 1972 on the federal lands in the Pacific offshore of Canada,



**Figure 44.** Estimate of in-place gas potential of the Tertiary structural play in Tofino Basin region. Median value of probabilistic assessment is  $266 \times 10^9 \text{ m}^3$  of in-place gas distributed in 41 fields.

only one well has been completed (Olympic Peninsula) since 1980. The introduction of sophisticated seismic techniques in the late 1970s contributed to locating this well, the only one with a significant gas flow to date. Ten wells represent evaluation of a very small fraction of the total prospective area within the three basins. As noted previously, it is common for many unsuccessful wells to be drilled in a frontier area before the first discovery is made.

*Distribution of resources in Canada.* Assuming an even distribution of gas potential resource in the play, roughly 85 per cent of the play is in Canadian jurisdiction. Therefore, about  $226 \times 10^9 \text{ m}^3$  of gas may occur in Canada (median value).

## BASIN COMPARISONS

The Queen Charlotte Basin region has been compared to the petroleum-producing Cook Inlet Basin (Haimala and Procter, 1982; Hamilton and Cameron, 1989; Thompson et al., 1991). Similarities between the two regions include the general ages and types of reservoir strata, source rocks, regional unconformities and structures, and common occurrences of surface hydrocarbon shows. The main difference between the two regions is in their Neogene structural history; the Cook Inlet Basin developed in a forearc setting and contains comparatively fewer strike-slip-related extensional faults than does the Queen Charlotte Basin. Total petroleum resources (produced and remaining) in the  $20\,000 \text{ km}^2$  Cook Inlet Basin are about  $2.2 \times 10^9$  bbl of oil and 10 TCF of gas (Magoon and Kirschner, 1990). Most of the Cook Inlet petroleum accumulations occur in Oligocene–Miocene conglomeratic sandstone units within thrust faulted, Pliocene anticlines (similar to many Pliocene structures in the Queen Charlotte Basin). The largest individual oil and gas fields in Cook Inlet are the McArthur River field, with original oil reserves of  $90 \times 10^6 \text{ m}^3$  ( $570 \times 10^6$  bbl), and the Kenai field, with original gas reserves of  $65 \times 10^9 \text{ m}^3$  (2.3 TCF). These field sizes are of comparable magnitude to the median size estimates of the largest oil and gas fields predicted for the Queen Charlotte Basin.

Some debate has occurred concerning the tectonic setting for Georgia Basin. England and Bustin (1998) implied that Georgia Basin occupies the inner forearc position in the convergent margin plate tectonic model, while Mustard (1994) proposed a foreland model for the Cretaceous Georgia Basin as a result of multiple source regions for the sedimentary succession and the presence of fold and thrust belts. The largest gas field of the eastern Cordilleran foreland fold and thrust belt comprises the Cretaceous gas pools of the Ricinus Field. The Cretaceous pools of this field have an initial in-place volume of  $28\,400 \times 10^6 \text{ m}^3$ , comparable to the largest predicted field size for the Georgia Cretaceous play ( $31\,977 \times 10^6 \text{ m}^3$ ). Comparing a proposed forearc Cretaceous

Georgia Basin with the Cook Inlet Basin (forearc setting) reveals predicted largest field sizes less than half the size of the Kenai gas field ( $32 \times 10^9 \text{m}^3$  vs.  $65 \times 10^9 \text{m}^3$ ). Analogous gas-bearing basins for the strike-slip pull-apart model proposed for Tertiary Georgia Basin are the Mackenzie Delta basin of northern Canada (Dixon et al., 1994) and the Willamette Basin of northern Oregon containing the Mist Field (Armentrout and Suek, 1985). Active underplating in the Tofino and Juan de Fuca Basin regions has produced local areas of normal, strike-slip and thrust faulting in the overlying sedimentary rocks. Occasional thrust-faulted anticlines in the Tertiary succession reflect trap-forming structures analogous to similar circum-Pacific convergent margin forearc basins (Japan Trench; Aleutian Trench).

Other northeast Pacific region petroleum fields with past oil or gas production include three small onshore fields in Washington and Oregon. The commercial Mist Lake gas field was discovered in the forearc Willamette Basin of northern Oregon in 1979 (Armentrout and Suek, 1985). The Mist Lake gas reservoir occurs in Eocene quartzofeldspathic sandstone with compositional and reservoir characteristics similar to Neogene sandstone in the Queen Charlotte and Georgia basins.

The Queen Charlotte Basin region has also been compared to the southern California continental borderland, based on similarities in Neogene tectonic history and structural characteristics (Rohr and Dietrich, 1992). Numerous petroleum-bearing Neogene strike-slip basins occur in the California borderland region, including the oil-rich Los Angeles Basin, with reserves of over  $10^{10}$  bbl (Biddle, 1992). Differences in types of petroleum source rocks in the California and Queen Charlotte basins preclude making direct petroleum endowment comparisons between the two regions. However, from a general perspective of basin types, strike-slip basins are known worldwide to be above average in hydrocarbon richness, on a sedimentary rock volume or area basis (Price, 1994). Factors that contribute to hydrocarbon enrichment in strike-slip basins include high paleo-heat-flow and extensive faulting of source and reservoir rocks, two features that characterize the Queen Charlotte Basin region. Tertiary Georgia Basin has some strike-slip basin characteristics (Mustard and Rouse, 1994) and may be comparable to some southern California basins in terms of structural styles.

There are no direct geological analogues for the British Columbia offshore region in other Canadian frontiers. However, by comparing the magnitude of resource estimates with other Canadian frontier regions, a ranking of potential can be achieved. Note that recoverable resources are quoted here for comparison purposes. Compared to the median recoverable resource estimate of  $2.6 \times 10^9$  bbl of oil and 20 TCF of gas in the Queen Charlotte Basin region, median estimates of resource endowment for Canadian East Coast


and Arctic basins include  $4.7 \times 10^9$  bbl of oil and 13 TCF of gas for the offshore Newfoundland Jeanne D'Arc Basin (Procter et al., 1983; Sinclair et al., 1992), 18 TCF of gas and  $1 \times 10^9$  bbl of oil/condensate for the Scotian Shelf Basin (Wade et al., 1989), and  $7 \times 10^9$  bbl of oil and 68 TCF of gas for the Beaufort-Mackenzie Basin (Dixon et al., 1994). Median resource estimates for recoverable gas in the Georgia Basin are 5.8 TCF and in the Tofino Basin region 8.5 TCF (recovery factor: 0.9). In comparing Canada's east and west coast basins, the present assessment indicates the Queen Charlotte Basin may have a gas resource endowment comparable to the Scotian Shelf Basin and an oil resource endowment about half that of the Jeanne D'Arc Basin. Gas resource estimates for the combined Tofino-Georgia basins are comparable to the Jeanne D'Arc Basin endowment.

## CONCLUSIONS

The oil and gas resource potential of Canada's west coast basins has been evaluated through regional petroleum play assessments. The quantitative assessments were derived using the Geological Survey of Canada's (PETRIMES) assessment methodology system. The assessments included analyses of 10 conceptual plays, each of which incorporated the calculation or estimation of field size parametric data, numbers of prospects and exploration risks. Oil and gas volumes reported for these conceptual plays are total statistical estimates of the resource present 'in the ground', not the gas volume that is economically producible. Individual field-size determinations will be important in identifying which plays are attractive for exploration programs.

Median estimates for total oil and gas potential for all Canadian west coast basins are  $1560 \times 10^6 \text{m}^3$  of in-place oil and  $1228 \times 10^9 \text{m}^3$  of in-place gas (Fig. 24, 45). In-place oil potential is restricted exclusively to the Queen Charlotte Basin region; gas potential encompasses all Canadian west coast basins.

The potential for significant petroleum accumulations in the Queen Charlotte Basin and environs is indicated by the combined presence of abundant reservoir strata, good petroleum source rocks, numerous and diverse structural and stratigraphic traps and common occurrence of oil and gas shows. Quantitative assessments of six petroleum plays in the Queen Charlotte Basin region provide estimates of total resource potential of  $1560 \times 10^6 \text{m}^3$  ( $9.8 \times 10^9$  bbl) of in-place oil and  $734 \times 10^9 \text{m}^3$  (25.9 TCF) of in-place gas (median values). In terms of number of predicted fields and energy-equivalent volumes, estimated gas resources are more abundant than oil. The ranges of oil and gas estimates from high to low probability reflect the level of uncertainty in assessing petroleum potential for this region. However, in comparative terms, the estimates from the current



**Figure 45.** Estimate of total gas potential for west coast basins. Median value of probabilistic assessment is  $1228 \times 10^9 \text{ m}^3$  of in-place gas.

assessment are substantially higher than those derived in the Geological Survey of Canada's 1982 assessment. The higher resource estimates in this assessment reflect several factors, including more optimistic evaluations of the number of prospects, and volume and quality of potential reservoirs. In addition, the indications of a Jurassic-source Miocene reservoir petroleum system occurring offshore is considered significant, since large oil accumulations will probably occur as part of such a petroleum system.

Historical natural gas production from Pleistocene sandstone confirms the existence of petroleum accumulations within Georgia Basin. Although the ultimate play potential and field sizes are small for the Pleistocene play, sufficient favourable play conditions are present for significant gas accumulations in the Tertiary and Cretaceous sedimentary successions. The combination of ample reservoir-quality material, adequate gas source rock, and abundant petroleum-trapping configurations provide the necessary elements for potential petroleum accumulations. The total resource potential in the Georgia Basin is  $185 \times 10^9 \text{ m}^3$  (6.5 TCF) of in-place raw gas (median). Currently available geochemical information indicates there is probably little or no oil resource potential in the Georgia or Tofino basins.

The potential for gas resources in the Tofino Basin is indicated by gas seeps and gas shows in onshore portions of the basin on the Olympic Peninsula. The total potential for the Tofino Basin assessment region is estimated (at a median value) to be  $266 \times 10^9 \text{ m}^3$  (9.4 TCF) of in-place gas.

Significant upside potential for natural gas was recognized in three plays. The most attractive plays for natural gas exploration, in decreasing order in terms of potential and largest field size, are 1) the Queen Charlotte Pliocene play, 2) the Queen Charlotte Miocene play, and 3) the Tofino–Winona–Juan de Fuca Tertiary structural play. Oil plays, in decreasing order of potential, are 1) the Queen Charlotte Miocene play, 2) the Queen Charlotte Pliocene play, and 3) the Queen Charlotte Cretaceous play.

This assessment provides a favourable geological basis for further petroleum evaluation and exploration in the British Columbia coastal region. The complex geology and anticipated high exploration risks associated with the plays suggest that considerable amounts of new seismic data and many exploration wells may be required to properly evaluate the region's oil and gas potential. The present assessment suggests that substantial petroleum resources remain to be discovered in the sedimentary basins on the Pacific margin of Canada.

## REFERENCES

### Armentrout, J.M. and Suek, D.H.

1985: Hydrocarbon exploration in western Washington and Oregon; American Association of Petroleum Geologists, Bulletin, v. 69, no. 4, p. 627–643.

### Barrie, J.V.

1988: Surficial geology of Hecate Strait, British Columbia continental shelf; Geological Survey of Canada, Open File 1682.

### Biddle, K.T.

1992: The Los Angeles Basin; in *Active Margin Basins*, (ed.) K.T. Biddle; American Association of Petroleum Geologists, Memoir 32, p. 5–23.

### Bird, T.D., Barclay, J.E., Campbell, R.I., and Lee, P.J.

1994: Triassic gas resources of the Western Canada Sedimentary Basin, Part I. Geological play analysis and resource assessment; Geological Survey of Canada, Bulletin 483, 66 p.

### Bow Valley Industries

1984: Well history report, Bow Valley et al. Naden b-A27-J; British Columbia Department of Mines and Petroleum Resources, Open File WA06110.

### British American Oil Company Ltd.

1965: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 1275.

**British Petroleum Resources Canada Ltd.**

1987a: Well history report, BP Yellow Point d-84-C; British Columbia Department of Mines and Petroleum Resources, Open File WA06451.

1987b: Well history report, BP Laurel Harmac c-36-F; British Columbia Department of Mines and Petroleum Resources, Open File WA06452.

**Bustin, R.M.**

1990: Stratigraphy, sedimentology, and petroleum source rock potential of the Georgia Basin, southwest British Columbia and northwest Washington State; *in* Current Research, Part F; Geological Survey of Canada, Paper 90-1F, p. 103–108.

1995: Organic maturation and petroleum source rock potential of Tofino Basin, southwestern British Columbia; *Bulletin of Canadian Petroleum Geology*, v. 43, no. 2, p. 177–186.

**Bustin, R.M. and England, T.D.J.**

1991: Petroleum source rock potential of the Nanaimo Group, western margin of the Georgia Basin, southwestern British Columbia; *in* Current Research, Part A; Geological Survey of Canada, Paper 91-1A, p. 143–145.

**Bustin, R.M., Vellutini, D., and Goodarzi, F.**

1990: Petroleum source rock characteristics of the Tertiary Skonun Formation, Queen Charlotte Islands, Hecate Strait and Queen Charlotte Sound, British Columbia; *in* Current Research, Part F; Geological Survey of Canada, Paper 90-1F, p. 87–93.

**Campbell, R.B., Dodds, C.J., Yorath, C.J., and Sutherland Brown, A.**

1991: Part A. Insular Belt; *in* Structural styles, Chapter 17 of *Geology of the Cordilleran Orogen in Canada*, (ed.) H. Gabrielse and C. J. Yorath; Geological Survey of Canada, *Geology of Canada*, no. 4, p. 574–581 (*also* Geological Society of America, *The Geology of North America*, v. G-2, p. 574–581).

**Canadian Superior Oil Ltd.**

1962: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 1049.

**Carlisle, D. and Susuki, T.**

1974: Emergent basalt and submergent carbonate-clastic sequences including the Upper Triassic Dilleri and Welleri Zones on Vancouver Island; *Canadian Journal of Earth Sciences*, v. 11, p. 254–279.

**Chase, R.L., Tiffin, D.L., and Murray, J.W.**

1975: The western Canadian continental margin; *in* *Canada's Continental Margins and Offshore Petroleum Exploration*, (ed.) C.J. Yorath, E.R. Parker, and D.J. Glass; Canadian Society of Petroleum Geologists, *Memoir 4*, p. 701–722.

**Clague, J.J.**

1994: Quaternary stratigraphy and history of south-coastal British Columbia; *in* *Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia*, (ed.) J.W.H. Monger; Geological Survey of Canada, *Bulletin 481*, p. 181–192.

**Clague, J.J., Luternauer, J.L., and Hebda, R.J.**

1983: Sedimentary environments and postglacial history of the Fraser Delta and lower Fraser Valley, British Columbia; *Canadian Journal of Earth Sciences*, v. 20, p. 1314–1326.

**Clowes, R.M., Baird, D.J., and Dehler, S.A.**

1997: Crustal structure of the Cascadia subduction zone, southwestern British Columbia, from potential field and seismic studies; *Canadian Journal of Earth Sciences*, v. 34, p. 317–335.

**Clowes, R.M., Brandon, M.T., Green, A.G., Yorath, C.J., Sutherland Brown, A., Kanasevich, E.R., and Spencer, C.**

1987: LITHOPROBE- southern Vancouver Island: Cenozoic subduction complex imaged by deep seismic reflections; *Canadian Journal of Earth Sciences*, v. 24, p. 31–51.

**Coney, P.J., Jones, D.L., and Monger, J.W.H.**

1980: Cordilleran suspect terranes; *Nature*, v. 288, p. 329–333.

**Daly, R.A.**

1912: Geology of the North American Cordillera at the forty-ninth parallel; Geological Survey of Canada, *Memoir 38*, 857 p.

**Davis, E.E. and Hyndman, R.D.**

1989: Accretion and recent deformation of sediments along the northern Cascadia subduction zone; *Geological Society of America, Bulletin*, v. 101, p. 1465–1480.

**Davis, E.E. and Riddihough, R.P.**

1982: The Winona Basin: structure and tectonics; *Canadian Journal of Earth Sciences*, v. 19, p. 767–788.

**Davis, E.E., Hyndman, R.D., and Villinger, H.**

1990: Rates of fluid expulsion across the northern Cascadia accretionary prism: constraints from new heat flow and multichannel seismic data; *Journal of Geophysical Research*, v. 95, no. B6, p. 8869–8889.

**Dehler, S.A. and Clowes, R.M.**

1992: Integrated geophysical modelling of terranes and other structural features along the western Canadian margin; *Canadian Journal of Earth Sciences*, v. 29, p. 1492–1508.

**Dickinson, W.R.**

1976: Sedimentary basins developed during evolution of Mesozoic–Cenozoic arc-trench system in western North America; *Canadian Journal of Earth Sciences*, v. 13, p. 1268–1287.

**Dickinson, W.R. and Seely, D.R.**

1979: Structure and stratigraphy of forearc regions; *American Association of Petroleum Geologists, Bulletin*, v. 63, p. 2–31.

**Dietrich, J.R.**

1995: Petroleum resource potential of the Queen Charlotte Basin and environs, west coast Canada; *Bulletin of Canadian Petroleum Geology*, v. 43, no. 1, p. 20–34.

**Dietrich, J.R., Higgs, R., Rohr, K.M., and White, J.M.**

1993: The Tertiary Queen Charlotte Basin: a strike-slip basin on the western Canadian continental margin; *in* *Tectonic Controls and Signatures in Sedimentary Successions*, (ed.) L. Frostick and R. Steel; International Association of Sedimentologists, *Special Publication Number 20*, p. 161–169.

**Dietrich, J. R., Morrell, G.R., and Fortier, M.C.**

1992: Petroleum resource potential in the proposed area of Gwaii Haanas/South Moresby National Marine Park, British Columbia; Geological Survey of Canada, Open File 2557.

**Dixon J., Morrell, G.R., Dietrich, J.R., Procter, R.M., and Taylor, G.C.**

1994: Petroleum resources of the Mackenzie Delta and Beaufort Sea; Geological Survey of Canada, *Bulletin 474*, 44 p.

- England, T.D.J.**  
1990: Late Cretaceous to Paleogene evolution of the Georgia Basin, southwestern British Columbia; Ph.D. thesis, Memorial University of Newfoundland, 481 p.
- 1991: Late Cretaceous to Paleogene structural and stratigraphic evolution of Georgia Basin, southwestern British Columbia: implications for hydrocarbon potential; *Washington Geology*, v. 19, no. 4, p. 10–11.
- England, T.D.J. and Bustin, R.M.**  
1998: Architecture of the Georgia Basin, southwestern British Columbia; *Bulletin of Canadian Petroleum Geology*, v. 46, no. 2, p. 288–320.
- England, T.D.J. and Calon, T.J.**  
1991: The Cowichan fold and thrust system, Vancouver Island, southwestern British Columbia; *Geological Society of America, Bulletin*, v. 103, p. 336–362.
- England, T.D.J. and Hiscott, R.N.**  
1992: Lithostratigraphy and deep-water setting of the upper Nanaimo Group (Upper Cretaceous), outer Gulf Islands of southwestern British Columbia; *Canadian Journal of Earth Sciences*, v. 29, p. 574–595.
- England, T.D.J., Currie, L.D., Massey, N.W.D., Roden-Tice, M.K., and Miller D.S.**  
1997: Apatite fission-track dating of the Cowichan fold and thrust system, southern Vancouver Island, British Columbia; *Canadian Journal of Earth Sciences*, v. 34, p. 635–645.
- England, T.D.J., Piggott, N., and Douglas, T.R.**  
1989: Hydrocarbon charge control on the prospectivity of the Georgia Basin, Vancouver Island, British Columbia [abstract]; *in* Exploration Update '89: CSEG/CSPG Annual Meeting, Program with Abstracts.
- Fogarassy, J.A.S. and Barnes, W.C.**  
1988: Petroleum reservoir aspects of Middle to Upper Cretaceous and Tertiary strata of the Queen Charlotte Islands; *in* Some Aspects of the Petroleum Geology of the Queen Charlotte Islands, (ed.) R. Higgs; Canadian Society of Petroleum Geologists, Field Trip Guide, p. 22–25.
- 1991: Stratigraphy and diagenesis of the Middle to Upper Cretaceous Queen Charlotte Group, Queen Charlotte Islands, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.S. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 279–294.
- Fowler, M.G., Snowdon, L.R., Brooks, P.W., and Hamilton, T.S.**  
1987: Biomarker characterisation and hydrous pyrolysis of bitumen from Tertiary volcanics, Queen Charlotte Islands, British Columbia, Canada; *Organic Geochemistry*, v. 13, p. 715–725.
- Galloway, W.E.**  
1974: Deposition and diagenetic alteration of sandstone in northeast Pacific arc-related basins: implications for graywacke genesis; *Geological Society of America, Bulletin*, v. 85, p. 379–390.
- Glover, S.L.**  
1935: Oil and gas possibilities of western Whatcom County; Washington Division of Geology, Report of Investigations 2, 69 p.
- Gordy, P.L.**  
1988: Evaluation of the hydrocarbon potential of the Georgia Depression; British Columbia Ministry of Energy, Mines and Petroleum Resources, 80 p.
- Green, A.G., Clowes, R.M., and Yorath, C.J.**  
1985: LITHOPROBE seismic reflection profiles from southeastern Vancouver Island; Geological Survey of Canada, Open File 1180, 24 p.
- Green, A.G., Clowes, R.M., Yorath, C.J., Spencer, C., Kanasewich, E.R., Brandon, M.T., and Sutherland Brown, A.**  
1986: Seismic reflection imaging of the subducting Juan de Fuca Plate; *Nature*, v. 319, p. 210–213.
- Haggart, J.W.**  
1991: A synthesis of Cretaceous stratigraphy, Queen Charlotte Islands, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 253–277.
- 1992: Progress in Jurassic and Cretaceous stratigraphy, Queen Charlotte Islands, British Columbia; *in* Current Research, Part A; Geological Survey of Canada, Paper 92-A, p. 361–365.
- 1993: Latest Jurassic and Cretaceous paleogeography of the northern Insular Belt, British Columbia; *in* Mesozoic paleogeography of the Western United States – II, (ed.) G. Dunne and K. McDougall; Society of Economic Paleontologists and Mineralogists, Pacific Section, Book 71, p. 463–475.
- Haimala, N.E. and Procter, R.M.**  
1982: Hydrocarbon potential of offshore British Columbia; Geological Survey of Canada, Open File 824, 28 p.
- Hamilton, T.S. and Cameron, B.E.B.**  
1989: Hydrocarbon occurrences on the western margin of the Queen Charlotte Basin; *Bulletin of Canadian Petroleum Geology*, v. 34, no. 4, p. 443–466.
- Hamilton, T.S. and Dostal, J.**  
1994: Middle Tertiary eruptive rocks in the Vancouver area; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W.H. Monger; Geological Survey of Canada, Bulletin 481, p. 171–179.
- Hamilton, T.S. and Ricketts, B.D.**  
1994: Contour map of the sub-Quaternary bedrock surface, Strait of Georgia and Fraser Lowland; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W.H. Monger; Geological Survey of Canada, Bulletin 481, p. 193–196.
- Higgs, R.**  
1989: Sedimentology and implications for hydrocarbon exploration of the “Hippa Beds”, Queen Charlotte Islands, British Columbia; *in* Current Research, Part H; Geological Survey of Canada, Paper 89-1H, p. 53–58.
- 1991: Sedimentology, basin-fill architecture and petroleum geology of the Tertiary Queen Charlotte Basin, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 337–371.



- Hurst, P.D.**  
1991: Petroleum geology of the Bellingham Basin, Washington and evaluation of the AH-EL and partners Birch Bay No. 1 well; *Washington Geology*, v. 19, no. 4, p. 16–18.
- Hyndman, R.D., Yorath, C.J., Clowes, R.M., and Davis, E.E.**  
1990: The northern Cascadia subduction zone at Vancouver Island: seismic structure and tectonic history; *Canadian Journal of Earth Sciences*, v. 27, p. 313–329.
- Isachsen, C., Armstrong, R.L., and Parrish, R.R.**  
1985: U-Pb, Rb-Sr, and K-Ar geochronometry of Vancouver Island igneous rocks; *in* Programmes and Abstracts, A Symposium on Deep Structure of Southern Vancouver Island; Results of LITHOPROBE Phase 1; Geological Association of Canada, Victoria Section, p. 21–22.
- Johnson, S.Y.**  
1982: Stratigraphy, sedimentology, and tectonic setting of the Eocene Chuckanut Formation, northwest Washington; Ph.D. thesis, University of Washington, Seattle, Washington, 222 p.  
1984: Stratigraphy, age, and paleogeography of the Eocene Chuckanut Formation, northwest Washington; *Canadian Journal of Earth Sciences*, v. 21, no. 1, p. 92–106.  
1991: Sedimentation and tectonic setting of the Chuckanut Formation, northwest Washington; *Washington Geology*, v. 19, no. 4, p. 12–13.
- Johnston, W.A.**  
1921: Sedimentation of the Fraser River delta; Geological Survey of Canada, Memoir 125, 46 p.  
1923: Geology of Fraser River delta map-area; Geological Survey of Canada, Memoir 135, 87 p.
- Kay, M.**  
1951: North American geosynclines; Geological Society of America, Memoir 48, 143 p.
- Key Oil and Gas (1955) Ltd.**  
1958: Well history report, Hercon Key Evans No. 1; British Columbia Department of Mines and Petroleum Resources, Open File WA00162.
- King, P.B.**  
1969: Tectonic map of North America; United States Geological Survey, scale 1:5,000,000.
- Klemme, H.D.**  
1984: Field-size distribution related to basin characteristics; *in* Petroleum Resource Assessment, (ed.) C.D. Masters; International Union of Geological Sciences, Publication No. 17, p. 95–121.
- Lee, P.J.**  
1993: Two decades of Geological Survey of Canada petroleum resource assessments; *Canadian Journal of Earth Sciences*, v. 30, p. 321–332.
- Lee, P.J. and Tzeng, H.P.**  
1989: The petroleum exploration and resource evaluation system (PETRIMES): working reference guide; Institute of Sedimentary and Petroleum Geology, Calgary, Alberta, 258 p.
- Lee, P.J. and Wang, P.C.C.**  
1990: An introduction to petroleum resource evaluation methods; Canadian Society of Petroleum Geologists, 1990 Convention on Basin Perspectives, Short Courses Program: SC-2 Petroleum Resource Evaluation, 108 p.
- Lewis, T.J., Bentkowski, W.H., and Hyndman, R.D.**  
1992: Crustal temperatures near the Lithoprobe Southern Canadian Cordillera Transect; *Canadian Journal of Earth Sciences*, v. 29, p. 1197–1214.
- Lewis, P.D., Haggart, J.W., Anderson, R.G., Hickson, C.J., Thompson, R.I., Dietrich, J.R., and Rohr, K.M.**  
1991: Triassic to Neogene geologic evolution of the Queen Charlotte region; *Canadian Journal of Earth Sciences*, v. 28, p. 854–868.
- Lingley, W.S. and von der Dick, H.**  
1991: Petroleum geochemistry of Washington – a summary; *Washington Geology*, v. 19, no. 4, p. 23–27.
- Lowe, C. and Dehler, S.A.**  
1995: Crustal thickness beneath the Queen Charlotte Basin, Canada: results of a seismically constrained gravity inversion. *Journal of Geophysical Research*, v. 100, no. B12; p. 24, 331–24, 345.
- Luternauer, J.L., Barrie, J.V., Christian, H.A., Clague, J.J., Evoy, R.W., Hart, B.S., Hunter, J.A., Killeen, P.G., Kostaschuk, R.A., Mathewes, R.W., Monahan, P.A., Moslow, T.F., Mwenifumbo, C.J., Olynyk, H.W., Patterson, R.T., Pullan, S.E., Roberts, M.C., Robertson, P.K., Tarbotton, M.R., and Woeller, D.J.**  
1994: Fraser River delta: geology, geohazards and human impact; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W. Monger; Geological Survey of Canada, Bulletin 481, p. 197–220.
- Lyatsky, H.V.**  
1991: Regional geophysical constraints on crustal structure and geologic evolution of the Insular Belt, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 97–106.
- Lyatsky, H.V. and Haggart, J.W.**  
1993: Petroleum exploration model for the Queen Charlotte Basin, offshore British Columbia; *Canadian Journal of Earth Sciences*, v. 30, no. 5, p. 918–927.
- Lynch, J.V.G.**  
1991: Georgia Basin Project: stratigraphy and structure of Gambier Group rocks in the Howe Sound–Mamquam River area, southwest Coast Belt, British Columbia; *in* Current Research, Part A; Geological Survey of Canada, Paper 91-1A, p. 49–57.  
1992: Deformation of Early Cretaceous volcanic-arc assemblages, southern Coast Belt, British Columbia; *Canadian Journal of Earth Sciences*, v. 29, p. 2706–2721.
- Macauley, G.**  
1983: Source rock-oil shale potential of the Jurassic Kunga Formation, Queen Charlotte Islands; Geological Survey of Canada, Open File 921.
- Magoon, L.B. and Kirschner, C.E.**  
1990: Geology and petroleum resource potential of six onshore Alaska provinces; United States Geological Survey, Open File Report 88-450T.

- Masters, C.D.**  
1984: Petroleum resource assessment; International Union of Geological Sciences, Publication No. 17, 157 p.
- McFarland, C.R.**  
1983: Oil and gas exploration in Washington, 1900–1982; Washington Department of Natural Resources, Division of Geology and Earth Resources, Information Circular 75, 119 p.
- Moen, W.S.**  
1969: Mines and mineral deposits of Whatcom County, Washington; Washington Department of Natural Resources, Division of Mines and Geology, Bulletin 57, 134 p.
- Monger, J.W.H.**  
1991a: Late Mesozoic to Recent evolution of the Georgia Strait – Puget Sound region, British Columbia and Washington. *Washington Geology*, v. 19, no. 4, p. 3–7.  
1991b: Georgia Basin Project: structural evolution of parts of southern Insular and southwestern Coast belts, British Columbia; *in* Current Research, Part A, Geological Survey of Canada, Paper 91-1A, p. 219–228.
- Monger, J.W.H. and Journeay, J.M.**  
1994: Basement geology and tectonic evolution of the Vancouver region; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W.H. Monger; Geological Survey of Canada, Bulletin 481, p. 3–25.
- Monger, J.W.H., Price, R.A., and Templeman-Kluit, D.J.**  
1982: Tectonic accretion and the origin of two major metamorphic and plutonic belts in the Canadian Cordillera; *Geology*, v. 10, p. 70–75.
- Monger, J. W. H., Souther, J. G., and Gabrielse, H.**  
1972: Evolution of the Canadian Cordillera: a plate tectonic model; *American Journal of Science*, v. 272, p. 577–602.
- Monger, J.W.H., Wheeler, J.O., Tipper, H.W., Gabrielse, H., Harms, T., Struik, L.C., Campbell, R.B., Dodds, C.J., Gehrels, G.E., and O'Brien, J.**  
1991: Part B. Cordilleran terranes; *in* Chapter 8 of Geology of the Cordilleran Orogen in Canada, (ed.) H. Gabrielse and C.J. Yorath; Geological Survey of Canada, Geology of Canada, no. 4, p. 281–327 (*also* Geological Society of America, The Geology of North America, v. G-2, p. 281–327).
- Muller, J.E.**  
1977: Evolution of the Pacific margin, Vancouver Island and adjacent regions; *Canadian Journal of Earth Sciences*, v. 14, no. 9, p. 2062–2085.
- Mustard, P.S.**  
1991: Stratigraphy and sedimentology of the Georgia Basin, British Columbia and Washington State; *Washington Geology*, v. 19, no. 4, p. 7–9.  
1994: The Upper Cretaceous Nanaimo Group, Georgia Basin; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W.H. Monger; Geological Survey of Canada, Bulletin 481, p. 27–95.
- Mustard, P.S. and Rouse, G.E.**  
1991: Sedimentary outliers of the eastern Georgia Basin margin, British Columbia; *in* Current Research, Part A; Geological Survey of Canada, Paper 91-1A, p. 229–240.
- 1994: Stratigraphy and evolution of Tertiary Georgia Basin and subjacent Upper Cretaceous sedimentary rocks, southwestern British Columbia and northwestern Washington State; *in* Geology and Geological Hazards of the Vancouver Region, Southwestern British Columbia, (ed.) J.W.H. Monger; Geological Survey of Canada, Bulletin 481, p. 97–169.
- Niem, A.R. and Snaveley, P.D.**  
1991: Geology and preliminary hydrocarbon evaluation of the Tertiary Juan de Fuca Basin, Olympic Peninsula, northwest Washington; *Washington Geology*, v. 19, no. 4, p. 27–34.
- Orchard, M.J. and Forster, P.J.L.**  
1991: Conodont colour and thermal maturity of the Late Triassic Kunga Group, Queen Charlotte Islands, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 453–464.
- Pacht, J.A.**  
1984: Petrologic evolution and paleogeography of the Late Cretaceous Nanaimo Basin, Washington and British Columbia: implications for Cretaceous tectonics; *Geological Society of America, Bulletin*, v. 95, p. 766–778.
- Pacific Petroleum Ltd.**  
1959: Well history report, Charter et al. Saturna No. 1; British Columbia Department of Mines and Petroleum Resources, Open File WA00292.
- Petal Company Ltd.**  
1959: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report No. 823.
- Platte River Associates**  
1995: BasinMod I-D; Basin modelling software system, Platte River Associates.
- Podruski, J.A., Barclay, J.E., Hamblin, A.P., Lee, P.J., Osadetz, K.G., Procter, R.M., and Taylor, G.C.**  
1988: Conventional oil resources of Western Canada (light and medium), Part I: Resource endowment; Geological Survey of Canada, Paper 87-26, 125 p.
- Potential Gas Committee**  
1990: Definitions and procedures for estimation of potential gas resources; Potential Gas Agency, Colorado School of Mines.
- Price, L.C.**  
1994: Basin richness and source rock disruption: a fundamental relationship; *Journal of Petroleum Geology*, v. 17, no. 1, p. 5–38.
- Prims, J., Furlong, K.P., Rohr, K.M.M., and Govers, R.**  
1997: Lithospheric structure along the Queen Charlotte margin in western Canada: constraints from flexural modeling; *Geophysical Research Letters*, v. 17, p. 94–99.
- Procter, R.M., Taylor, G.C., and Wade, J.A.**  
1983: Oil and natural gas resources of Canada; Geological Survey of Canada, Paper 83-31, 59 p.
- Reinson, G. E., Lee, P.J., Warters, W., Osadetz, K.G., Bell, L.L., Price, P.R., Trollope, F., Campbell, R.I., and Barclay, J.E.**  
1993: Devonian gas resources of the Western Canada Sedimentary Basin; Part I: Geological play analysis and resource assessment; Geological Survey of Canada, Bulletin 452, 127 p.

**Rice, D.D.**

1986: Oil and gas assessment – methods and applications; American Association of Petroleum Geology, Studies in Geology, no. 21, 267 p.

**Richfield Oil Corporation**

1958: Well history report, Richfield-Mic Mac-Homestead Tow Hill No. 1; British Columbia Department of Mines and Petroleum Resources, Open File WA00380.

1959a: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 899.

1959b: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 939.

1962a: Well history report, Richfield Pure Abbotsford 16-17-16; British Columbia Department of Mines and Petroleum Resources, Open File DA00884.

1962b: Well history report, Richfield Pure Sunnyside 16-13; British Columbia Department of Mines and Petroleum Resources, Open File WA00913.

1963: Well history report, Richfield Pure Point Roberts 6-3-5; British Columbia Department of Mines and Petroleum Resources, Open File WA01075.

**Risley, D.E., Martin, G.C., Lynch, M.B., Flett, T.O., Larson, J.A., and Horowitz, W.L.**

1992: Geologic report for the Gulf of Alaska Planning Area, (ed.) R.F. Turner; U.S. Department of the Interior, OCS Report MMS 92-0065.

**Rohr, K.M.M. and Currie, L.**

1997: Queen Charlotte basin and Coast Mountains: Paired belts of subsidence and uplift caused by a low-angle normal fault; *Geology*, v. 25, p. 819–822.

**Rohr, K. and Dietrich, J.R.**

1990: Deep seismic survey of Queen Charlotte Basin; Geological Survey of Canada, Open File 2258.

1991: Deep seismic reflection survey of the Queen Charlotte Basin, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 127–133.

1992: Strike-slip tectonics and development of the Tertiary Queen Charlotte Basin, offshore western Canada: evidence from seismic reflection data; *Basin Research*, v. 4, p. 1–19.

**Royalite Oil Company**

1949: Well history report, Queen Charlotte No. 1; British Columbia Department of Mines and Petroleum Resources, Open File.

**Shell Canada Ltd.**

1968a: Well history report, Shell Anglo Tyee N-39; Energy, Mines, and Resources Canada, Open File.

1968b: Well history report, Shell Anglo Sockeye B-10; Energy, Mines, and Resources Canada, Open File.

1968c: Well history report, Shell Anglo Sockeye E-66; Energy, Mines, and Resources Canada, Open File.

1968d: Well history report, Shell Anglo Osprey D-36; Energy, Mines, and Resources Canada, Open File.

1968e: Well history report, Shell Anglo Auklet G-41; Energy, Mines, and Resources Canada, Open File.

1968f: Well history report, Shell Anglo Prometheus H-68; Energy, Mines, and Resources Canada, Open File.

1968g: Well history report, Shell Anglo Pluto I-87; Energy, Mines, and Resources Canada, Open File.

1968h: Well history report, Shell Anglo Zeus D-14; Energy, Mines, and Resources Canada, Open File.

1968i: Well history report, Shell Anglo Zeus I-65; Energy, Mines, and Resources Canada, Open File.

1969a: Well history report, Shell Anglo Harlequin D-86; Energy, Mines, and Resources Canada, Open File.

1969b: Well history report, Shell Anglo South Coho I-74; Energy, Mines, and Resources Canada, Open File.

1969c: Well history report, Shell Anglo Murrelet L-15; Energy, Mines, and Resources Canada, Open File.

1969d: Well history report, Shell Anglo Apollo J-14; Energy, Mines, and Resources Canada, Open File.

1969e: Well history report, Shell Anglo Cygnet J-100; Energy, Mines, and Resources Canada, Open File.

**Shouldice, D.H.**

1971: Geology of the western Canadian continental shelf; *Bulletin of Canadian Petroleum Geology*, v. 19, no. 2, p. 405–436.

**Sinclair, I.K., McAlpine, K.D., Sherwin, D.F., and McMillan, N.J.**

1992: Petroleum resources of the Jeanne D'Arc Basin and environs; Part I: Geological Framework; Geological Survey of Canada, Paper 92-8, p. 1–38.

**Singh, S.C., Minshull, T.A., and Spence, G.D.**

1990: Velocity structure of a gas hydrate reflector; *Science*, v. 260, p. 204–207.

**Snavely, P.D.**

1987: Tertiary geologic framework, neotectonics, and petroleum potential of the Oregon-Washington continental margin; *in* Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins – Beaufort Sea to Baja California, (ed.) D.W. Scholl, A. Grantz, and J.G. Vedder; Circum-Pacific Council for Energy and Mineral Resources, Earth Science Series, v. 6, p. 305–336.

**Snowdon, L.R., Fowler, M.G., and Hamilton, T.S.**

1988: Sources and seeps: organic geochemical results from the Queen Charlotte Islands; *in* Some Aspects of the Petroleum Geology of the Queen Charlotte Islands; Canadian Society of Petroleum Geologists, Field Trip Guide, p. 37–43.

**Spence, G.D., Clowes, R.M., and Ellis, R.M.**

1985: Seismic structure across the active subduction zone of western Canada; *Journal of Geophysical Research*, v. 90, p. 6754–6772.

- Spence, G.D., Hole, J.A., Asudeh, I., Ellis, R.M., Clowes, R.M., Yuan, T., and Rohr, K.M.M.**  
 1991: A seismic refraction study in the Queen Charlotte Basin, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 135–149.
- Stacey, R.A.**  
 1975: Structure of the Queen Charlotte Basin. *In* Canada's Continental Margins and Offshore Petroleum Exploration, C.J. Yorath, E.R. Parker, and D.J. Glass (eds.). Canadian Society of Petroleum Geologists, Memoir 4, p. 723–741.
- Sutherland Brown, A.**  
 1968: Geology of the Queen Charlotte Islands, British Columbia; British Columbia Department of Energy, Mines, and Petroleum Resources, Bulletin 54, 226 p.
- Sutherland Brown, A. and Yorath, C.J.**  
 1985: LITHOPROBE profile across southern Vancouver Island: geology and tectonics; *in* Field Guides to Geology and Mineral Deposits in the Southern Canadian Cordillera; Geological Society of America, Cordilleran Section Meeting, Vancouver, British Columbia, p. 8–1 to 8–23.
- Texaco Exploration Canada**  
 1968: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 1474.  
 1969: Assessment report; British Columbia Department of Mines and Petroleum Resources, Report 1530.
- Thompson, R.I., Haggart, J.W., and Lewis, P.D.**  
 1991: Late Triassic through early Tertiary evolution of the Queen Charlotte Basin, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 3–29.
- Tipper, H.W., Woodsworth, G.J., and Gabrielse, H.**  
 1981: Tectonic assemblage map of the Canadian Cordillera; Geological Survey of Canada, Map 1505A.
- Union Oil Company of Canada Ltd.**  
 1971: Well history report, Union Port Louis; British Columbia Department of Mines and Petroleum Resources, Open File WA02954.
- Vance, J.A.**  
 1975: Bedrock geology of San Juan County; *in* Geology and water resources of the San Juan Islands, (ed.) R.H. Russel; Washington Department of Ecology, Water-Supply Bulletin 46, p. 3-19.
- van der Heyden, P.**  
 1992: A middle Jurassic to early Tertiary Andean-Sierran arc model for the Coast Belt of British Columbia; *Tectonics*, v. 17, p. 163–174.
- Vellutini, D. and Bustin, R.M.**  
 1991a: Source rock potential of Mesozoic and Tertiary strata of the Queen Charlotte Islands; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 381–409.
- 1991b: Organic maturation and source rock potential of Mesozoic and Tertiary strata, Queen Charlotte Islands, British Columbia; *in* Evolution and Hydrocarbon Potential of the Queen Charlotte Basin, British Columbia, (ed.) G.J. Woodsworth; Geological Survey of Canada, Paper 90-10, p. 411–452.
- Wade, J.A., Campbell, G.R., Procter, R.M., and Taylor, G.C.**  
 1989: Petroleum resources of the Scotian Shelf; Geological Survey of Canada, Paper 88-19, 26 p.
- Wheeler, J.O. and McFeeley, P.**  
 1991: Tectonic assemblage map of the Canadian Cordillera and adjacent parts of the United States of America; Geological Survey of Canada, Map 1712A, scale 1:2,000,000.
- Wheeler, J.O., Brookfield, A.J., Gabrielse, H., Monger, J.W.H., Tipper, H.W., and Woodsworth, G.J.**  
 1991: Terrane map of the Canadian Cordillera; Geological Survey of Canada, Map 1713A, scale 1:2,000,000.
- White, D.A. and Gehman, H.M.**  
 1979: Methods of estimating oil and gas resources; American Association of Petroleum Geologists, Bulletin, v. 63, no. 12, p. 2183–2192.
- White, D.J. and Clowes, R.M.**  
 1984: Seismic investigation of the Coast Plutonic Complex – Insular Belt boundary beneath the Strait of Georgia; Canadian Journal of Earth Sciences, v. 21, p. 1033–1049.
- White, J.M.**  
 1991: Evidence of Paleogene sedimentation on Graham Island, Queen Charlotte Islands, west coast, Canada; Canadian Journal of Earth Sciences, v. 21, p. 1033–1049.
- White, W.H.**  
 1959: Cordilleran tectonics in British Columbia; American Association of Petroleum Geologists, Bulletin, v. 43, p. 60-100.
- Woodsworth, G.J.**  
 1988: Karmutsen Formation and the east boundary of Wrangellia, Queen Charlotte Basin, British Columbia; *in* Current Research, Part E; Geological Survey of Canada, Paper 88-1E, p. 209–212.
- Yorath, C.J.**  
 1980: The Apollo structure in the Tofino Basin, Canadian Pacific continental shelf; Canadian Journal of Earth Sciences, v. 17, p. 758–775.  
 1987: Petroleum geology of the Canadian Pacific continental margin; *in* Geology and Resource Potential of the Continental Margin of Western North America and Adjacent Ocean Basins – Beaufort Sea to Baja California, (ed.) D.W. Scholl, A. Grantz, and J.G. Vedder; Circum-Pacific Council for Energy and Mineral Resources, Earth Science Series, v. 6, p. 283–304.  
 1991: Upper Jurassic to Paleogene assemblages; *in* Chapter 9 of Geology of the Cordilleran Orogen in Canada, (ed.) H. Gabrielse and C.J. Yorath; Geological Survey of Canada, Geology of Canada, no. 4, p. 329–371 (*also* Geological Society of America, The Geology of North America, v. G-2, p. 329–371).
- Yorath, C.J. and Chase, R.L.**  
 1981: Tectonic history of the Queen Charlotte Islands and adjacent areas – a model; Canadian Journal of Earth Sciences, v. 20, p. 135–159.

**Yorath, C.J. and Hyndman, R.D.**

1983: Subsidence and thermal history of Queen Charlotte Basin; Canadian Journal of Earth Sciences, v. 20, p. 135-159.

**Yorath, C.J., Clowes, R.M., Sutherland Brown, A., Brandon, M.T., Massey, N.W.D., Green A.G., Spencer, C., Kanasewich, E.R., and Hyndman R.D.**

1985a: LITHOPROBE – Phase I: Southern Vancouver Island: preliminary analyses of reflection seismic profiles and surface geological studies; *in* Current Research, Part A; Geological Survey of Canada, Paper 85-1A, p. 543–554.

**Yorath, C.J., Green, A.G., Clowes, R.M., Sutherland Brown, A., Brandon, M.T., Kanasewich, E.R., and Spencer, C.**

1985b: LITHOPROBE, southern Vancouver Island: seismic reflection sees through the Juan de Fuca Plate; *Geology*, v. 13, p. 759-762.

**Yorath, C.J., Sutherland Brown, A., Campbell, R.B., and Dodds, C.J.**

1991: The Insular Belt; *in* Chapter 9 of Geology of the Cordilleran Orogen in Canada, (ed.) H. Gabrielse and C.J. Yorath; Geological Survey of Canada, Geology of Canada, no. 4, p. 354–360 (*also* Geological Society of America, The Geology of North America, v. G-2, p. 354–360).

## APPENDIX A - INPUT DATA FOR PETROLEUM ASSESSMENTS

The following tables present the probability distributions of reservoir parameters, number of prospects, and marginal probabilities of geological risk factors used as input for various conceptual statistical analyses discussed in this paper. These estimates are based on subjective opinion, partly constrained by reservoir data and information from analogous petroleum-bearing basins.

### 1. QUEEN CHARLOTTE CRETACEOUS OIL PLAY

**Table A-1a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	2	50	100
Formation thickness	m	1	30	180	200
Porosity	decimal fraction	0.08	0.15	0.2	0.25
Trap fill	decimal fraction	0.1	0.3	0.8	1.0
Oil saturation	decimal fraction	0.5	0.65	0.75	0.8
Shrinkage factor	decimal fraction	0.65	0.80	0.9	0.95

**Table A-1b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		X
Presence of reservoir facies	0.5		X
Adequate seal	0.8		X
Adequate timing	0.9		X
Adequate source	0.5		X
Adequate recovery	0.9		X
Adequate play conditions	0.95	X	

**Table A-1c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	300	500	1000

## 2. QUEEN CHARLOTTE CRETACEOUS GAS PLAY

**Table A-2a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	2	50	100
Formation thickness	m	1	30	180	200
Porosity	decimal fraction	0.08	0.15	0.2	0.25
Trap fill	decimal fraction	0.05	0.2	0.7	0.9
Gas saturation	decimal fraction	0.5	0.65	0.75	0.8
Formation volume factor	decimal fraction	0.002	0.004	0.009	0.01

**Table A-2b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		X
Presence of reservoir facies	0.5		X
Adequate seal	0.8		X
Adequate timing	0.9		X
Adequate source	0.4		X
Adequate recovery	0.9		X
Adequate play conditions	0.95	X	

**Table A-2c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	300	500	1000

### 3. QUEEN CHARLOTTE MIOCENE OIL PLAY

**Table A-3a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	3	100	300
Formation thickness	m	100	300	800	1000
Porosity	decimal fraction	0.1	0.25	0.3	0.35
Trap fill	decimal fraction	0.01	0.05	0.2	0.25
Oil saturation	decimal fraction	0.5	0.65	0.75	0.8
Shrinkage factor	decimal fraction	0.65	0.80	0.9	0.95

**Table A-3b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		X
Presence of reservoir facies	0.9		X
Adequate seal	0.9		X
Adequate timing	0.8		X
Adequate source	0.4		X
Adequate preservation	0.7		X
Adequate recovery	0.9		X
Adequate play conditions	0.9	X	

**Table A-3c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	90	300	400



#### 4. QUEEN CHARLOTTE MIOCENE GAS PLAY

**Table A-4a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	3	100	300
Formation thickness	m	100	300	800	1000
Porosity	decimal fraction	0.1	0.25	0.3	0.35
Trap fill	decimal fraction	0.01	0.05	0.2	0.25
Gas saturation	decimal fraction	0.5	0.65	0.75	0.8
Formation volume factor	decimal fraction	0.002	0.004	0.009	0.01

**Table A-4b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		X
Presence of reservoir facies	0.9		X
Adequate seal	0.9		X
Adequate timing	0.8		X
Adequate source	0.5		X
Adequate preservation	0.8		X
Adequate recovery	0.9		X
Adequate play conditions	0.9	X	

**Table A-4c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	90	300	400

## 5. QUEEN CHARLOTTE PLOIOCENE OIL PLAY

**Table A-5a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	5	150	300
Formation thickness	m	200	700	1200	1500
Porosity	decimal fraction	0.15	0.28	0.35	0.38
Trap fill	decimal fraction	0.01	0.02	0.2	0.25
Oil saturation	decimal fraction	0.5	0.65	0.75	0.9
Shrinkage factor	decimal fraction	0.65	0.80	0.9	0.95

**Table A-5b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.8		X
Adequate seal	0.7		X
Adequate timing	0.5		X
Adequate source	0.4		X
Adequate recovery	0.9		X
Adequate play conditions	0.7	X	

**Table A-5c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	40	200	300

## 6. QUEEN CHARLOTTE PLIOCENE GAS PLAY

**Table A-6a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	5	150	300
Formation thickness	m	200	700	1200	1500
Porosity	decimal fraction	0.15	0.28	0.35	0.38
Trap fill	decimal fraction	0.01	0.02	0.2	0.25
Gas saturation	decimal fraction	0.5	0.65	0.75	0.8
Formation volume factor	decimal fraction	0.0025	0.005	0.009	0.01

**Table A-6b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.8		X
Adequate seal	0.7		X
Adequate timing	0.6		X
Adequate source	0.6		X
Adequate recovery	0.9		X
Adequate play conditions	0.9	X	

**Table A-6c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	40	200	300

## 7. GEORGIA CRETACEOUS STRUCTURAL GAS PLAY

**Table A-7a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	10	50	60
Formation thickness	m	10	60	200	250
Porosity	decimal fraction	0.03	0.05	0.12	0.15
Trap fill	decimal fraction	0.01	0.2	0.5	1.00
Gas saturation	decimal fraction	0.5	0.75	0.9	0.95
Formation volume factor	decimal fraction	0.002407	0.004211	0.019051	0.02

**Table A-7b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.75		X
Presence of porosity	0.2		X
Adequate seal	0.95		X
Adequate source	0.9		X
Adequate play conditions	0.75	X	

**Table A-7c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	60	450	900

## 8. GEORGIA TERTIARY STRUCTURAL GAS PLAY

**Table A-8a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	10	50	60
Formation thickness	m	1	15	20	30
Porosity	decimal fraction	0.08	0.15	0.34	0.37
Trap fill	decimal fraction	0.0	0.1	0.2	1.00
Water saturation	decimal fraction	0.1	0.1	0.1	0.1
Formation volume factor	decimal fraction	0.002	0.004	0.019	0.02

**Table A-8b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.75		X
Presence of porosity	0.9		X
Adequate seal	0.85		X
Adequate play conditions	0.98	X	

**Table A-8c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	60	150	300

## 9. GEORGIA PLEISTOCENE STRATIGRAPHIC GAS PLAY

**Table A-9a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	0.1	0.3	0.5	1.5
Formation thickness	m	1	11	20	30
Porosity	decimal fraction	0.1	0.25	0.32	0.37
Trap fill	decimal fraction	0.5	0.75	0.9	1.00
Gas saturation	decimal fraction	0.01	0.25	0.9	1.0
Formation volume factor	decimal fraction	0.05	0.07	0.1	0.5

**Table A-9b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.9		X
Presence of porosity	0.9		X
Adequate seal	0.5		X

**Table A-9c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	10	200	500

## 10. TOFINO TERTIARY STRUCTURAL GAS PLAY

**Table A-10a**  
Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile		1.0	0.5	0.01	0.0
Area of closure	km <sup>2</sup>	1	25	145	200
Formation thickness	m	20	200	500	1100
Porosity	decimal fraction	0.07	0.15	0.21	0.32
Trap fill	decimal fraction	0.01	0.05	0.1	0.15
Gas saturation	decimal fraction	0.5	0.8	0.9	0.95
Formation volume factor	decimal fraction	0.002	0.006	0.011	0.012

**Table A-10b**  
Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.6		X
Presence of reservoir facies	0.6		X
Presence of porosity	0.9		X
Adequate seal	0.9		X
Adequate source	0.3		X

**Table A-10c**  
Probability distribution for number of prospects

Geological variable	Probability in upper percentiles	Probability in upper percentiles	Probability in upper percentiles
Percentile	0.99	0.5	0.0
Number of prospects	100	500	800

