Natural Resources Canada | Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA

DESCRIPTIVE NOTES

INTRODUCTION

The Manitouwadge greenstone belt comprises a highly deformed remnant of upper amphibolite-facies supracrustal rocks adjacent to the boundary between the Wawa and Quetico subprovinces of the Archean Superior Province (Fig. 2). These 1:10 000 (Fig. 1) and 1:25 000 (Map 1917A) maps of the Manitouwadge greenstone belt are the result of new mapping with supporting structural studies, petrology, geochemistry, and geochronology, funded by the Canada–Ontario Northern Ontario Development Agreement (NODA) from 1991–1996.

The Manitouwadge greenstone belt is host to major volcanogenic Cu-Zn-Ag deposits which produced 63.9 Mt of ore collectively. Production came to an end with the closure of the Geco mine (Noranda Mineral Inc., Geco Division) in 1995. The known economic deposits were discovered in the 1950s on the basis of surface exposures and conventional geophysical methods that detected orebodies under thin overburden. The belt is still highly prospective terrain, but further exploration will benefit from an improved understanding of the primary depositional setting and its subsequent modification. The 1:10 000 map, focused on the area of known economic mineralization, is based on detailed new mapping supplemented by compilation of existing published and unpublished maps and, in some cases, subsurface information from drill holes and underground workings. The 1:25 000 map shows the regional setting and structure of the greenstone belt, and its relationship to the Wawa–Quetico subprovince boundary.

The Manitouwadge greenstone belt lies in the volcano-plutonic Wawa Subprovince, immediately south of the boundary with the migmatitic metasedimentary Quetico Subprovince (Fig. 2). The Superior Province was initially subdivided into subprovinces on the basis of structural trends and style, for example, contrasting the linear east-west trends of the Quetico Subprovince with the curvilinear trends of the Wawa and Wabigoon subprovinces, as defined by remnants of greenstone belts lying between batholithic domes (Stockwell, 1964). Lithological, metamorphic, and geophysical characteristics were subsequently also used as distinguishing features (Stockwell, 1970; Card and Ciesielski, 1986). The tectonic significance of the subprovinces and their belt-like configuration has stimulated much debate. Langford and Morin (1976) proposed assembly of the Superior Province by successive accretion of island arcs and inter-arc sedimentary basins, based on comparison with the belt structure of the Canadian Cordillera. More recent refinements of arc-accretionary models have been proposed for the Wabigoon, Quetico, and Wawa subprovinces by Percival (1989), Percival and Williams (1989), and Williams (1990), among others. Arc-accretionary models commonly describe the Quetico Subprovince as an accretionary complex that accumulated southward from the margin of the Wabigoon arc above a northerly dipping subduction zone, with collision of the Wawa arc from the south amalgamating the three subprovinces.

The Manitouwadge project illustrates the importance of detailed study in constraining regional tectonic models. The geological evolution of the Manitouwadge greenstone belt is related to its proximity to the Wawa–Quetico subprovince boundary, and the end result is a complex geometry from which the structural evolution and primary stratigraphic relationships need to be unravelled.

REGIONAL SETTING

Together with the Fariles Lake and Moshkinabi belts to the east (Williams and Breaks, 1990a, 1996), the Manitouwadge belt is part of a semicontinuous supracrustal suite (Manitouwadge–Hornpayne assemblage, Williams et al., 1991) along the northern margin of the Wawa Subprovince. The greenstone belts are deformed by east-northeasterly trending folds, including the Jim Lake synform, Blackman Lake antiform, and Manitouwadge synform (Map 1917A). Gradients in deformation state and metamorphic grade are in part related to the synformic boundary. The folds tighten toward the north, and their axial traces curve, becoming transitional to the steeply dipping east-west trends of the Quetico Subprovince. The regional metamorphic grade increases from upper amphibolite facies in the Manitouwadge belt toward a zone of orthopyroxene-bearing granulite rocks in the Quetico Subprovince (Williams and Breaks, 1990a, b; Pan et al., 1994). All supracrustal rocks in the map area have been regionally metamorphosed and are generally characterized by upper amphibolite to granulite-facies assemblages.

The Manitouwadge greenstone belt is bounded to the west and south by foliated multiphase plutonic rocks of the Black Pic batholith which extend south to the Schreiber–Hemlo greenstone belt (Williams et al., 1991). Between the Manitouwadge and Hemlo–Schreiber belts, foliations in the batholith define a structural dome (Williams and Breaks, 1990a, b).

The regional map pattern is dominated by F_2 folds, including the Manitouwadge synform which plunges to the east at about 25°. The supracrustal rocks are thickest in the hinge region of the Manitouwadge synform and along its southern limb, the area shown in detail on Figure 1. Away from the hinge region, supracrustal rocks are increasingly attenuated and invaded by intrusions.

PREVIOUS MAPPING

The first systematic mapping of the Manitouwadge greenstone belt was undertaken by Pye (1957) for the Ontario Department of Mines. He outlined the general shape of the belt and recognized mafic metavolcanic rocks along the inner and outer margins of the folded supracrustal sequence. At the time of Pye's mapping, most of the presently known orebodies had already been discovered and, in many cases, his detailed maps of mine properties are the only published record of geological relationships since obliterated by mining. Pye regarded the mineralization as Algoman replacement that postdated all intrusive activity, with the possible exception of diabase dykes. He interpreted most supracrustal rocks as metasedimentary. Milne (1974) remapped the area of major economic interest and reinterpreted the felsic rocks to be of volcanic, volcaniclastic, and sedimentary origin. Regional mapping by Williams and Breaks (1990a, 1996) provided a regional context and structural interpretation for the Manitouwadge belt and the Wawa–Quetico boundary region. The trace of the orthopyroxene-in isograd, and the locations of several iron-formation layers in the Quetico Subprovince (Map 1917A) are partly compiled from their work.

In addition to published maps, detailed geological mapping and geophysical surveys of claim groups have been undertaken by private-sector companies and prospectors. Unpublished claim maps were made available by Noranda Mining and Exploration Inc., Minnova Inc., and Breakwater Resources Ltd., during the course of this mapping. The area near Gaug and Agam lakes (Fig. 1) was in part compiled from detailed mapping by Keast (1991) and Bates (1993) for Grange Inc.

TECTONOSTRATIGRAPHY AND GEOCHRONOLOGY

Manitouwadge greenstone belt

The stratigraphy of the Manitouwadge greenstone belt was defined in the hinge region and along the southern limb of the F_2 Manitouwadge synform in the thickest preserved section of supracrustal rocks. Deformation and metamorphic recrystallization have resulted in a paucity of way-up indications on outcrop scale. However, in the area of Willroy (W1–W6) and Geco (G1–G3) mines, Zn-Cu ratios of orebodies and the position of footwall hydrothermal alteration zones and of a subvolcanic trondhjemite (see below) indicate southward overgoing. The stratigraphic sequence in this area dips subvertically, and shows a progression from mafic volcanic rocks at the base (north), overlain by interlayered felsic volcanic rocks and iron-formation, overlain by metagreywacke (south). This sequence is repeated along the southern limb of the Manitouwadge synform by the F_2 Agam Lake syncline. Two belts of volcanic rocks, hereafter referred to as "inner" and "outer" with respect to the easterly plunging F_2 fold, show a progression from mafic to felsic rocks toward the greywacke (see Fig. 3–9 for representative illustrations of the lithological units). Whereas economic mineralization is known only in the inner volcanic belt, felsic rocks are interlayered with iron-formation in both. The synvolcanic trondhjemite (unit Ato) which intrudes the base of the sequence, in the present configuration occupies a structural position in the core of the F_2 Manitouwadge synform. In the inner belt, three discrete mappable bodies of coarsely quartz-porphyritic felsic rocks (unit Ato) are interlayered with iron-formation (units Aip, Aie, Aib) and spatially associated with massive sulphide deposits. In the outer volcanic belt, similar felsic rocks (unit Ato) are interlayered with iron-formation. Mafic rocks (units Am, Amg, Amv) are more abundant in the outer volcanic belt than in the inner, occurring in the latter as screens or septa in synvolcanic trondhjemite and interleaved with felsic metavolcanic rocks along the transitional contact zone.

The Dead Lake suite (unit Ad), an enigmatic package of interleaved foliated gabbro, diorite, and layered mafic to intermediate rocks of unknown origin, forms several semicontinuous zones in the inner trondhjemite. The Dead Lake suite includes a distinctive group of strongly magnetic, melanocratic rocks characterized by hornblende-magnetite ± plagioclase ± garnet ± clinopyroxene ± syngabbro minerals (Fig. 10). The melanocratic rocks show local transitions on centimetre- to outcrop-scale to mesocratic, garnet ± magnetite ± hornblende trondhjemite, which is transitional to typical leucocratic trondhjemite (Zaleski et al., 1995b). Similar rocks are present in sporadic exposures in the mafic volcanic unit near Gaug Lake in the outer volcanic belt, suggesting that the Dead Lake suite is also repeated by the Agam Lake synform.

Uranium-lead zircon geochronology on felsic volcanic and subvolcanic rocks supports the correlation of volcanic units across the Agam Lake syncline. Aphyric felsic breccia from the outer volcanic belt gave an age of 2722 ± 2 Ma (Zaleski et al., 1994, 1995), within error of 2720 ± 2 Ma muscovite schist at the Geco mine interpreted as an altered felsic volcanic rock (Davis et al., 1994). Foliated trondhjemite sampled north of the Willroy and Geco mines gave an age of 2720 ± 2 Ma, confirming our field interpretation of subvolcanic intrusion.

Extensive zones of orthoamphibole-cordierite-garnet gneiss (unit Aog) near the mafic-felsic transition in the inner volcanic belt, and to a lesser extent in the outer volcanic belt, are interpreted as synvolcanic hydrothermally altered, alkali-depleted, Fe-Mg-enriched rocks, modified by high-grade metamorphic recrystallization and deformation (Fig. 11). The precursors to alteration, mafic and interlayered mafic-felsic rocks near the transitional mafic-felsic contact, are still recognizable in less altered enclaves mostly on the margins of the unit and to the northwest further from known deposits. Near the contact with orthoamphibole-cordierite-garnet gneiss, trondhjemite locally contains disseminated garnet and foliated seams of orthoamphibole, garnet, and biotite, suggesting that the trondhjemite was also affected by incipient hydrothermal alteration. On the basis of these observations, the trondhjemite was interpreted as part of a dynamic hydrothermal system in which the intrusion provided heat for driving hydrothermal circulation, and continued to rise intrusting its own halo of hydrothermally altered rocks (Zaleski and Peterson, 1995). Surface mapping and subsurface data show that orthoamphibole-garnet rocks form a semiconformable sheet of regional extent, mantling synvolcanic trondhjemite and folded by the F_2 Manitouwadge synform. Orthoamphibole-garnet gneiss extends semicontinuously for at least 30 km from Rabbitskin Lake on the northern limb of the Manitouwadge synform, around the hinge of the fold to the Willroy–Geco area, east to the Hucamp zone of subeconomic mineralization (H17), and beyond to the Gavan mines (H18) and Falconbridge zones (H19). In the outer volcanic belt, orthoamphibole-garnet assemblages are found locally in a discontinuous zone in mafic metavolcanic rocks near the northern contact with felsic metavolcanic rocks and metagreywacke.

The orthoamphibole-cordierite-garnet rocks, equivalent to synvolcanic footwall alteration zones associated with volcanogenic massive sulphide deposits in less metamorphosed terranes, lie north of the Geco mine (NC), Willroy (W1–W6), and Geco (G1–G3) deposits. The Geco orebodies increase in Zn/Cu from north to south, characteristically a stratigraphic-upwards zonation in volcanogenic massive sulphide deposits (Sutfel et al., 1971). The position of the footwall alteration with respect to the deposits and the southward increase in Zn/Cu have been used to infer southward youngingin the Willroy–Geco area (Sutfel et al., 1971; Friesen et al., 1982). The northern position of the synvolcanic trondhjemite intrusion is consistent with exposure of progressively deeper levels of a volcano-plutonic edifice in that direction in the inner volcanic belt.

Quartz-muscovite ± sillimanite schist, sillimanite-knot felsic schist, and quartzose schist (units Aam, Aas, Aaq) are interpreted as the products of aluminous-potassic and silicic synvolcanic hydrothermal alteration (Fig. 12; Zaleski and Peterson, 1995). Sillimanite-muscovite-quartz schist is found close to massive sulphide deposits, mainly to the south of orebodies in the Willroy–Geco area and enveloping the Geco main orebody (S1). Two zones of sillimanite-bearing felsic schist converge to the east near the Geco mine, where they are interpreted to be separated by an early, low-angle fault (Sunset Lake fault, see below). The northern zone grades easterly from sillimanite-knot schist to sillimanite-muscovite-quartz schist consisting of alternating thin quartz-rich and muscovite-rich layers with sillimanite forming sprays on the foliation surface. Near Willroy, the southern belt consists of thinly layered quartzofeldspathic schist that becomes more muscovitic and silicic to the east, merging with the northern belt and grading to a sulphidic muscovite-biotite quartzose schist in the Geco area. Felsic volcanic rocks lying along strike and in less altered enclaves are interpreted as the protolith of the altered rocks.

Near the hinge region of the Manitouwadge synform in the Willecho area (E1–E3), sillimanite-knot felsic schist is interleaved (on a metre scale) with nonsillimanitic felsic schist, straight gneiss, and sheared pegmatite. Similar rocks occur sporadically to the south of Willecho, associated with iron-formation that can be traced continuously to the Willroy area. At a few localities, thinly layered quartz-porphyritic felsic tuff or tuffaceous metasediment was found to contain abundant sillimanite that coalesces into sillimanitic layers.

Three types of iron-formation (units Aib, Aia, Aig) are present in the Manitouwadge belt: a) quartz-magnetite-silicate banded iron-formation, b) magnetite-silicate iron-formation, and c) sulphidic iron-formation consisting of disseminated or stockwork pyrite and/or pyrrhotite in quartz-magnetite-silicate host rocks. Quartz-magnetite-silicate banded iron-formation is by far the most abundant type (Fig. 7, 8). Magnetite-silicate iron-formation is mainly present near the contacts of banded iron-formation, especially near the northern contact (the stratigraphic footwall) of the thickest iron-formation. Sulphidic iron-formation rarely forms mappable units, but occurs mainly in the transition from quartz-magnetite-silicate banded iron-formation to massive sulphide (e.g. Timms and Marshall, 1955).

Monotonous metagreywacke and biotite-hornblende schist (Fig. 9a) lie between the inner and outer metavolcanic belts, occupying the centre of the F_2 Agam Lake syncline along the southern limb of the F_2 Manitouwadge synform. The maximum age of sedimentation of 2693 Ma, constrained by U-Pb dating of detrital zircon grains, is at least 25 Ma younger than the age of felsic volcanism associated with Cu-Zn mineralization (Zaleski et al., 1996). The nature of the contact with adjacent metavolcanic rocks is obscured by intrusions, metamorphic recrystallization, and deformation, but it is either an unconformity or an early (pre- to syn-D_2) fault.

Very fine-grained to aphanitic, laminated felsic rocks, referred to as straight gneiss (Fig. 13), were mapped as semicontinuous outcrop-scale layers and lenses. In the Willroy–Geco area, straight gneiss lies in a zone along which iron-formation and other units are truncated and, across which the sequence quartz-porphyritic felsic rock, iron-formation (with massive sulphide mineralization), sillimanite-muscovite-quartz schist, is repeated. Based on these observations, straight gneiss is interpreted as annealed mylonite marking early ductile faults (see below). The protolith to straight gneiss is mainly felsic volcanic rocks and, in some cases, pegmatite.

Away from the hinge region of the Manitouwadge synform, supracrustal rocks are increasingly attenuated and invaded by intrusions, and unambiguous identification of protolith becomes difficult. Undivided mafic to intermediate rocks, locally with garnetiferous zones, define the northern limb of the Manitouwadge synform and related macroscopic folds to the north. Orthoamphibole-bearing rocks, well exposed west of Rabbitskin Lake, trend easterly into the east-northeasterly trending Rabbitskin Lake fault. Further east along the the northern limb of the Manitouwadge synform, orthoamphibole-garnet rocks outcrop west of the hinge region of the Blackman Lake antiform associated with sphalerite-bearing iron-formation at the Jim Lake occurrence (A51). In the vicinity of the Blackman Lake and Jim Lake folds, the greenstone belt is represented by layered metavolcanic rocks with minor iron-formation, and by foliated medium- to coarse-grained amphibolite that could be recrystallized volcanic, subvolcanic, or plutonic rocks. The Jim Lake synform is partly defined by zones of mafic to intermediate inclusions in foliated tonalite and pegmatite. Inside the Manitouwadge synform, the Dead Lake suite forms a marker unit that can be traced in outcrop and as an aeromagnetic high (Geological Survey of Canada, 1997) along the northern limb of the synform and around the Blackman Lake antiform.

On the southern limb of the Manitouwadge synform, the easternmost exposure of orthoamphibole-bearing rocks lies about 4 km east of Wiwun Lake. To the east, exposure is poor; however, a prominent aeromagnetic anomaly continues the same trend to the Falconbridge zone (H19), and the unit was intersected by drilling (unpub. data, Noranda Mines Ltd., Geco Division, 1984). The outer belt of mafic metavolcanic rocks can be traced as a southerly diverging zone of inclusions in foliated plutonic rocks. South of Thompson Lake, the area of the Banana Lake antiform is poorly exposed and no supracrustal rocks outcrop. Foliations in plutonic rocks are folded, and supracrustal rocks are exposed to the south in the Fariles Lake belt (Williams and Breaks, 1990a, 1996). East of Thompson and Banana lakes near the eastern extremity of the map area, orthoamphibole-garnet-cordierite gneiss, mafic schist, and garnet ± magnetite ± hornblende ± clinopyroxene rocks resembling the Dead Lake suite define macroscopic folds with a northerly trending enveloping surface.

QUETICO SUBPROVINCE

Metagreywacke in the Quetico Subprovince is indistinguishable from that in Manitouwadge greenstone belt, except for the difference in metamorphic grade, as shown by pervasive migmatitic segregations in the Quetico metagreywacke (Fig. 9b). Locally, thin siliceous and sulphide iron-formation, ranging from inches to more than 10 m in width (e.g. McKay, 1994), are intercalated with migmatitic metagreywacke. Iron-formation widths as shown on the map are exaggerated for legibility. In the area of the Jim Lake synform, there is a compositional gradation from hornblende-dominated schist, interpreted to be of volcanic or volcaniclastic origin, to hornblende-biotite metagreywacke. The depositional age of Quetico greywacke, constrained by U-Pb dating of detrital zircon grains, is less than 2690 Ma, permitting correlation of Quetico greywacke with greywacke on the southern limb of the Manitouwadge synform in the Wawa Subprovince (Zaleski et al., 1996).

STRUCTURE AND METAMORPHISM

The structural evolution can be modelled by four phases of ductile deformation (D_1-D_4). The deformation history is based on field observations of planar (S) and linear (L) fabric elements, their inter-relationships, and their relationships with respect to mesoscopic folds (F_1), coupled with map-scale interpretations. Our deformation sequence is generally compatible with that of Williams and Breaks (1989, 1990a, 1996), with differences arising from our detailed mapping of supracrustal rocks and extensive integration of geological and aeromagnetic data (Geological Survey of Canada, 1997).

The oldest fabric elements recognized include modified bedding, contacts, volcanic flow horizons, and transposed sheets or layering in plutonic rocks and in metasomatically altered rocks. These are subparallel to tectonic fabric elements, that is, subparallel to laminations in straight gneiss (annealed mylonitic layering, S_1), to the dominant mineral foliation (S_2), or to the oldest migmatitic segregations (S_2) in the Quetico Subprovince. The dominant foliation (S_2) is typically a composite fabric comprising transposed layering and laminations (S_0-S_1), and layer-parallel mineral schistosity (S_2). The schistosity (S_2), and a strong lineation (L_2) on the schistosity surface, are defined by the preferred crystallographic orientation and shapes of high-grade metamorphic minerals including mica, amphibole, cordierite, and sillimanite. Mineral lineations (L_2) are commonly parallel to a stretching lineation defined by rodding of mineral aggregates and elongate sillimanite knots, and prolate primary features such as clasts, pillows, and phenocrysts.

The dominant S_2 planar and L_2 linear fabrics are widely distributed and their relationships to mesoscopic and macroscopic folds were used to classify fold generations. In general, F_2 folds deform compositional layering and modified bedding, and the dominant foliation and lineation are parallel to F_2 axial planes and fold axes, respectively. Key relationships supporting an episode of D_1 deformation are preserved locally in the hinge regions of F_2 folds where folded mylonitic layering (S_1) is oblique to S_2 axial planar fabrics. F_2 folds, including the Manitouwadge synform, deform the dominant D_2 fabrics and, in some cases, are associated with weakly developed crenulation microfolds (F_3) and crenulation cleavage (S_3). F_2 folds are typically kinks at high angle to the dominant foliation and, locally, are strongly developed in schistose rocks and associated with crenulation cleavage (S_3).

Metamorphic grade increases northward culminating in an orthopyroxene-granulite zone in the Quetico Subprovince (Williams and Breaks, 1990a). Thermobarometry in granulite assemblages indicates peak metamorphic conditions of 680–770°C and 4–6 kbar (Pan et al., 1994). In the southern Manitouwadge belt, thermobarometry has resulted in estimates of 600°C and 3–5 kbar (Pan and Fleet, 1992) and 650°C and 5–7 kbar (Petersen, 1984). The regional gradient is marked in the Manitouwadge belt by a change from muscovite-sillimanite-quartz schist on the southern limb of the Manitouwadge synform to sillimanite-microcline-quartz schist near the hinge region. In the absence of evidence for melting, this transition is consistent with the high temperatures and with low pressures of less than 4–5 kbar (e.g. Spear, 1993).

With increasing metamorphic grade toward the north, the dominant planar fabric is increasingly recrystallized and is commonly defined by weak mineral foliation and transposed compositional layering, which in the Quetico Subprovince, includes modified bedding and migmatitic segregations. Mineral and/or stretching lineations are subtle or absent. Especially in the Quetico Subprovince, mesoscopic folds (F_2) that deform early migmatitic layering (S_2) show development of axial planar mineral foliations (S_2). A hornblende ± biotite foliation (post-D_2 pre-D_3) transects the dominant fabrics at a high angle.

D_1 and D_2 are particularly important for understanding stratigraphic relationships in the area of known economic deposits, the geometry of the 'Geco mine horizon', and the internal complexity of the greenstone belt. In the Willroy–Geco area, the sequence, quartz-porphyritic felsic rocks, iron-formation (with massive sulphide deposits), sillimanite-muscovite-quartz schist, is repeated across an interpreted D_1 fault (Garnet Lake fault) marked by zones of straight gneiss. The Garnet Lake fault divides the Nama Creek–Willroy–Geco area into two blocks and results in repetition of mineralized horizons (Zaleski and Peterson, 1995; Peterson and Zaleski, 1996). The fault geometry and presence of mylonitic layering (annealed S_1) are consistent with a ductile fault with relatively minor displacement (hundreds of metres to 1 km) at low angle to stratigraphic contacts. The fault continues eastward in sillimanite-muscovite-quartz schist through the Geco area, to highly strained iron-formation and straight gneiss exposed near the Hucamp zone (H17) east of Wiwun Lake. Northwest of Willroy, the D_1 fault separates an area of complex folding, involving iron-formation and felsic rocks in the Willecho area and south, from an area of more continuous orthoamphibole-garnet gneiss and trondhjemite to the northeast. North of the Willecho deposits, quartz-porphyritic felsic rocks, iron-formation, and sillimanite-knot felsic schist are repeated by the F_2 fold that is transected by the dominant S_2 fabric and, hence, probably an F_1 fold.

D_3 deformation produced folds with curved hinge lines, including sheath folds, on map and outcrop scale (Peterson and Zaleski, 1994, 1999). The southernmost iron-formation in the inner volcanic belt is considerably thickened by F_3 folding and, in the Geco area, it is much more continuous than conformable tonalite dykes or sheets of probable D_2 age. In the complex area between Nama Creek and Willecho, a macroscopic F_3 fold repeats iron-formation and the D_1 Garnet Lake fault in a closed structure (approximately 4 km long and 1 km wide). The intrafolial, rather than repetitive, character of the structure and the nearly coaxial relationship of D_2 and D_3 lineations and mesoscopic fold axes are best explained by an F_3 sheath fold, rather than an F_2-F_3 interference pattern (Peterson and Zaleski, 1999). The Willecho deposits and subeconomic sulphide occurrences (#7 and #8) lie near or on the folded repetition of the fault.

On the southern limb of the Manitouwadge synform, the lithological sequence of mafic volcanic rocks, felsic volcanic rocks interlayered with iron-formation, sedimentary rocks, was interpreted as a stratigraphic succession repeated by an early syncline (Suffel et al., 1971; Robinson, 1979), here called the F_2 Agam Lake syncline. A possible consequence of F_3 fold repetition of sequences across the southern limb of the F_2 Manitouwadge synform is that the mineralized and altered intervals of the Willroy–Geco area may be repeated in the outer volcanic belt, assuming that they were not removed by erosion or faulting along the contact with younger greywacke. The sporadic zones of orthoamphibole, garnet, and minor cordierite in mafic rocks in the outer belt near the contact with interlayered felsic rocks and iron-formation are interpreted as synvolcanic alteration zones correlative with altered rocks in the Willroy–Geco area.

The dominant foliation and mineral lineations, defined by peak metamorphic minerals in the Manitouwadge belt and folded by F_2 folds on map and outcrop scale, are consistent with D_2 fabric development during peak metamorphism. Toward the north, evidence of recrystallization subsequent to D_2 deformation increases. In the Quetico Subprovince and near the subprovince boundary, migmatitic segregations are both folded by, and cut across, folds attributed to D_2 deformation. Ubiquitous mesoscopic Z-folds and dextral kinematic indicators support oblique dextral shear during D_2 deformation. Minor structures deform migmatitic layering and are transected by migmatitic segregations, suggesting that peak metamorphism and migmatization were associated with progressive deformation and the development of F_2 folds near the subprovince boundary. These observations imply a protracted metamorphic history, and diachronous cooling, with cooling from peak temperatures predating D_2 deformation in the Manitouwadge belt and postdating D_2 deformation in the Quetico Subprovince (Zaleski et al., 1995a). This interpretation is supported by U-Pb zircon ages on plutonic rocks. Strong D_2 fabrics in the 2687 ± 2 Ma Loken Lake pluton, folded by the Manitouwadge synform, are cut by 2680 ± 4/-3 Ma foliated granite, bracketing the D_2 fabrics and, by inference peak metamorphism, to the interval ca. 2687–2680 Ma (Zaleski et al., 1996). In the Wawa–Quetico boundary region, the 2679 ± 2 Ma Everest Lake pluton and the 2680 ± 2 Ma Nama Creek pluton show evidence of migmatization.

D_4 fabrics are well developed in synvolcanic trondhjemite, early phases of the Black Pic batholith and in the Loken Lake pluton. Macroscopic F_2 synformal 'keels', account for repetitions of supracrustal rocks about pluton-cored antiformal culminations. For example, our preferred interpretation of the mafic and orthoamphibole-cordierite-garnet rocks east of Thompson and Banana lakes involves repetition of altered rocks and trondhjemite by an F_2 synformal keel. The Loken Lake pluton, characterized by an L_4 or S_4 fabric, is interpreted to occupy an F_2 antiformal fold.

Easterly plunging F_2 folds, the Manitouwadge synform, Blackman Lake antiform, and Jim Lake synform, dominate the regional map pattern and involve migmatitic greywacke continuous with that of the Quetico Subprovince. The axial surface of the Manitouwadge synform dips to the south, resulting in shallow to moderate southerly dips of layering and foliation on the northern limb, and near-vertical to moderate southerly dips on the southern limb (Fig. 14). F_2 folds deform D_1 fabrics and, in general in the Manitouwadge belt, S_1 axial planar fabrics are weakly developed. Although L_1 linear fabrics are typically nearly coaxial with F_2 folds, they show systematic variation consistent with reorientation by F_2.

D_2 deformation is mainly manifested as kinks and crenulations, open warps of F_2 axial traces, and transposition of F_2 axial surfaces, for example in the Wawa–Quetico boundary region. The northwestern contact of the Manitouwadge belt to the Nama Creek pluton was the locus of post-D_2 ductile shearing, sinistral in map view, along the D_2 Nama Creek shear zone. Near this zone, structural and aeromagnetic trends are discontinuous, and outcrop-scale F_2 folds and earlier fabrics are strongly transposed into L > S tectonite. In the southeastern area of the Manitouwadge belt, the Banana Lake antiform, previously considered to be an F_2 fold (Williams and Breaks, 1996), is tentatively reinterpreted to be an F_1 fold. Its orientation and style are consistent with either an F_1 or an F_2 fold. However, D_2 development is supported by the 2687 ± 2 Ma Loken Lake pluton, which postdates D_1 deformation and yet contains a tectonic fabric (S_2) folded by the Banana Lake antiform.

A number of brittle-ductile faults were previously identified by Pye (1957), Milne (1974), and Williams and Breaks (1990a). The Agam Lake fault (Pye, 1957) forms a pronounced strike-parallel topographic lineament, mostly in metagreywacke, but juxtaposing metagreywacke and mafic volcanic rocks along its western extremity. In metagreywacke, it corresponds to a change in aeromagnetic signature, suggesting possible involvement of volcanic rocks in the subsurface. The Rabbitskin Lake fault forms a broad lineament defined by topographic and aeromagnetic lows. The inferred fault apparently truncates or attenuates orthoamphibole-bearing rocks, and may be responsible for the strong attenuation of the northern limb of the Manitouwadge synform, implying sinistral movement in map view. The Agam Lake fault and Rabbitskin Lake fault, and two faults west and southwest of Garnet Lake, are apparently folded by the Manitouwadge synform.

High-angle faults are marked by low aeromagnetic and low topographic lineaments, and by displacement of lithological units and aeromagnetic trends. The northerly trending Castaway and Fox Creek faults show apparent sinistral displacement and, in the case of the Fox Creek Fault, intersected by underground workings at the Geco mine, 60 m of left-lateral and east-side-up displacement is recorded (Pye, 1957; Brown et al., 1960). In general, the trends of faults are subparallel to those of Proterozoic Matachewan and Marathon dykes, that is northwesterly and northerly, respectively. Diabase dykes are commonly intruded along faults, or along some segments of faults and, in some cases, dykes and faults cannot be discriminated on the basis of aeromagnetic signature.

AGE AND STRUCTURAL RELATIONSHIPS OF PLUTONIC ROCKS

The Black Pic batholith comprises a range of plutonic rocks from diorite to granite to layered gneissic rocks, commonly with several generations of intrusions represented on a single outcrop. The magmatic and deformation history of the region was partly addressed by U-Pb dating of zircons in plutonic rocks (Zaleski et al., 1996). The oldest recognized phase of the Black Pic batholith, 2689 ± 2 Ma monzodiorite (Fig. 15), which varies to granodiorite and quartz diorite, has a tectonic foliation folded by the Manitouwadge synform. The contact to the Manitouwadge belt is a transitional zone of mafic metavolcanic rocks and monzodiorite sheets involved in tight folds and, in many cases, cut by younger foliated tonalite and aplite-pegmatite. Inside the Manitouwadge synform, the 2687 ± 2 Ma Loken Lake pluton is a foliated microline-megacrystic granodiorite, commonly with an L > S tectonic fabric (Fig. 16). Both the Black Pic monzodiorite and Loken Lake pluton are interpreted as pre- to syn-D_2 intrusions.

Several intrusions, all with tectonic fabrics interpreted as late-D_2 to syn-D_3, have emplacement ages of ca. 2680–2675 Ma. The Nama Creek pluton, 2680 ± 2 Ma microline-megacrystic granodiorite to monzodiorite, forms a long sinuous body along the margin of Manitouwadge belt and is folded by the F_3 Blackman Lake antiform. The foliation of the 2677 ± 2 Ma Banana monzodiorite pluton is folded around the F_3 Banana Lake antiform. Everest Lake pluton underlies an elongate area, slightly modified here from Williams and Breaks (1990a), along the Wawa–Quetico boundary. Its northern contact consists of a wide zone in which monzodiorite to diorite sheets are interleaved and deformed together with Quetico paragneiss (Fig. 17). Both the paragneiss and the monzodiorite sheets have abundant *in situ* anatectic segregations, implying that migmatization, at least in part, postdated the 2679 ± 2 Ma emplacement of the pluton.

WAWA–QUETICO SUBPROVINCE BOUNDARY

The Wawa–Quetico boundary in the Manitouwadge area is transitional on the basis of lithological, structural, and metamorphic criteria. The similarity in depositional age constraints and in composition between metagreywacke in the Manitouwadge greenstone belt and that in the Quetico Subprovince, and the folding of the latter together with metavolcanic rocks in the Jim Lake antiform, argue for correlation of the sedimentary sequences across the subprovince boundary. There is a northward change from macroscopic F_2 folds with broad hinge regions and northeasterly trending axial planes that dip toward the south, to tight F_2 folds with subvertical dips and east-west trends parallel to the dominant structural trends of the Quetico Subprovince (Fig. 14). The increased transposition and increased D_2-D_3 strain toward the Wawa–Quetico boundary may partly reflect the contribution of the thermal gradient to ductility contrasts during metamorphism.

ECONOMIC GEOLOGY

The Geco deposit was by far the largest in the Manitouwadge belt (Table 1A), producing 55.9 Mt of ore during its lifetime with an overall grade of 1.9% Cu, 3.8% Zn, and 47 g/t Ag (H. Lockwood, written comm., 1994). Combined production from the Willroy, Nama Creek, and Willecho deposits, similar deposits of the Willroy mine in the 1970's, was 8.7 Mt with a grade of 0.9% Cu, 4.9% Zn, and 50 g/t Ag. Except for Nama Creek, each deposit comprised several massive, semimassive, and stockwork sulphide orebodies of varying Cu, Zn, and Pb grades and ratios. In terms of classification based on Cu-Zn-Pb, the Manitouwadge belt collectively fits Franklin's (1986) Cu-Zn group 1a, which includes the deposits of the Abitibi greenstone belt, the Flin Flon–Snow Lake belt, and other areas dominated by basalt-volcanism (Zaleski and Peterson, 1995).

On the basis of Cu-Zn-Pb proportions, nature of mineralization and metamorphism, and Economic Minerals of Canada, Part A, (ed.) R.J.W. Douglas; Geological Survey of Canada, Economic Geology Report 1, 44–54.

Suffel, G.G., Hutchinson, R.W., and Ridler, R.H.
1971: Metamorphism of massive sulfides at Manitouwadge, Ontario, Canada; Society of Mining Geologists of Japan, Special Issue No. 3, p. 235–240.

Timms, P.D. and Marshall, D.
1959: The geology of the Willroy mines base metal deposits; Proceedings of the Geological Association of Canada, v. 11, p. 55–65.

Williams, H.R.
1990: Subprovince accretion tectonics in the south-central Superior Province; Canadian Journal of Earth Sciences, v. 27, p. 570–581.

Williams, H.R. and Breaks, F.W.
1989: Geological studies in the Manitouwadge–Hornpayne area; Ontario Geological Survey, Miscellaneous Paper 146, p. 79–91.
1990a: Geology of the Manitouwadge–Hornpayne area; Ontario Geological Survey, Open File Map 142, scale 1:50 000.
1990b: Geological studies in the Manitouwadge–Hornpayne area; Ontario Geological Survey, Miscellaneous Paper 151, p. 141–147.
1996: Geology of the Manitouwadge–Hornpayne region, Ontario; Ontario Geological Survey, Open File Report 5953, 138 p.

Williams, H.R., Stott, G.M., Heather, K.B., Muir, T.L., and Sage, R.P.
1991: Wawa Subprovince; in Geology of Ontario, Special Volume 4, Part 1, p. 485–539.

Wolfson, I.
1991: Assessment work report, One Otter Lake East area 1991, Schreiber–Hemlo District; Thunder Bay Resident Geologist's Office, Assessment File 2.14620, Noranda Minerals Inc., 49 p., 2 maps (MDI-42F05SE-0001).

Zaleski, E. and Peterson, V.L.
1995: Depositional setting and deformation of massive sulfide deposits, Iron formation and associated alteration in the Manitouwadge greenstone belt, Superior Province, Ontario; Economic Geology, v. 90, p. 2244–2261.

Zaleski, E., Peterson, V.L., Lockwood, H., and van Breemen, O.
1995a: Geology, structure and age relationships of the Manitouwadge greenstone belt, northwestern Ontario; in Field guidebook: Institute on Lake Superior Geology, 41st Annual Meeting, May 13–18, 1995, Marathon, Ontario, Proceedings Volume 41, Part 2b, 77 p.

Zaleski, E., Peterson, V.L., and van Breemen, O.
1994: Geological, geochemical, and age constraints on base metal mineralization in the Manitouwadge greenstone belt, northwestern Ontario; in Current Research 1994-C; Geological Survey of Canada, p. 225–235.
1995b: Geological and age relationships of the margins of the Manitouwadge greenstone belt and the Wawa–Quetico subprovince boundary, northwestern Ontario; Current Research 1995-C, Geological Survey of Canada, p. 35–44.

Zaleski, E., van Breemen, O., and Peterson, V.L.
1996: Geological evolution of the Manitouwadge greenstone belt and Wawa–Quetico subprovince boundary, Superior Province, Ontario, constrained by U-Pb zircon dates of supracrustal and plutonic rocks; Canadian Journal of Earth Sciences, v. 36, p. 945–966.

ACKNOWLEDGMENTS

Hugh Lockwood, Battle Mountain Gold, formerly Chief Geologist at Noranda's Geco mine was instrumental in the initiation of the Manitouwadge project, and contributed greatly to its success by facilitating the exchange of information and the development of ideas. Otto van Breemen, Geological Survey of Canada, has worked with us on geochronology and contributed to our understan