

Geology of the Meadowbank Iron Formation-Hosted Gold Deposit

Sherlock, R.L.¹; Alexander, R.B.²; March, R.²; and Kellner, J.²
 1. Canada-Nunavut Geoscience Office, Iqaluit, Nunavut.; 2. Cumberland Resources Ltd., Vancouver, B.C.
 GSC Open File 3149




Figure 1: Regional geology of northwestern Hudson Bay showing the distribution of supracrustal rocks. The Meadowbank deposit is located within the Woodburn Lake group, approximately 70 km north of Baker Lake, Nunavut.




Figure 2: Simplified geological map of the Woodburn Lake group in the Meadowbank area, modified from Henderson and Henderson (1994) and Zaleski et al. (1997a; 1999a).

Common Legend

- Granitoid rocks (2.60-2.62 Ga), massive to weakly foliated
- Gabbro, massive plagioclase-amphibole intrusive
- Quartzite, quartz-pebble (oligomictic) conglomerate
- Ultramafic schist, talc-amphibolite, locally spinifex-textured
- Banded iron formation, magnetite-quartz-amphibole ± pyrrhotite, pyrite and gold
- Intermediate-felsic volcaniclastic rocks; dacite-rhyodacite in composition, locally showing well-defined bedforms and blue quartz phenocrysts (2.72-2.71 Ga)
- Intermediate-felsic volcanic rock; dacite-rhyodacite in composition, massive unit likely flow or shallow intrusions, locally abundant blue quartz phenocrysts
- Apparent dip of S/S₂ composite fabric, extrapolated from drill sections
- S/S₂ composite fabric, dipping & vertical
- F/F₂ fold axis
- S fabric
- F, fold axis




Figure 3: Detailed geology of the Meadowbank gold deposits, at 1:7,500 scale.




Figure 4: Geology of the Third Portage area showing the axial trace of the two fold sets.




Figure 5: Cross sections through the Meadowbank gold deposits. A-A' is through the North Portage deposit; B-B' is a section through the Third Portage deposit; and C-C' is through the Goose Island deposit. In all three areas gold mineralization is best developed in high strain zones localized at or near the structural footwall of the ultramafic rocks within the iron formation-felsic volcaniclastic sequence.

INTRODUCTION

The geology of northwest Hudson Bay consists of Archean and Proterozoic supracrustal sequences and plutonic rocks (Figures 1 & 2). The Archean supracrustal sequences (e.g. Woodburn Lake and Prince Albert groups) are intruded by (ca. 2.60 Ga) orogenic granites, which are in turn unconformably overlain by pre-Hudsonian Proterozoic basins (e.g. Amer and Hurwitz groups). Post-Hudsonian basins (e.g. Baker Lake and Thelon Basins) unconformably overlie both sequences forming an overlap assemblage (Hoffman, 1990). The Woodburn Lake group, which hosts the Meadowbank gold deposits, is Archean in age, but has been overprinted by the Paleoproterozoic Hudsonian orogeny (Henderson et al., 1991), the significance of which with respect to gold metallogeny is uncertain.

The Meadowbank area (Sherlock et al., 2001; Figures 2 & 3) has been the focus of exploration activities since its discovery by Asamera Minerals Inc. in 1987 (Armitage et al., 1996). Most of the work in recent times (1995-1999) has focused on definition of the Third Portage, North Portage, Goose Island and the Bay Zone deposits, collectively referred to as the Meadowbank deposits. Gold in the Meadowbank deposits is hosted by complexly deformed iron formations and felsic-intermediate volcaniclastic rocks. Currently a classified resource of about 2 million ounces of gold has been defined (Cumberland Resources Inc. Press Release, March 29, 2000).

This map (Figure 3) was produced through a combination of surface mapping, trench mapping and extrapolation of geology, as interpreted from drill sections, to the surface.

LITHOLOGIES

The Woodburn Lake group (Figures 2 & 3) includes quartzite and oligomictic conglomerate, ultramafic to mafic volcanic rocks, felsic to intermediate volcanic rocks and derived sediments with interbedded iron formation (Henderson et al., 1991; Zaleski et al., 1997b; 1999b; Sherlock et al., 2001). Several phases of deformation have affected the stratigraphy with four events recognized regionally (Pehrsson et al., 2000) two of which have affected the geometry of the Meadowbank gold deposits.

Felsic-Intermediate Volcanic Rocks (IV)

The felsic to intermediate volcanic rock package is the dominant lithology recognized in the Meadowbank area. Its age ranges from 2.72 to 2.71 Ga (Davis and Zaleski, 1998; Zaleski et al., 2000). Two subtypes are recognized:

- Massive felsic flows to subvolcanic intrusives (Figure 3) are recognized to the east, in the structural footwall of the mineralized intervals at Third Portage. This unit is blue quartz and plagioclase feldspar phytic with a groundmass of fine-grained quartz and biotite.
- Bedded volcaniclastic rocks (Figure 3) comprise the bulk of the map area. Bedforms range from 1 cm to 3 m thick and rarely show unambiguous internal sedimentary structures, such as graded beds. The beds are well sorted and homogeneous in composition, but are not graded, suggesting mass flow emplacement. Alternatively these may represent agglomerated volcaniclastic beds. Beds are composed mainly of quartz and feldspar grains, which display a granular clastic texture, along with an assemblage of epidote-biotite-chlorite-muscovite. Locally blue quartz and plagioclase feldspar phenocrysts are recognized, similar to the massive units.

Iron Formation

Intervals of iron formation comprising an assemblage of magnetite, quartz and amphibole (granule- and concretion-like) are interbedded with the felsic-intermediate volcanic rocks. Several iron formations have been identified, including the East BIF, Central and West BIF (Figures 2 & 3). Of these only the Central BIF is surficial. In areas of low strain, beds within the iron formation are typically 0.2 to 10 cm thick and consist of alternating nonmetallic layers of magnetite, quartz & amphibole. To the south, near Goose Island, garnet and biotite are developed in the iron formation, suggesting higher metamorphic grades. Locally where auriferous, magnetite has been replaced by pyrrhotite and lesser pyrite.

Quartzite

Massive to bedded and foliated quartzite are exposed (Figures 2 & 3) in the western portion of the Meadowbank area in the structural hangingwall of the immediate deposit stratigraphy. The bedded material has centimeter-scale bed forms defined by selvages of muscovite. The foliated quartzite lacks bedforms and has correspondingly more muscovite, defining the foliations.

Ultramafic Rocks

The ultramafic rocks comprise a massive sequence of talc-amphibolite-chlorite with textural variations in the size and color of the amphiboles. This unit does not crop out in the immediate Meadowbank area but is commonly intersected in drill cores. Only rarely are primary features such as spinifex textures and foliations visible. Locally, within the ultramafic sequence, mafic volcanic intervals have been identified.

Intrusive Rocks

The supracrustal rocks at Meadowbank are bracketed to the east and west by large biotite-bearing granitoid bodies (Figure 2). These rocks, range in age from 2621 ± 2 to 2599 ± 5 Ma (Ashton, 1989) and are dated at 2812 ± 4 (Roddick et al., 1992) in the immediate Meadowbank area. These intrusives locally preserve the dominant S₁ fabric described below. Locally, quartz and feldspar phytic dykes are intersected in drill core between the Third and North Portage deposits. These bodies are discontinuous, having been rotated into the transposition fabric, and locally contain Cu-Au mineralization (Sherlock et al., 2001). Narrow undeformed lamprophyre dykes are also recognized in drill cores.

METAMORPHISM

The metamorphic assemblages at Meadowbank are characterized by chlorite-muscovite-biotite-calcite in the volcanosedimentary strata; interpreted to reflect regional greenschist facies metamorphic conditions. The iron formations are characterized by an assemblage of quartz-amphibole-magnetite ± sulphides. To the south, at Goose Island, the iron formations are characterized by a quartz-amphibole-biotite-garnet mineral assemblage. This suggests a transition from greenschist, at Third Portage, to lower amphibolite metamorphic assemblages at Goose Island with the garnet isograd transecting Third Portage and Goose Island deposits.

STRUCTURE

The volcanosedimentary sequence at the Meadowbank area is polydeformed with four events recognized regionally (Pehrsson et al., 2000) two of which have significantly affected the geometry of the mineralized bodies. We have adopted the deformation nomenclature of Pehrsson et al. (2000).

A composite transposition fabric (S/S₂) is the principal fabric observed in the Meadowbank area. The degree of transposition observed indicates that compositional layering preserved throughout the area should not be considered primary. The earliest event recognized D (D₁) in the terminology of Pehrsson et al. (2000) is considered here as a progressive event with refolding or buckling of the transposition fabric, generating light to isoclinal folds (F₁) which exhibit moderately steeply dipping enveloping surfaces in the Third Portage area (Sherlock et al., 2001). The axial trace of these early folds is shown in Figure 4. The culmination of the progressive deformation event involves the development of penetrative planar fabrics and local high strain zones. These high strain zones, often exploit lithologic contacts where rheologic contrasts have focused strain gradients into the volcanosedimentary rocks. These fabrics are commonly spatially related to gold mineralization. In the North Portage, Third Portage and Goose Island sections (Figure 5) the strata are folded about a NE verging antiform-synform pairs. The high strain zones are often localized along fold limbs at the ultramafic-volcanosedimentary contacts due to contrasts in the mechanical properties of the strata. Garnet porphyroblasts are commonly deformed by these high strain zones.

D₂ deforms and refolds earlier deformation fabrics. F₂ geometries are typically open to close, angular to subrounded and often "S" shaped in the Meadowbank area. Axial surfaces are upright, NE-SW trending and doubly plunging and locally controlled by the orientation of the earlier fabrics. The axial trace of F₂ is shown in Figure 4. Fold interference patterns are common resulting from the superposition of upright F₁ folds on light and inclined F₂ folds. A spaced to penetrative axial planar cleavage (S₂) is locally well developed and overprints earlier fabrics as well as crenulating amphibole rosettes. Significantly D₂ folds and deforms mineralized envelopes, suggesting that it is a post-mineralization event.

MINERALIZATION

Gold is contained within sulphide-bearing iron formation (Armitage et al., 1996; Kerswill et al., 1998; Kerswill, 2000; Sherlock et al., 2001), characterized by disseminated to semimassive pyrrhotite-pyrite which preferentially replaces magnetite (Armitage et al., 1996). Gold is also contained within felsic-intermediate volcanic package as disseminated pyrrhotite and pyrite along with narrow blue-grey quartz-sulphide veins. Gold mineralization often shows a strong spatial relationship to high strain zones described above.

Armitage et al. (1996), based on whole rock and mineral chemistry, has shown that auriferous iron formations are characterized by a mineral assemblage of cummingtonite and biotite with pyrrhotite-pyrite-magnetite and quartz. Barren, but sulphidic, iron formations are characterized by an assemblage of garnet ± hornblende + sillpionomelane with pyrrhotite, pyrite, quartz and magnetite. Whereas barren, nonsulphide-bearing, iron formations are characterized by quartz and magnetite. Based on these mineral assemblages and their relationship to gold mineralization, Armitage et al. (1996) has suggested that mineralization is related to syn-deformation metasomatic fluids that have altered the iron formation with the introduction of Mg-K-Ca-S-As-Cu and Au. Based on the fabric relations recognized gold, and associated sulphides, were introduced into dilation sites during progressive D₂ deformation (Sherlock et al., 2001). The mechanical contrast between the ultramafic and volcaniclastic rocks partitioned strain which allowed fluid influx and the resulting alteration assemblages to be concentrated in the higher strain corridor adjacent to the ultramafic rocks. The progressive nature of deformation event resulted in the various relationships between sulphide minerals and deformation fabrics.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the support provided by Cumberland Resources Ltd. We also would like to thank Bill Barclay, John Kerswill, Sally Pehrsson, Lori Wilkinson and Eva Zaleski for helpful discussions on the Meadowbank geology. This open file has benefited from a critical review by Sally Pehrsson and David Scott.

REFERENCES

Ashton, K.E. 1988. Precambrian geology of the southeastern Amer Lake area (66N41'), near Baker Lake, N.W.T. Ph.D. thesis, Queen's University, 335 p.

Armitage, A.E., James, R.S., and Goff, S.P. 1996. Gold mineralization in Archean banded iron formation, Third Portage Lake area, Northwest Territories, Canada. *Exploration and Mining Geology*, 5, 5: 1-15.

Davis, W.J., and Zaleski, E. 1998. Geochronological investigations of the Woodburn Lake group, western Churchill province, Northwest Territories: preliminary results. In *Radiogenic Age and Isotopic Studies: Report 11*, Geological Survey of Canada, Current Research 1998-F, 89-97.

Henderson, J.R., and Henderson, M.N. 1994. *Geology of the Whiteshell-Tehok Lakes Area*. District of Keewatin, Northwest Territories (parts of 56D, 56E, 56A and 56H). Geological Survey of Canada, Open File 2923, scale 1:100 000.

Henderson, J.R., Henderson, M.N., Pryor, L.L., and Crosswell, R.G. 1991. *Geology of the Whiteshell-Tehok Lakes area*. District of Keewatin and Archean supracrustal belt with iron formation hosted gold mineralization in the central Churchill Province, Geological Survey of Canada, Current Research, Part C, paper 91-1C, p. 149-156.

Hoffman, P.F. 1990. Subdivision of the Churchill Province and extent of the Trans-Hudson Orogen, in Lewy, J.F. and Stauffer, M.R., eds. *The Early Proterozoic Trans-Hudson Orogen of North America*. Geological Association of Canada, Special Paper 37, p. 15-39.

Kerswill, J.A., Goff, S.P., Wilkinson, L., Jenner, G.A., Kjarsgaard, B.A., Bretzfeld, R., and Samaras, C. 1998. An update on the metallogeny of the Woodburn Lake group, western Churchill Province, Northwest Territories, Geological Survey of Canada Current Research 1998-Cp, 29-41.

Kerswill, J.A., Goff, S.P., Kjarsgaard, B.A., Jenner, G.A., and Wilkinson, L. 2000. Highlights of recent metallogenic investigations in western Churchill Province, Nunavut, Canada: implications for mineral exploration in Archean greenstone belts. (extended abstract) in *Geoscience 2000: The Millennium Geoscience Summit*, CD-ROM, Calgary, Abstract 736.

Kerswill, J.A. 2000. Iron formation-hosted gold deposits: a view from Nunavut with emphasis on "Lupin-Blau" deposits. [extended abstract with maps] in *Abstract Volume, Short Course on Geology and Mineral Deposits of Nunavut Territory*, Society of Economic Geologists, London Student Chapter, March 3, 2000, 10 p.

Pehrsson, S., Wilkinson, L., Zaleski, E., Kerswill, J., and Alexander, B. 2000. Structural geometry of the Meadowbank deposit area, Woodburn Lake group: implications for a major gold deposit in the Western Churchill Province. Extended abstract in *GeoCanada 2000*, Calgary Alberta Digital Release.

Roddy, J.C., Henderson, J.R., and Chapman, H.J. 1992. U-Pb ages from the Archean Whiteshell-Tehok lakes supracrustal belt, Churchill Province, District of Keewatin, Northwest Territories. *Radiogenic Age and Isotopic Studies*, report 8, Geological Survey of Canada, paper 1992-2, p. 31-40.

Sherlock, R.L., Alexander, R.B., March, R., Kellner, J., and Barclay, W.A. 2001. Geologic setting of the Meadowbank iron formation-hosted gold deposits. *Geological Survey of Canada, Current Research 2001-C11*, Geological Survey of Canada, 10 p.

Zaleski, E., Corrigan, D., Kjarsgaard, B.A., Kerswill, J.A., Jenner, G.A., and Henderson, J.R. 1997a. *Geology, Woodburn Lake group, Meadowbank River to Tehok Lake (66N41', 56E41')*. District of Keewatin (Nunavut), Northwest Territories. Geological Survey of Canada, Open File 3461, scale 1:50 000.

Zaleski, E., Corrigan, D., Kjarsgaard, B.A., Kerswill, J.A., Jenner, G.A., and Henderson, J.R. 1997b. Preliminary results of mapping and structural interpretation from the Woodburn project, western Churchill Province, Northwest Territories. *Geological Survey of Canada, Current Research 1997-Cp*, 91-100.

Zaleski, E., Duke, N.L., L'Heureux, R., and Wilkinson, L. 1999a. *Geology, Woodburn Lake group, Amalik Lake to Tehok Lake, Kivalliq Region, Nunavut*. Geological Survey of Canada, Open File 3745, scale 1:50 000.

Zaleski, E., L'Heureux, R., Duke, N., Wilkinson, L., and Davis, W.J. 1999b. Komatiitic and felsic volcanic rocks overlain by quartzite, Woodburn Lake group, Meadowbank River Area, Western Churchill Province, Northwest Territories (Nunavut). *Geological Survey of Canada, Current Research 1999-Cp*, p. 9-18.

Zaleski, E., Davis, W.J., and Wilkinson, L. 2000. Sasamitocover relationships, unconformities and depositional cycles of the Woodburn Lake group, western Churchill Province Nunavut. [abstract] 28th Yellowknife Geoscience Forum, November 2000, N.W.T. Chamber of Mines, p. 91-92.

