


LEGEND

Conglomerate.....	
Limestone / Dolomitic limestone.....	
Carbonaceous shale.....	
Coal.....	
Siderite concretion bed or calcrete concretions.....	
Bentonite bed.....	
Oolitic bed.....	
Stromatolite bed or individual stromatolites.....	
Lens-shaped bed.....	
Discontinuous scour / gutter fills.....	
Fault.....	
Fractures with slickensides (either structural or pedogenic).....	
Fining-upward Trend.....	
Coarsening-upward Trend.....	
Paleocurrent Indicators.....	
Copper Sulfide Mineralization.....	
Erosive base with rip-ups and granules.....	
Scoured Base.....	
Ball and Pillow.....	
Rip-up Interclasts.....	
Breccia / Flat Pebble Conglomerate.....	
Trough Cross bedding.....	
Ripple Cross Lamination.....	
Climbing Ripples.....	
Low Angle Lamination.....	
Planar Tabular Crossbedding.....	
Inclined Bedding Surfaces (IBS) or Lateral Accretion Surfaces (LA).....	
Inclined Heterolithic Stratification (IHS).....	
Contorted Lamination.....	
Hummocky Cross Stratification (HCS).....	
Water Escape Structure.....	
Roots.....	
Bioturbation / Burrowing.....	
Vertical Burrows (eg. Skolithos).....	
Desiccation Cracks.....	
Fossil shells (pelecypod, gastropod, brachiopod).....	
Dinosaur bone fragments.....	
Carbonized wood fragments.....	
Gypsum nodule bed.....	
Evaporite crystal molds.....	

UPPER ORDOVICIAN/LOWER SILURIAN - SOUTHERN ONTARIO
CABOTHEAD/DYER BAY/FOSSIL HILL FMS
Brooke Roadcut, 3 km NW of Owen Sound
41A/10 Owen Sound 023373
Lat. 44°35'30"N Long. 80°58'10"W
General strike 330°-340°
Dip <1°SW



FOSSIL HILL
DYER BAY
CABOT HEAD

- buff to brown fine-crystalline dolostone, thick bedded, abundant fossils of insitu corals and *Pentamerus* brachs, sharp irregular erosional base.

- slightly c-up sequence thin bedded pale grey very fine- to fine-crystalline dolostone, horizontal lamination, soft sediment deformation, generally non fossiliferous with few thin bioclastic beds, sharp top
- green slightly silty mudstone, non calcareous, poorly exposed, bioturbated.