

QUATERNARY

SURFICIAL DEPOSITS

POST LAST GLACIATION

NONGLACIAL ENVIRONMENTS

- O ORGANIC DEPOSITS: organic matter > 1 m thick; formed by the accumulation of vegetation in poorly drained depressions (swamps and bogs); usually forms flat terrain
Ca Talus (scree): accumulations of blocks, commonly exceeding 2 m in diameter; as much as 50 m thick; forming aprons and fans below cliffs
Cca Rock Glaciers (relict): rock debris deformed by the down-slope flow of buried or interstitial ice, forming pronounced transverse and longitudinal ridges and furrows
Csr Rock slide deposits: chaotic landscape of irregular and stacked bedrock blocks; prominent in areas of steeply dipping, poorly-indurated sandstone and shale-rich beds in the Mattson and Fantiasque formations
Csd Debris slide deposits: accumulations of unconsolidated material; internal structure of material is not maintained; where sufficient moisture is present, the slide may become a flow producing characteristic levees along its lateral margins and a spatulate form at the base of slope
Cpr Bedrock slump deposits: large rotational blocks in bedrock, shallow to 10's of metres thick; internal structure of material may be retained; often traceable upslope to active scarps; where sufficient moisture is present the slump may produce a flow at its base, forming a characteristic spatulate form; prominent in areas of steeply dipping, poorly-indurated sandstone and shale-rich beds in the Mattson and Fantiasque formations; associated with the largest mass movements in the region
A Alluvial deposits: well sorted gravel and sand with detrital organic beds, including concentrations of logs, > 1 m thick; Ap, floodplains and meandering valley floors, forming meander scars and point bars; Al, terraces along valley wall sides
Af Alluvial fan: poorly sorted gravel and sand with organic detritus and buried soils; fans are commonly crossed by debris flow channels and levees and subject to shifting stream courses; > 1 m thick

POSTGLACIAL OR LATE WISCONSINAN

PROGLACIAL AND GLACIAL ENVIRONMENTS

- G Proglacial outwash: Gd, braided outwash deltas; Gdt, delta terraces; Gf, fans; Gp, outwash plains and mantling valley floors; Gt, level outwash terraces
I Ice contact stratified drift: deposited behind or at the ice margin; topography is undulating, irregular, or ridged; It, forming lateral moraine terraces and delta terraces; Ik, kettle holes; Il, hummocky moraine kame fields, or ice block disintegration terrain; Ir, eskers or crevasse fillings
Tb Till blanket: > 2 m thick; forming undulating topography that obscures underlying bedrock structure; Tbk, distinctly kettled
Tv Till veneer: < 2 m thick and discontinuous; surface mimics underlying bedrock structure

PRE-QUATERNARY

BEDROCK

- R Sedimentary bedrock, undifferentiated. The north-south aligned Kotanelee and La Biche anticlines dominates the map sheet, and are comprised largely of moderate to shallow-dipping (<25°), Lower Carboniferous lower to upper Mattson Formation strata (calcareous quartz arenite, siltstone and shale, with minor limestone, dolostone and coal). The eastern edge of the map sheet encompasses the eastern half of the valley system; the two ranges is comprised of shallow-dipping (<20°) Lower Carboniferous Scattered, Garbut and Chinkh formations (strata include shale, siltstone and sandstone). Permian Fantiasque and Tika formations are prominent along the eastern margins of the Kotanelee and La Biche ranges, and are comprised of diverse strata that includes chert, siltstone, limestone, dolostone, and sandstone. Devonian and Carboniferous Bess River Formation (mostly shale with some sandstone) is exposed in the valley bottom in the west-central part of the map sheet as well as in the small valley extending southward through the central-upper half of the map.

NOTE: In areas where the surficial cover forms a complex mosaic, the area is coloured according to the predominant unit and labelled with hyphenated letters in descending order of cover

MAP SYMBOLS



- Geological boundary (defined, gradational)
Scarp
Cirque; peaks and sharp ridges formed by glacial erosion
Moraine
Esker
Striae (glacial flow direction known, unknown)
Fluting or drumlinoid ridge parallel to ice flow (direction of flow known, unknown)
Proglacial meltwater channel (direction of flow known)
Lateral meltwater channel (barb points upslope and down flow)
Observation
Drift geochemistry sample site
Canadian Shield erratic

NOTES:
Mass Wasting is the collective term given to the range of processes and resultant landforms that relate to the gravitational downslope movement of rock and/or unconsolidated material without the direct conveyance by water, air or ice. Water and ice are, however, often key components in initiating and perpetuating mass wasting by reducing the strength of materials and in their plastic and fluid behaviour.
Different types of mass wasting are distinguished by the type of material involved (e.g., bedrock, talus, till), the mode of deformation (e.g., creep, slide, slump, flow, speed of movement, morphology of the moving mass, and water content.
Creep is the slow (mm to cm per year), often imperceptible, downslope movement of soil, talus or other unconsolidated material. Creep occurs episodically in response to solifluction, seasonal wetting and drying, or freeze-thaw cycles and may include the plastic deformation of clay-rich soils. While more prevalent on steep slopes, creep can occur on slopes <5°. Evidence of creep is seen where tree trunks or structures (e.g., hydro poles) are tilted downslope, soil accumulates on retaining walls, and cracks develop in the soil perpendicular to the slope. Creep is also responsible for the formation of gelification lobes, prominent, small-scale (metres in length, centimetres thick), periglacial landforms found along the upper reaches of local mountain ranges (but not included in the regional surficial geology mapping).
Slides are the rapid, downslope movement of bedrock or unconsolidated material. Failure occurs along bedding and/or fracture planes in bedrock, and along bedrock contacts, or structural and sedimentological boundaries within unconsolidated material. Slides can be initiated at shallow or considerable depths. Slumps involve the rotational movement of bedrock and/or unconsolidated material along failure planes. Slumps may occur as individual blocks or amorphous masses inflicting water content and structural integrity of the falling material. Slumps often extend progressively up-slope through time, and can be associated with active scarp or headwall retreat. Slumps can be initiated by failure along bedding, fractures, or sedimentological planes, by infiltration of surface water, through lateral incision and undercutting of slopes by streams, or excavation activities (e.g., road building, pipeline trenching). Slumps are prominent in areas of steeply dipping, poorly-indurated sandstone and shale-rich beds in the Mattson and Fantiasque formations, and are associated with the largest mass movements in map area.
While different earth surface materials and geological settings are often strongly associated with various types of mass wasting, predicting their occurrence, magnitude and rate of deformation is often not possible. Some areas that are prone to mass wasting include regions of steeply dipping bedrock, poorly indurated and shale-rich bedrock, and along steep slopes, soil accumulation, and areas of retaining walls. Activities such as road building, pipeline trenching, logging and seismic exploration can also initiate mass wasting, particularly where they undercut slopes, or act to destabilize surficial materials.

Glacial History: The Babiche Mountain map area was glaciated during the last (late Wisconsinan) glaciation (ca. 25-10 000 years ago) by the continental Laurentide Ice Sheet flowing from the northeast (Keweenaw Sector) and by the Cordilleran Ice Sheet flowing from the west. The Laurentide Ice Sheet dispersed disintegrating granite blocks or amorphous masses inflicting water content and structural integrity of the falling material. Slumps often extend progressively up-slope through time, and can be associated with active scarp or headwall retreat. Slumps can be initiated by failure along bedding, fractures, or sedimentological planes, by infiltration of surface water, through lateral incision and undercutting of slopes by streams, or excavation activities (e.g., road building, pipeline trenching). Slumps are prominent in areas of steeply dipping, poorly-indurated sandstone and shale-rich beds in the Mattson and Fantiasque formations, and are associated with the largest mass movements in map area.
While different earth surface materials and geological settings are often strongly associated with various types of mass wasting, predicting their occurrence, magnitude and rate of deformation is often not possible. Some areas that are prone to mass wasting include regions of steeply dipping bedrock, poorly indurated and shale-rich bedrock, and along steep slopes, soil accumulation, and areas of retaining walls. Activities such as road building, pipeline trenching, logging and seismic exploration can also initiate mass wasting, particularly where they undercut slopes, or act to destabilize surficial materials.

OPEN FILE DOSSIER PUBLIC 1558
GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA
2003
Open files are products that have not gone through the normal publication process.
Les dossiers publics sont des produits qui n'ont pas passé soumise au processus officiel de publication de la CGC.

Recommended citation: Smith, L.R. 2003. Surficial Geology, Babiche Mountain (95C/8), Yukon Territory - Northwest Territories; Geological Survey of Canada, Open File 1558, 1 map, scale 1:50 000.



CONTOUR INTERVAL 100 FEET / Elevations in Feet above Mean Sea Level

OPEN FILE 1558
SURFICIAL GEOLOGY
BABICHE MOUNTAIN
YUKON TERRITORY - NORTHWEST TERRITORIES
Scale 1:50 000/Échelle 1/50 000

Universal Transverse Mercator Projection / Système de référence géodésique nord-américain, 1983

Compilation by L.R. Smith based on fieldwork and studies of vertical air photographs 2000-2002. THIS MAP IS A PRODUCT OF THE CENTRAL FORELAND NATMAP PROJECT.
Surficial geology from field work by L.R. Smith 2000-2002. Additional data from T.E. Kubilj 1996.
Digital cartography by L.R. Smith.
Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.
Base map at the same scale published by Surveys and Mapping Branch in 1971.

Table with 3 columns: File Number, Name, and Coordinates. Includes entries like Tika Creek, Chinkh Creek, Mount Flett, Brown Lake, Babiche Mountain, Fisherman Lake, Mount Merrill, Mount Martin, and Betalamea Lake.