

LEGEND

This legend is common to Open Files 1508 to 1612, and 1628 to 1621. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

SURFICIAL DEPOSITS

QUATERNARY

HOLOCENE

FLUVIAL SEDIMENTS: alluvium, gravel and sand, 2–20 m thick.

- Ap** Alluvial plains: active braided floodplains; includes active proglacial outwash.
- At** Alluvial terraces
- Af** Alluvial fans

MARINE AND GLACIAL MARINE SEDIMENTS: gravel, silt, sand, and clay, 1–20 m thick, deposited in deltaic and beach environments during regression of the postglacial sea.

- Mr** Beach sediments: gravel and sand, 1–5 m thick, forming ridges and swales.
- Mt** Deltaic sediments: clay, silt, sand, and gravel, 6–20 m thick, forming coarsening upward sequences under dissected terraces.
- Mv** Deepwater proglacial silt veneers: silt, clay silt, and fine sand with dropstones, 1–2 m thick.
- Mb** Deepwater proglacial silt blankets: silt, clay silt, and fine sand with dropstones and minor gravel, 2–10 m thick.

GLACIAL LAQUSTRINE SEDIMENTS: clay, silt, sand, and gravel deposited in glacial dammed lakes in deepwater, beach and deltaic environments.

- Lr** Beach sediments: sand and gravel, 1–5 m thick, forming beach ridges.
- Lt** Deltaic sediments: clay, silt, sand, and gravel, 5–20 m thick, forming coarsening upward sequences under dissected terraces.
- Lv** Deepwater proglacial silt veneers: silt, clay silt, and fine sand with dropstones, 1–2 m thick.
- Lb** Deepwater proglacial silt blankets: silt, clay silt, and fine sand with dropstones, 2–5 m thick.

GLACIOFLUVIAL SEDIMENTS: gravel and sand, 1–10 m thick, deposited behind, at, and in front of the ice margin.

- Gp, Lt** Proglacial outwash: gravel and sand, 1–10 m thick, forming braided floodplains. Gp: terraces, Lt: fans, Gt.
- Gr, Gh** Ice contact stratified drift: gravel and sand, 1–5 m thick, forming eskers, Gr; and kames, Gh.

EARLY HOLOCENE AND WISCONSINAN


- Tm** End moraine: 5–80 m high, composed of or mantled by till, deposited in subglacial and ice marginal environments; lithic composition generally reflects underlying bedrock.
- Tv** Till veneer: 0.5–2 m thick and discontinuous.
- Tvw** Till veneer: 0.5–2 m thick, surface armored by stones due to washing by subglacial meltwater.
- Tb** Till blanket: 2–10 m thick forming an undulating blanket with drumlins and ribbed moraines in places.
- Tbr** Till blanket: 2–10 m thick forming ribbed (frozen) moraines.

BEDROCK

PRE-QUATERNARY


- R** **ROCK:** rock of various compositions and ages (Jackson and Sangster, 1987) variably modified by glacial erosion during the Quaternary and with patchy till cover; hilly and hummocky surfaces, ice moulded in places, with lake basins in subglacially scoured regions; smooth surfaces exhibiting little or no sign of glacial erosion in postglacial intervals (Dyke, 1983); cliffs resulting from glacial over-steepening in places veneered by thin till, commonly bouldery.

Geological boundary (defined, assumed)
 Areas covered by periglacial icefields during the Little Ice Age (indicated by a white pattern)
 Kettle (large)
 Glacial lake spillway
 Glacial lake limit
 Marine limit
 Escarpment
 Lateral meltwater channel; barb on upslope side
 Subglacial and proglacial meltwater channel (small)
 Esker
 Kame
 Ice contact face
 Ribbed moraine
 Lateral moraine
 End moraine
 Margin of dispersal train; teeth toward axis, steep side of teeth face down ice
 Drumlinoid hill
 Drag-and-tail
 Ice moulded bedrock
 Striae (ice flow direction known, unknown)
 Crossed striae (numbers indicate relative age, 1 being the oldest)
 Field observation site: boundary diameter (bd), boundary gravel (bg), clay (c), diameter (d), gravel (g), gravely sand (gs), mud (m), muddy sand (ms), rock (r), sand (s), sandy gravel (sg), stony mud (sm), till (t)
 Field observation site: material as above near rock outcrop
 Marine limit elevation (metres)



Geology by A.S. Dyke, 2002
 Field data provided by De Beers Canada Inc., 2002
 Digital cartography by M.M. Proulx, Earth Sciences Sector Information Division (ESS Info)
 This map was produced from processes that conform to the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001:2000 standard

OPEN FILE 1611
 SURFICIAL GEOLOGY
GIFFORD FIORD (EAST)
 BAFFIN ISLAND
 NUNAVUT



Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.
 Digital base map from data compiled by Geomatics Canada, modified by ESS Info
 Locational accuracy of the base appears to be ±100m based on plotting of GPS measured field site locations
 Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area
 Mean magnetic declination 2003, 39°18'W, decreasing 42.8' annually
 Elevations in metres above mean sea level

47 E13 OF 1601	47 E14 OF 1600	47 E15 OF 1599	47 E16 OF 1598
47 E12 OF 1602	47 E11 OF 1603	47 E10 OF 1604	47 E9 OF 1605
47 E5 OF 1608	47 E6 OF 1608	47 E7 OF 1607	47 E8 OF 1606
47 E4 OF 1610	47 E3 OF 1611	47 E2 OF 1612	47 E1 OF 1613
47 D13 OF 1628	47 D14 OF 1629	47 D15 OF 1630	47 D16 OF 1631

OPEN FILE DOSSIER PUBLIC 1611
 GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA
 2003

Open files are products that have not gone through the GSC normal publication process.
 Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la GSC.

Recommended citation:
 Dyke, A.S.
 2003: Surficial geology, Gifford Fjord (East), Baffin Island, Nunavut, Geological Survey of Canada, Open File 1611, scale 1:50 000.

