Biotite±hornblende monzogranite, syenogranite; pink, fine- to medium-grained;

Psammite, semipelite; grey- to rusty-brown, flaggy; centimetre- to metre-scale

millimetre- to centimetre-scale black and green layers; metre-scale layers of

laterally continuous layers; abundant melt pods; local interlayers of quartzite, pelite

Hornblende-biotite±clinopyroxene amphibolite; fine- to medium-grained; alternating

coarser-grained amphibolite, semipelite and pelite; may be derived from a volcanic

Biotite±hornblende tonalite, granodiorite; grey to tan, fine- to medium-grained; locally

with ciorite inclusions; varies from weakly to strongly foliated, locally an L-tectonite.

Biotite±hornblende quartzofeldspathic orthogneiss; leucocratic gneiss of plutonic

origin; granodioritic to monzogranitic; alternating grey to white, black, pink, fine- to

medium-grained; moderately to well foliated, locally layered with concordant

syenogranitic leucosome; locally contains amphibolite and tonalite bands,

massive to moderately foliated; locally grades into megacrystic granite.

MARY RIVER GROUP (units AMa - AMp)

Bedrock outcrop massive .

Oblique-slip fault (defined) .

Mesoscopic M fold axis

Mesoscopic Z fold axis

Mesoscopic fold axial plane

(arrow indicates plunge) . .

(arrow indicates plunge) . .

(defined, approximate) . . .

Synform (upright) (arrow indicates plunge) .

Antiform (upright) (arrow indicates plunge) .

(defined, approximate)

Glacial striae; direction of ice movement undetermined .

Syncline (upright, overturned) (arrow indicates plunge) .

Anticline (upright, overturned) (arrow indicates plunge)

Synform, inferred beneath Quaternary cover (upright)

Syncline, inferred beneath Quaternary cover (upright, overturned)

Anticline, inferred beneath Quaternary cover (upright, overturned)

Cordierite±andalusite isograd (ornament on high-temperature side)

Location of field photograph with corresponding Figure number .

Sillimanite-K-feldspar±melt pod isograd (ornament on high-temperature side)

Bedding (upright, tops known) .

Bedding (overturned, tops known)

D_{2P} thrust fault (defined); teeth on hanging wall .

protolith and associated sedimentary rocks.

gabbro/anorthosite boudins; locally migmatitic.

LEGEND

This legend is common to maps OF 4432 and OF 4433. All units and symbols may not appear on this

Unconsolidated glacial drift with associated marine, lacustrine, fluvial, and bog

FRANKLIN INTRUSIONS: tholeiitic diabase dykes; medium-grained with ophitic

Ultramafic rocks; serpentinized peridotite and hornblendite; foliated to schistose; dark

Biotite-allanite±hornblende monzogranite, granodiorite; commonly grades into

quartz diorite, and psammite (unit PPD); local weak foliation.

biotite-allanite±hornblende syenogranite; massive, fine- to coarse-grained, pink to

white: abundant crosscutting veins and sheets of associated biotite-muscovite-garnet

±tourmaline pegmatite; locally contains inclusions of layered orthogneiss (unit Agn),

LONGSTAFF BLUFF FORMATION: (metamorphic mineral units PPLb - PPLs)

metamorphic assemblages; locally includes bodies of biotite-allanite±hornblende

monzogranite, granodiorite, syenogranite or biotite-garnet±muscovite±sillimanite

(fibrolite)±cordierite monzogranite, and associated biotite-muscovite-garnet

Psammite, semipelite, pelite, arkosic- and lithic-wacke; interbedded; thin- to

thick-bedded, light- to dark-grey; graded beds including inverse metamorphic

biotite-muscovite-cordierite±andalusite metamorphic assemblages; locally

includes bodies of biotite-allanite±hornblende monzogranite, granodiorite,

syenogranite or biotite-garnet±muscovite±sillimanite (fibrolite)±cordierite

monzogranite, and associated biotite-muscovite-garnet±tourmaline pegmatite.

Psammite, semipelite, pelite, arkosic- and lithic-wacke; interbedded; thin- to

thick-bedded, light- to dark-grey; graded beds; minor hornblende-bearing

calcsilicate beds and concretions; biotite-muscovite±garnet metamorphic

LONGSTAFF BLUFF FORMATION: Arkosic- and lithic-wacke; interbedded with

ASTARTE RIVER FORMATION: sulphidic schist; rusty-weathering; graphitic,

BRAVO LAKE FORMATION: dominantly basalt; pillowed, fragmental and massive

flows; light- to dark-green; mafic and ultramafic cumulates; metre-scale layers, dark

FLINT LAKE FORMATION: marble, dolomite and calcsilicate; chiefly white to grey or

iron-formation; locally includes bodies of biotite-allanite±hornblende monzogranite,

DEWAR LAKES FORMATION: quartzite and feldspathic quartzite, semipelite; grey,

white, and black; laminated, bedded, and massive, locally crossbedded; may include

Biotite±hornblende K-feldspar megacrystic monzogranite, granodiorite; pink to buff;

magnetite rich laminae; locally includes iron-formation; chiefly oxide facies with

silicate facies; metallic grey; fine- to coarse-grained; laminated to bedded; may

biotite-garnet±muscovite±sillimanite (fibrolite)±cordierite monzogranite, and

include marble, dolomite and calculicate; locally includes bodies of

K-feldspar megacrysts in a finer-grained matrix of plagioclase, quartz, biotite±hornblende: varies from weakly to strongly foliated, locally an L-tectonite

gradational into granitic and granodioritic rocks lacking megacrysts.

associated biotite-muscovite-garnet±tourmaline pegmatite.

biotite-allanite±hornblende monzogranite, granodiorite, syenogranite or

buff weathering; may include semipelite, pelite, quartzite and carbonate facies

granodiorite, syenogranite and associated biotite-muscovite-garnet±tourmaline

PPLa psammite, semipelite, pelite; thin- to thick-bedded, white, gritty surface; graded beds;

green to brown; minor volcaniclastic sedimentary beds; millimetre- to

iron-formation; gabbro; peridotite; layered peridotite-gabbro sills.

centimetre-scale laminations, dark grey to white; minor quartzite, semipelite,

minor hornblende-bearing calculicate beds and concretions.

pyrrhotite-pyrite schist and slate; sulphide-facies iron-formation.

grading; minor hornblende-bearing calculicate beds and concretions;

Psammite, semipelite, pelite, arkosic- and lithic-wacke; interbedded; thin- to

thick-bedded, light- to dark-grey; graded beds; minor hornblende-bearing

calcsilicate beds and concretions; biotite-sillimanite-K-feldspar±melt pod

deposits. Chiefly marine muds on coastal areas. Includes extensive felsenmeer in

map. The prefix 'meta' applies to all lithologies in units Agn to PPLs

texture; narrow chilled margins.

green or brown weathering.

±tourmaline pegmatite.

PILING GROUP (units PPD - PPLs)

_____ proximity to Barnes Ice Cap.

PALEOPROTEROZOIC



DESCRIPTIVE NOTES

INTRODUCTION

This map is one of a set of two bedrock geological maps which present results of 1:100 000 scale mapping by the Geological Survey of Canada (GSC) and the Canada-Nunavut Geoscience Office (C-NGO), with logistical support provided by the Polar Continental Shelf Project, in the Clyde River (OF 4432) and Blanchfield Lake (OF 4433) map areas, central Baffin Island (Fig. 1). An overview of the principal Archean and Paleoproterozoic plutonic and supracrustal units mapped during the three-year (2000–2002) partnered field project are presented in Corrigan et al. (2001) and Scott et al. (2002a, 2003). Field work in 2002 led to the completion of bedrock mapping in NTS 27 C (west in 2001, 2002, and 2003 (St-Onge et al. 2001a, b: 2002a, b, c). From north to south, the open file maps provide a continuous, wide transect across the southeastern margin of the Archean Rae craton, the Paleoproterozoic Piling Group, and the northern margin of the Paleoproterozoic arc complexes of south-central Baffin Island including the ca. 1.86-1.85 Ga Cumberland batholith.

The first systematic mapping of the area was undertaken at reconnaissance scale between 1965 and 1970 (Jackson, 1969, 1978, 1984, 2000; Jackson and Taylor, 1972), and in 1974 and 1975 (Morgan et al., 1975, 1976; Jackson and Morgan, 1978; Morgan, 1983). More detailed work was subsequently initiated in the southeastern portion of the project area, in parts of map sheets NTS 37 A, 27 B, 27 C, and 27 D (Tippett, 1980, 1984; Henderson et al., 1988, 1989; Henderson and Henderson, 1994). The most recent work in proximity to the current project area consists of bedrock mapping and U-Pb geochronology of Archean rocks in the Eque Bay area (Bethune and Scammell, 1997), and a lake sediment and water regional geochemical survey southwest of the Barnes Ice Cap Results of recent Quaternary mapping are reported in Dredge (2002), electromagnetic work in Evans et al. (2003), teleseismic monitoring in Snyder (2003), petrological studies in Allan and Pattison (2003), and petro-stratigraphic

studies in Stacev and Pattison (2003). The work presented below is based on systematic bedrock foot and helicopter traverses with approximately three

GEOLOGICAL SETTING

The Central Baffin Project area (Fig. 1) is centered on the northern margin of the eastern segment of the ca. 1.8 Ga Trans-Hudson Orogen (Hoffman, 1988; Lewry and Collerson, 1990), a Himalayan-scale collisional mountain belt that is exposed from Greenland in the east, across Baffin Island and beneath Hudson Bay, to Manitoba and Saskatchewan in the west. The northern part of the area (St-Onge et al., 2001a, b, 2002d) (OF 4432, OF 4433) comprises various orthogneisses, metamorphosed sedimentary and minor volcanic rocks of the Mary River Group, and younger felsic plutonic rocks, all of Archean age and ascribed to the Bae craton (Jackson, 1969, 2000; Bethune and Scammell, 1997, in press; Corrigan et al., 2001; Scott et al., 2002a). The central part of the study area is underlain by siliciclastic, carbonate, and mafic volcanic rocks of the Paleoproterozoic Piling Group (Morgan et al., 1975, 1976; Scott et al., 2002a and references therein), a succession that has recently been described in more detail (St-Onge et al., 2002a, b, c; Scott et al., 2002a). Various felsic plutonic rocks, ranging in age from 1.90-1.85 Ga and including the Cumberland patholith (Wodicka et al., 2002), are exposed in the southern part of the area (St-Onge et al., 2002a, b, c). Collectively, this assemblage of rocks is thought to represent a Paleoproterozoic continental margin succession (Morgan et al., 1975) originally deposited on the southern flank of the Archean Rae craton and subsequently deformed during Trans-Hudson orogenesis (see Corrigan et al., 2001, Scott et al., 2002a, and references therein) varies across the project area. A primary unconformable relationship is preserved along much of the northern margin of the belt, however, where exposed in southern domal structural culminations (Fig. 2), the unconformity is strongly

deformed. Northeast of Nauja Bay (Fig. 2), Archean orthogneisses are locally thrust onto Piling Group rocks, apparently in association with late, thick-skinned folding The primary distribution of rock types within the Piling Group varies dramatically across the strike of the belt: carbonate strata are extensively exposed on the northern margin, whereas mafic volcanic rocks are exposed on the southern margin (Scott et al., 2002a; Fig. 1). Regional metamorphic grade increases outward from the central portion rocks (St-Onge et al., 2002d). In the following sections, the rocks are described from lowest to highest structural evels, based on the results of new 1:100 000-scale mapping on NTS map sheets 27 C (west half) and 37 D (east half),

ARCHEAN RAE CRATON (units Agn-Agb) The rocks of the Rae craton in the Clyde River (OF 4432) and Blanchfield Lake (OF 4433) map areas comprise

as well as targeted re-examination of key areas across the entire project area

dominantly banded granodioritic to monzogranitic orthognelss, various granodioritic to monzogranitic and rare tonalitic plutons, and siliciclastic and minor mafic volcanic rocks of the Mary River Group.

The most common component of the orthogneiss (unit Agn) is biotite \pm hornblende granodiorite to monzogranite, with ndividual layers varying in thickness from several centimetres to several metres (Fig. 3). Biotite \pm hornblende tonalite layers are present locally; rare mafic layers range in composition from hornblende ± biotite diorite to clinopyroxenite A sample collected from a relatively homogeneous biotite monzogranite layer of orthogneiss in the vicinity of Dewar Lakes (Fig. 2) yielded a U-Pb crystallization age of 2827 + 8/-7 Ma (Wodicka et al., 2002). Similar ages have been reported from the southern margin of the Rae craton west of the map area (Bethune and Scammell, 1997, in press, and references therein), supporting the assertion that the southern domal, structural culminations in the Dewar Lakes area represent erosional windows that expose the southernmost extent of the Rae craton (Scott et al., 2002a; St-Onge

PLUTONIC ROCKS (units Atn, Agr-Agb)

The layering in the orthogneiss (unit Agn) is truncated in numerous localities by weakly- to moderately foliated plutoni rocks that range in composition from rare syenogranite to biotite monzogranite to less common tonalite. South of Generator Lake (OF 4432), a hornblende biotite mafic tonalite (unit Atn) is exposed structurally below an extensive panel of Mary River Group metasedimentary rocks (see below). Several hundred metres away from the supracrustal rocks, the tonalite is medium-grained and massive to weakly foliated (Fig. 20). Near the base of the metasedimentary rocks, the tonalite is an L-S tectonite characterized by a strong down-dip hornblende mineral lineation and abundant muscovite (Fig. 5). Immediately below the panel of Mary River Group rocks, the muscovite-defined fabric is most strongly developed, and is parallel to that developed in the overlying metasedimentary rocks. The progressive ncrease in the muscovite content of the tonalite may result from potassium metasomatism in proximity to the supracrustal rocks, in which case the tonalite could be viewed as the structural basement to the Mary River Group in his location. Alternatively, the muscovite-rich zone may represent a paleo-weathering horizon, suggesting that the contact with the overlying metasedimentary rocks may originally have been an unconformable depositional surface, and that the tonalite represents the stratigraphic basement to the Mary River Group. The highly strained nature of the

of the mafic tonalite has been collected for U-Pb geochronology. A variety of monzogranite plutons are present throughout the map area (Fig. 2). A sample of biotite allanite monzogranite (unit Agr) has been dated at 2738 + 25/-21 Ma, and a potassium feldspar megacrystic monzogranite (unit Agk) has an age of 2719 ± 4 Ma (Wodicka et al., 2002). Combined with U-Pb results presented by previous workers (Jackson et al., 1990; see Wodicka et al., 2002 for a compilation of results), the interval 2.74-2.72 Ga is emerging as a period of extensive felsic plutonism in the southern Rae craton. Dykes of hornblende-blottle \pm clinopyroxene gabbro (unit Agb) crosscut the blottle monzogranite (unit Agr) west of Blanchfield Lake (OF 4433). A similar dyke in the Eque Bay area has yielded an age of 2717 \pm 16/-13 Ma (Bethune

MARY RIVER GROUP (units Ama-AMp) Archean siliciclastic and volcanic rocks in the map area are assigned to the Mary River Group (Jackson, 1969, 2000;

Bethune and Scammell, 1997, in press), a succession of rocks that are found throughout northern Baffin Island and correlated with similar rocks (the Prince Albert Group) along strike to the southwest on the Melville Peninsula (Frisch, The most common siliciclastic rock type is brown-weathering psammite, with subordinate quartzite, semipelite and pelite (unit Amp). Development of coarse mica is ubiquitous, imparting a strong schistosity to all units. Coarse knots of sillimanite (+ quartz) are commonly developed in beds with more pelitic compositions (Fig. 6). Exposures of these rocks are commonly rust-stained, due to the oxidation of widespread, fine-grained disseminated pyrite. The brown weathering colour, coarse mica and generally rusty nature of these rocks readily distinguishes them from the siliciclastic rocks of the Piling Group (described below). Mafic rocks of the Mary River Group are relatively rare in the map area, in contrast with the area along strike to the

west (Bethune and Scammell, 1997, in press). Most are generally massive to moderately foliated amphibolite (unit

AMa), and characterized by hornblende + biotite; clinopyroxene was only rarely observed. At several localities,

irregular chloritic banding suggestive of pillow selvedges was observed; well-preserved pillows have been described along strike by Bethune and Scammell (1997, in press).

PALEOPROTEROZOIC PILING GROUP (units PPD-PPLs) Rocks of the Paleoproterozoic Piling Group are part of an extensive package (Jackson and Taylor, 1972) that is exposed as far east as the coast of West Greenland (Karrat Group; Taylor, 1982) and southwestward onto the Melville Peninsula (Penrhyn Group; Henderson, 1983). Because detailed descriptions of the constituent units and references to previous work have been previously presented (Scott et al., 2002a; St-Onge et al., 2002d), we focus here on newly

DEWAR LAKES FORMATION (unit PPD)

The stratigraphically lowest unit of the Dewar Lakes Formation (unit PPD) in the project area comprises white- to greyweathering quartzite and feldspathic quartzite, locally interbedded with psammite and pelite. The unconformable nature of the basal contact along much of the northern margin of the belt, as well as a three-part subdivision of the Dewar Lakes Formation, are described below. Of paramount importance to the understanding of the tectonic evolution and significance of the Piling Group is the recognition of its primary relationship to the underlying Rae craton. At numerous, widely spaced localities along the northern margin of the central belt (Fig. 2), clean exposures of the contact were identified and examined (Fig. 7). In each case, the basal contact of the Dewar Lakes Formation truncates lithological and structural elements in the underlying basement orthogneiss, but is conformable with overlying clastic material (commonly fine- to mediumgrained guartzite). Clear primary sedimentary features, such as planar bedding, were commonly observed, suggesting that the overall degree of strain at the contact is negligible. Consequently, we propose that much of the northern contact of the Piling Group against the underlying contiguous Archean rocks (Fig. 2) is an undisturbed unconformity. Recognition of this relationship clearly ties deposition of the Piling Group rocks to the Rae craton,

The lowest member of the Dewar Lakes Formation comprises thinly bedded quartzite, which is commonly rich in coarse muscovite and, grey- to pink-weathering psammite (Fig. 8); the thickest exposures of this member are up to ~50 metres. Overlying the lowest member are medium to thickly bedded quartzite (Fig. 9) with sillimanite and rare mica. This middle member of the Dewar Lakes Formation is by far the most widespread; exposures of tens to hundreds of metres in thickness are common. The upper member of the Dewar Lakes Formation is characterized by hin beds of quartzite to psammite and rusty semipelite to pelite (Fig. 10). The relatively higher proportion of pelitic

supporting earlier assertions (Morgan et al., 1975; Scott et al., 2002a) that these rocks collectively represent a

material in the upper member is consistent with a postulated deeper, more distal(?) depositional setting (Scott et al., Significant variations in overall thickness of the Dewar Lakes Formation have been identified throughout the project area (Fig. 2). At localities north of Flint Lake, the total thickness of the formation ranges from less than a metre to several tens of metres. In striking contrast, in excess of ~ 1000 m of quartzite is exposed at the west end of Flint Lake, and similar or greater thicknesses occur south of the McBeth River. In light of the recognition of an unconformity at the base of the formation, these variations need not be structural in nature, and are consequently interpreted as preserved primary features. One possible explanation is that the areas west of Flint Lake and south of the McBeth River were primary sedimentary depocentres, possibly deltas located at the mouths of rivers flowing off of the Rae

Detrital zircons from a quartzite sample from this formation have a bimodal age distribution: one population is Neoarchean, the other is 2.18–2.16 Ga (Henderson and Parrish, 1992). The Archean detritus is similar in age to known felsic plutonic rocks that form extensive parts of the adjacent Rae craton (Bethune and Scammell, 1997; Nodicka et al., 2002) and constitute the depositional basement to the Piling Group. The younger grain population brackets the timing of deposition of the entire Piling succession to younger than 2.16 Ga; the source of this detritus is enigmatic, as extensive exposures of rocks of this age are not known in northeast Laurentia (cf. Scott et al., 2002b, and references therein). Detrital zircons from additional samples of the Dewar Lakes Formation will be analysed by SHRIMP II ion microprobe (N. Wodicka, pers. comm. 2002) to further delineate the age and provenance of these rocks

FLINT LAKE FORMATION (unit PPF) White to grey weathering dolostone, marble (Fig. 11) and calcsilicate units of the Flint Lake Formation (unit PPF) are interlayered with minor to rare siliciclastic material (Corrigan et al., 2001; Scott et al., 2002a). The exposed stratigraphic thickness of this formation varies considerably across the project area (Fig. 2). The thickest exposures (500–1000m), near the west end of Flint Lake, thin toward both the east and west. In the central part of the area, nundreds of metres are preserved.

exposures are commonly less than several tens of meters, but can be traced along strike for hundreds of metres. Relatively thick exposures are found in the northeast part of the area (OF 4432), where thicknesses of tens to The southernmost exposures, typically thin calc-silicate and marble beds collectively only several metres thick, are no farther than \sim 40–50 km from the northern exposures of continuous Rae craton. We suggest that the Flint Lake

Formation originally formed a relatively narrow belt $(75–100\,\mathrm{km})$ adjacent and parallel to the southern edge of the Rae craton, with significant along-strike variation in primary thickness. BRAVO LAKE FORMATION (unit PPBm) In the southern part of the map area (Fig. 2), mafic volcanic rocks as well as mafic and ultramafic sills interbedded with

sedimentary rocks are assigned to the Bravo Lake Formation (see also Stacey and Pattison, 2003). The stratigraphic base of the formation occurs below the mafic volcanic rocks that conformably overlie quartzite, psammite and rusty semipelite of the upper Dewar Lakes Formation (Scott et al., 2002a; St-Onge et al., 2002b, c). In the northeastern part of the map area (OF 4432), the northernmost exposures of the Bravo Lake Formation (unit PPBm) are relatively thin sequences (< 100 m) of thin flows (or sills) interbedded with (or intruded into) siliciclastic rocks of the upper member of Because rocks of both the Flint Lake and Bravo Lake formations conformably overlie the siliciclastic rocks of the upper Dewar Lakes Formation, we suggest that these lithologically distinct formations occupy the same stratigraphic position in the overall Piling Group succession. This is further supported by observations (see next paragraph) that oth the carbonate and mafic volcanic rocks are directly overlain by rusty-weathering graphitic pelite and semipelite

ASTARTE RIVER FORMATION (unit PPA) This formation (unit PPA) comprises dark, rusty-weathering graphitic pelite and semipelite with minor sulphide-facies

ascribed to the Astarte River Formation.

throughout. Biotite and graphite are characteristic in the lowest metamorphic grade rocks, whereas sillimanite and muscovite are present in the highest-grade rocks. The maximum overall thickness of this formation is estimated to be on the order of 500 m in the northern part of the map area, and pinching out in the vicinity of Dewar Lakes (Fig. 2) in the south (Scott et al., 2002a; St-Onge et al., 2002d). In the vicinity of Flint Lake (Fig. 2), rusty graphitic pelites of the Astarte River Formation are in sharp, conformable contact with the underlying marbles of the Flint Lake Formation (Scott et al., 2002a). Further south (east of Foley sland: Fig. 2) and to the east and southeast (OF 4432), the Astarte River rocks are deposited directly on psammite and rusty semipelite of the upper member of the Dewar Lakes formation (Fig. 13). In the southernmost parts of the area, the rusty graphitic pelites overlie mafic volcanic rocks of the Bravo Lakes Formation (Scott et al., 2002a). The dramatic change in the nature of the stratigraphic units underlying the Astarte River Formation further emphasizes the acrossstrike variation in the paleogeographic distribution of the lowest units of the Piling Group.

LONGSTAFF BLUFF FORMATION (units PPLa-PPLs)

The Longstaff Bluff Formation (units PPLa-PPLs) comprises a relatively homogeneous package of distinctly eldspathic psammite, subordinate semipelite and rare pelite turbidites (Scott et al., 2002a). Bedding thickness is typically 10 to 50 cm, but can be as thick as 2 to 3 metres in extreme cases. Graded bedding the thinner beds (Fig. 14); the majority of such observations indicate that the succession is upright, with overturned peds found only on oversteepened fold limbs (St-Onge et al., 2002d) (OF 4432, OF 4433). Calcsilicate pods, up to several tens of centimetres in diameter, are ubiquitous throughout the psammitic beds of this formation and interpreted as metamorphosed carbonate concretions (Fig. 15). Elongate undulose features on bedding planes that are interpreted as ripple marks (Fig. 16) have been observed at only a few locations and are typically oriented roughly

The Longstaff Bluff Formation turbidites abruptly overlie the Astarte River Formation in the north (Fig. 17), the Bravo Lake Formation in the south-central part of the area, and the Dewar Lakes formation quartzites in the south (Fig. 2). Neodymium isotopic analysis (Johns, 2002) has demonstrated that these rocks have a distinctly younger

PALEOPROTEROZOIC GRANITES (unit Pgr) Plutonic rocks intrusive both into the Paleoproterozoic Piling Group (Fig. 18) and units of the Archean Rae craton, comprise syenogranite to monzogranite dykes, sills and irregularly-shaped plutonic masses (OF 4432, OF 4433) that commonly contain muscovite, garnet and biotite (unit Pgr). The granites are weakly to non-foliated, pink to white and coarse-grained to pegmatitic. These intrusions occur throughout the map area, and were emplaced over a time interval ranging from the thermal peak of metamorphism to the latest thick-skinned folding events (N. Wodicka, pers.

NEOPROTEROZOIC DIABASE DYKES (unit NFdb) Fresh, undeformed diabase dykes (unit NFdb) of the northwest-trending 723 + 4/-2 Ma (Heaman et al. 1992) Franklin swarm are widespread across the map area. These vertical dykes range in width from several metres to hundreds of netres; the thickest dykes can be traced for tens of kilometres. Locally they anastomose. Dykelets a few metres in width are locally observed branching from the larger intrusions. The dykes generally exhibit a brown-weathered

The Archean tectonothermal history of the Rae craton has been described in detail by Bethune and Scammell (1997

DEFORMATION AND METAMORPHISM

surface and the larger ones, a rectangular joint pattern.

in press) based on work in the Eque Bay area which is immediately to the west of the current project area. Overviews of the area north of Flint Lake have recently been presented (Corrigan et al., 2001; St-Onge et al., 2001a, b). Tight, intrafolial isoclinal folds of bedding in siliciclastic rocks of the Dewar Lakes Formation represent evidence of the earliest Paleoproterozoic deformation in the map area (D_{1B}). In the vicinity of Dewar Lakes (Fig. 2), the contact between the Archean orthogneisses and overlying quartities is highly strained, and commonly a well-developed, orthwest-trending mineral lineation is present (St-Onge et al., 2002c). The underlying orthogneisses are general recrystallized, with planar, contact-parallel fabric development increasing toward the contact with the overlying quartzites. Shear-sense indicators are rare, making it difficult to systematically document the sense of displacement ectonic imbrication of the lowest formations of the Piling Group in the Flint Lake area (St-Onge et al., 2001b) and structural juxtaposition of rocks of the Dewar Lakes and Bravo Lake formations onto Longstaff Bluff Formation rbidites in the southern part of the area (St-Onge et al., 2002b, c; Fig. 2), are both interpreted to have occurred during D_{1P}) (Scott et al., 2002a). Macroscopic, east-northeast-trending folds of Piling Group strata, often transected by steep cleavage striking 10 degrees clockwise from the strike of bedding, are also ascribed to D.... The axes of these folds typically have shallow plunges, and axial surfaces that vary from upright to slightly inclined. The fold closures Map-scale, northeast-trending folds of Piling Group rocks and underlying basement units occur throughout the central Baffin Island area (Fig. 2) and have been ascribed to Dap. At several localities, orthogneisses of assumed Archean age have been overthrust northward onto rocks of the lower Piling group (Fig. 4) and interpreted as a consequence of overtightening of D_{2P} folds. The basement-involved, thick-skinned D_{2P} folds are refolded (D_{3P}) t northwest- to northeast-trending upright cross folds that generate the overall dome-and-basin geometry of the map area (Fig. 2) and are responsible for the basement-cored domes that occur in the Dewar Lakes (St-Onge et al., 2002c),

McBeth River (OF 4432) and Blanchfield Lake (OF 4433) areas. Mapping in the eastern part of the map area has allowed documentation of the closure of metamorphic mineral zones, essentially outlining a "thermal trough" roughly centered on the belt of Piling group rocks (OF 4432). Mineral assemblages in pelitic rocks progressively increase in metamorphic grade in a broadly concentric pattern (St-Onge et al., 2002d) (OF 4432, OF 4433). As illustrated by rocks of pelitic composition, assemblages progress from muscovite, biotite muscovite, biotite muscovite cordierite + andalusite (Fig. 21), locally muscovite-sillimanite, then K-feldspar sillimanite melt, and ultimately garnet cordierite K-feldspar melt (in the southern portion of the project area). The present, complex geometry of the mineral zones is interpreted as a consequence of the interference between the D_{ap} and D_{ap} fold events.

ECONOMIC GEOLOGY The Paleoproterozoic rocks of the project area hold elevated potential for a variety of mineral deposit types. These include Mississippi Valley-type Zn in the relatively thick carbonate units (Flint Lake Formation), and sedimentary exhalative (SEDEX) base metal deposits in the overlying graphitic pelite (Astarte River Formation). The mafic and ultramafic rocks of the Bravo Lake Formation are exploration targets for Ni-Cu-PGE magmatic sulphide deposits Upper-amphibolite facies siliciclastic rocks of the upper Dewar Lakes Formation have potential for Broken Hill-type

REFERENCES

p-Zn-Ag deposits. In the northeastern part of the area, syenogranitic pegmatites that intrude the lower part of the

Deformation history and metamorphism of a synformal depression, Longstaff Bluff Formation metaturbidites, central Baffin Island, Nunavut; Geological Survey of Canada, Current Research 2002-

Bethune, K.M. and Scammell, R.J. Geology, Koch Island area, District of Franklin, Northwest Territories, (Part of NTS 37 C), Geological Survey of Canada, Open File 3391, scale 1:50 000. Structure and metamorphism in the Eqe Bay area, north-central Baffin Island: distinction between Archean and Paleoproterozoic tectonism and evolution of the Isortoq fault zone. Canadian Journal of

Regional geochemical reconnaissance, an introduction to the interpretation of data from central Baffin Island, District of Franklin; Geological Survey of Canada, Paper 86-10, 22p.

Corrigan, D., St-Onge, M.R., and Scott, D.J. Geology of the northern margin of the Trans-Hudson Orogen (Foxe Fold Belt), central Baffin Island, Nunavut. Geological Survey of Canada, Current Research 2001-C23, 17p.

Surficial geology, Nadluarjuq Lake, Nunavut. Geological Survey of Canada, Open File 4287, scale 1:100 Evans, S., Jones, A.G., Spratt, J., and Katsube, J.,

Central Baffin Electromagnetic Experiment (CBEX2) - Phase 2; Geological Survey of Canada, Current Precambrian geology of the Prince Albert Hills, western Melville Peninsula, Northwest Territories; Geological Survey of Canada, Bulletin 346, 70 p.

Friske, P.W.B., Day, S.J.A., McCurdy, M.W., and Durham, C.C. Reanalysis of 1775 lake sediments from surveys on central Baffin Island, Nunavut (Northwest Territories), parts of NTS 27 B, 27 C, 37 A, and 37 D; Geological Survey of Canada, Open File 3716, scale 1:300 000.

Heaman, L.M., LeCheminant, A.N., and Rainbird, R.H. Nature and timing of the Franklin igneous events, Canada: Implications for Late Proterozoic mantle plume and the breakup of Laurentia. Earth and Planetary Science Letters, v. 109, p. 117–131. Henderson, J.R.

1983: Structure and metamorphism of the Aphebian Penrhyn Group and its Archean basement complex in the Lyon Inlet area. Melville Peninsula, District of Franklin; Geological Survey of Canada, Bulletin 324, 50 p.

Geochronology and structural geology of Early Proterozoic Foxe-Rinkian Orogen, Baffin Island, NWT;

Geology of the Dewar Lakes area, central Baffin Island, District of Franklin, N.W.T. (parts of 27 B and 37 A); gical Survey of Canada, Open File 2924, scale 1:100 000 Henderson, J.R., Grocott, J., Henderson, M.N., Falardeau, F., and Heijke, P.

Survey of Canada, Paper 88-1C, p. 101-108. Henderson, J.R., Grocott, J., Henderson, M.N., and Perreault, S. Tectonic history of the Lower Proterozoic Foxe-Rinkian Belt in central Baffin Island, N.W.T., in Current

Results of field work near Dewar Lakes, Baffin Island, N.W.T.; in Current Research, Part C; Geological

United plates of America, the birth of a craton: Early Proterozoic assembly and growth of Laurentia.

Research, Part C, Geological Survey of Canada, Paper 89-1C, p. 186-187.

Annual Reviews of Earth and Planetary Sciences, v. 16: 543-603. Reconnaissance of north-central Baffin Island: in Report of Activities, Part A; Geological Survey of Canada, Paper 69-1, pt. A. p. 171-176. McBeth Gneiss Dome and associated Piling Group, central Baffin Island, District of Franklin; in Rubidium-Strontium Isotopic Age Studies, Report 2, (ed.) R.K. Wanless and W.D. Loveridge; Geological Survey of

Geology, Clyde River, District of Franklin, Northwest Territories; Geological Survey of Canada, Map 1582A, with expanded legend and brief marginal notes, scale 1:250 000. Geology of the Clyde-Cockburn land map area, north-central Baffin Island, Nunavut. Geological Survey of

Precambrian metamorphism on Baffin and Bylot Islands; in Metamorphism in the Canadian Shield (ed.) J.A. Fraser and W.W. Heywood; Geological Survey of Canada, Paper 78-10, p. 249-267. Jackson, G.D. and Taylor, F.C.

Correlation of major Aphebian rock units in the northeastern Canadian Shield. Canadian Journal of Earth Sciences, v. 9, p. 1650-1669. Jackson, G.D., Hunt, P.A., Loveridge, W.D., and Parrish, R.R.

Reconnaissance geochronology of Baffin Island, N.W.T.; in Radiogenic Age and Isotopic Studies: Report 3, Geological Survey of Canada, paper 89-2, p. 123-148.

Nd Isotope Constraints on the Provenance and Tectonic Evolution of the Piling Group Metasedimentary Rocks, Baffin Island, Nunavut, Unpublished B.Sc. thesis, University of Saskatchewan, Saskatoon, 61 p. The Trans-Hudson Orogen: extent, subdivisions, and problems; in The Early Proterozoic Trans-Hudson Orogen of North America (eds.) J.F. Lewry and M.R. Stauffer; Geological Association of Canada, Special

Lake Gillian, District of Franklin, Northwest Territories; Geological Survey of Canada, Map 1560A, scale

Morgan, W.C., Bourne, J., Herd, R.K., Pickett, J.W., and Tippett, C.R. Geology of the Foxe Fold Belt, Baffin Island, District of Franklin; in Report of Activities, Part A; Geological Survey of Canada, Paper 75-1A, p. 343–347. Stratigraphy, structure and metamorphism of the west half of the Foxe Fold Belt, Baffin Island; in Report of

Activities, Part A; Geological Survey of Canada, Paper 76-1A, p. 387-391. Scott, D.J., St-Onge, M.R., and Corrigan, D. 2002a: Geology of the Paleoproterozoic Piling Group and underlying Archean gneisses, central Baffin Island (NTS 37 A, 27 B); Geological Survey of Canada, Current Research 2002-C17, 10 p.

Geology of the Archean Rae craton and Mary River Group, and of the Paleoproterozoic Piling Group, entral Baffin Island, Nunavut; Geological Survey of Canada, Current Research 2003-C26. Scott, D.J., Stern, R.A., St-Onge, M.R., and McMullen, S.M. U-Pb geochronology of detrital zircons in metasedimentary rocks from southern Baffin Island: implications for the Paleoproterozoic tectonic evolution of Northeastern Laurentia. Canadian Journal of Earth

Teleseismic investigations of the lithosphere beneath central Baffin Island; Geological Survey of Canada,

Stacev. J. and Pattison, D.R.M. The stratigraphy, structure, and petrology of a representative klippe of the Bravo Lake Formation, central Baffin Island, Nunavut; Geological Survey of Canada, Current Research 2002-C13.

2001a: Geology, MacDonald River, Nunavut; Geological Survey of Canada, Open File 3958, scale 1:100 000. 2001b: Geology, Flint Lake, Nunavut; Geological Survey of Canada, Open File 3959, scale 1:100 000. Geology, North Tweedsmuir Island, Nunavut; Geological Survey of Canada, Open File 4199, scale 1:100 2002b: Geology, Straits Bay, Nunavut; Geological Survey of Canada, Open File 4200, scale 1:100 000.

2002c: Geology, Dewar Lakes, Nunavut; Geological Survey of Canada, Open File 4201, scale 1:100 000. 2002d: Geology, Inuksulik Lake, Nunavut; Geological Survey of Canada, Open File 4317, scale 1:250 000. Precambrian Geology of the Canadian North Borderlands; in Geology of the North Atlantic Borderlands, (ed.) J.W. Kerr, A.J. Fergusson, and L.C. Machan; Canadian Society of Petroleum Geologists, Memoir 7, p

Tippett, C.R. 1980: A geological cross-section through the southern margin of the Foxe Fold Belt, Baffin Island, Arctic Canada, and its relevance to the tectonic evolution of the northeastern Churchill Province; PhD. thesis, Queen's University, Kingston, Ontario, 409 p. Geology of a Transect Through the southern Margin of the Foxe Fold Belt, Central Baffin Island, District of Franklin; Geological Survey of Canada, Open File 1110; 77 pages.

Wheeler, J.O., Hoffman, P.F., Card, K.D., Davidson, A., Sanford, B.V., Okulitch, A.V., and Roest, W.R. Geological Map of Canada; Geological Survey of Canada, Map 1860A, scale 1:5 000 000. Wodicka, N., St-Onge, M.R., Scott, D.J., and Corrigan, D.

Preliminary report on the U-Pb geochronology of the northern margin of the Trans-Hudson Orogen, central Baffin Island, Nunavut; Geological Survey of Canada, Current Research 2002-F7, 12 p.

provenance than those of the Dewar Lakes formation. Detrital zircons from samples of this formation will be analysed by SHRIMP II ion microprobe (N. Wodicka, pers. comm. 2002) to characterize the provenance of and bracket the timing of deposition. We tentatively interpret the rocks of the Longstaff Bluff formation as a molasse derived through




Figure 1. Geological map of northeastern Canada modified after Wheeler et al. (1996) outlining the surface extent of the Trans-Hudson Orogen. Red box outlines location and surface extent of 2000-2002 mapping project in central Baffin







Figure 2. Generalized bedrock geology of the project area, compiled from St-Onge et al. (2001a, b; 2002a, b, c), OF 4432 and OF 4433.























Geology by M.R. St-Onge, D. Corrigan, F. Berniolles, K. Dubach, S. Modeland, J. Robillard. J. Stacey, J. Chakungal, S. Gagné, C. Hogue, C. Huerbert, A. Simmons, E. de Kemp, and D. Snyder (2000), M.R. St-Onge, D. Corrigan, M. Allan, F. Berniolles, K. Dubach,

Geological compilation by M.R. St-Onge, D.J. Scott, and D. Corrigan (2002)

J. Gladstone, S. Johns, R. Keiser, B. Sharpe, J. Stacey, and N. Wodicka (2002),

Geological Survey of Canada; D.J. Scott (2000, 2002), Canada-Nunavut Geoscience Office;

D.M. Carmichael and H. Helmstaedt (2000), Queen"s University; D. Francis (2000),

McGill University

Digital cartography by E. Everett, Earth Sciences Sector Information Division (ESS Info)

This map was produced from processes in conformance with the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001: 2000 standards

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada



igure 16. Ripple marks in a sandy bed of relatively thin-bedded

urbidites, Longstaff Bluff Formation (unit PPLc), McBeth River area.

Pen is 14 cm long. Location shown on OF 4432. (GSC 2002-806).

Figure 7. Basal quartzite of the Dewar Lakes Formation (unit PPD

long. Location shown on OF 4432. (GSC 2002-795).

unconformably overlying layered monzogranite-granodiorite




Figure 17. Ledge-forming psammite semipelite turbidites (left) of the

Longstaff Bluff Formation (unit PPLs) overlying rusty pelite of the Astarte

River Formation (unit PPA), southeast of the McBeth River. Backpack is

55 cm tall. View to the northeast. Location shown on OF 4432. (GSC

the Dewar Lakes Formation (unit PPD), southeast end of Flint Lake.

Hammer is 35 cm long. Location shown on OF 4433. (GSC 2002-796).






Figure 18. Garnet-muscovite leucocratic monzogranite (unit Pg


intrusive into calcareous marble of the Flint Lake Formation (unit PP)

4433. (GSC 2002-811).

MacDonald River area. Hammer is 35 cm long. Location shown on OF










Figure 15. Psammite-pelite turbidites with concretions (light coloured oblate domains), Longstaff Bluff Formation (unit PPLs), southeast of the McBeth River. Pen is 14 cm long. Location shown on OF 4432. (GSC



PPLs). Pen is 14 cm long. Location shown on OF 4432. (GSC 2002-808).

Logistical support provided by the Polar Continental Shelf Project as part of its mandate to promote scientific research in the Canadian North. PCSP 005-02.

OPEN FILE Open files are products DOSSIER PUBLIC that have not gone through the GSC formal Les dossiers publics son GEOLOGICAL SURVEY OF CANADA
OMMISSION GÉOLOGIQUE DU CANADA
pas été soumis au 2003 FEUILLET 2 DE 2

Sheet 2 of 2, Legend, descriptive notes, references, and figures Recommended citation: St-Onge, M.R., Scott, D.J., and Corrigan, D.

Canada, Open File 4433, scale 1:100 000.

2003: Blanchfield Lake, Baffin Island, Nunavut; Geological Survey of