Natural Resources Canada / Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA

NATMAP
CARTNAT
Canada's National Geoscience Mapping Program
Le Programme national de cartographie géoscientifique du Canada

DESCRIPTIVE NOTES

INTRODUCTION

The accompanying map presents the preliminary geology of the Meadowbank gold deposit area, examined as part of the recent Woodburn mapping project of the Western Churchill Natmap program. The map is a product of 1:5 000 to 1:20 000 scale mapping on foot traverse by Wilkinson, Zaleski, and others as cited. Previous and recent studies encompassing the area include 1:5 000 to 1:100 000 scale maps of Ashton (1988), Henderson and Henderson (1994), Zaleski et al., (1997, 1999a), and Sherlock et al., (2001), and relevant reports/theses of Ashton (1981, 1982), Henderson et al., (1991), Armitage (1992 and references therein), Armitage et al., (1996), Kjarsgaard et al., (1997), Kerswill et al., (1998), Zaleski et al., (1999b, 2000) and Sherlock et al., (2001b). Early regional reconnaissance maps and reports covering parts of the area include Fraser (1988), and Schau (1983). Map compilation and interpretation by the authors also utilised archived data of Henderson (1990), the unpublished 1:20 000 scale map of Hughes and Legault (1991), and an unpublished report by Miller (1997).

The centre of the map area is situated ~80 km north of Baker Lake, Nunavut, in the Rae domain of the Canadian Shield (Fig. 1). Access is by float plane or helicopter from Baker Lake in the summer or seasonally by winter road. Bedrock exposure is moderate to poor away from lakeshore and moderate to good on the shoreline and islands of Third Portage Lake. A summary of the geologic units is presented below, along with a description of structures, metamorphism, geochemistry, and mineral potential. No stratigraphic succession is implied by the order of Archean supracrustal units in the legend. All the supracrustal rock units have been metamorphosed but for simplicity the prefix 'meta' is omitted from the legend and following descriptions. Physiographic names, including minor lakes, are informal.

GENERAL GEOLOGY

The Meadowbank deposit area is underlain by metamorphosed and polydeformed supracrustal rocks of the Woodburn group and two large domains of granitoid rocks. The Woodburn group is part of a 3000 km long Neoarchean supracrustal sequence stretching from northern Saskatchewan to Baffin Island in the Rae domain (Fig 1). The group comprises Neoarchean volcano-sedimentary rocks that have been divided into at least three distinct packages: 1) ca. 2.72–2.71 Ga ultramafic through felsic volcanic and volcaniclastic rocks with associated iron-formation; 2) quartzites, phyllites, conglomerates, and quartz-feldspar porphyry of uncertain association and 3) arkosic wackes and pelites also with interlayered iron-formation (Davis and Zaleski, 1998; Zaleski et al., 2001). Rocks of the Woodburn Lake group are polydeformed and have been overprinted by strong regional Paleoproterozoic deformation, possibly related to the Taltson-Thelon and/or Trans-Hudson orogens (Fig. 2). Pronounced transposition of some units and fold interference results in a complex map pattern with multiple repetition of units. Low pressure-high temperature metamorphism that accompanied the second regional deformation event produced a zonation from mid-greenschist to amphibolite grade, north to south, across the map area.

The Meadowbank gold deposit, which includes the Third Portage, Goose Island, Bay zone, and North Portage zone, with an estimated combined resource of 2.1 million ounces (Table 1; Cumberland Resources Ltd. press release, June 07, 2001), is dominantly hosted in oxide-silicate facies iron-formations of the 2.71 Ga volcano-sedimentary package (Zaleski et al., 1999a, 2001; Sherlock et al., 2001b). Anomalous gold occurrences were first discovered in 1983 by Wollex Exploration (a subsidiary of Comaplex Minerals Corp.). Joint venture partner Asamera further delineated the prospect through mapping and diamond drilling between 1986–91, when it was optioned by Agnico Eagle, Hecla Mining, and Lucky Eagle Mines. During the mid-1990's, partners Comaplex Minerals and Cumberland Resources Ltd. considerably expanded the known resource. Cumberland Resources, the sole owner since 1998, is presently pursuing additional gold prospects northeast of Tern Lake.

THE WOODBURN GROUP

Volcanic and related rocks

The oldest supracrustal rocks of the Meadowbank deposit area are ca. 2.71 Ga volcanic, reworked volcaniclastic and related sedimentary rocks of the Woodburn group (Davis and Zaleski, 1998). Where exposure or data are insufficient to subdivide units or facies variability occurs on too small a scale, these rocks are grouped into an undifferentiated unit, found west and southwest of Third Portage Lake (unit Awv). A dearth of facing directions and strong transposition of units precludes establishing an internal stratigraphic order within this succession.

Felsic to intermediate reworked (unit Awvc) and primary (unit Awi) volcaniclastic rocks make up the largest component of the volcanic sequence. The reworked volcaniclastic rocks are bedded to laminated wackes and tuffaceous wackes, characterized by sand to granule sized quartz eyes, subrounded to rounded plagioclase crystals, and angular to rounded, lithic clasts in a fine-grained, clastic, variably recrystallized matrix of qtz-pl-msc-chl-crb-ep. The lithic clasts comprise (in decreasing order of abundance): polycrystalline quartz (quartzite and possible vein quartz), quartz-plagioclase porphyry, bt-msc-qtz-pl schist, and rare polycrystalline pl-fs-qtz aggregates. Bedding is defined by cm- to m-scale compositional and local grain-size grading. Tuffaceous wacke typically has a greater proportion of subrounded to euhedral plagioclase crystal clasts and quartz-plagioclase porphyry clasts and a compositionally more homogeneous, well-sorted matrix. Unit Awvc is interpreted as a reworked volcaniclastic deposit owing to the presence and abundance of quartz eyes, plagioclase crystals, and quartz-plagioclase porphyry clasts, the compositional similarity between the matrix and lithic clasts, and abundance of or locally observed gradational contacts with felsic to intermediate volcanic rocks.

Laminated volcaniclastic wackes with amphibole porphyroblasts (unit Awlam) are restricted to the southern part of the map area, above the amphibolite zone boundary. They are interpreted as metamorphic equivalents of unit Awvc with which they are contiguous, and can locally be difficult to distinguish from mafic intrusions where coarse, decussate amphiboles obliterate the layering.

Felsic to intermediate volcanic rocks (unit Awi) underlie the Dogleg Lake, Camp Island, and Horse Peninsula areas. Crystal and lapilli tuffs are the predominant component with subsidiary massive to layered quartz-plagioclase porphyry and probable volcanic flows. The tuffs are characterized by embayed and more rarely bipyramidal 0.5–5mm quartz crystals, broken and euhedral < 5mm plagioclase crystals, rare hornblende phenocrysts (pseudomorphed by actinolite) or dacite lapilli and crystals in a fine- to very fine-grained qtz-pl-msc ± chl or bt matrix. Rare, poorly sorted, volcanic breccias with > 25% blocks supported by an aphanitic qtz-pl-msc ± chl groundmass are noted on Horsehead Peninsula. Bedded, fine-grained tuffaceous volcanic rocks, characterized by cm- to dm-scale variations in the abundance of chlorite and muscovite and proportions of crystal clasts or lapilli, are distinguished from unit Awvc by the presence and abundance of delicate euhedral crystal clasts, lesser muscovite content, scarcity of lithic clasts, and a more homogeneous matrix. It is common for all layered or bedded units of this suite to have 2–7% disseminated and/or porphyroblastic carbonate. Occurrences of felsic volcanic rocks with up to 3% disseminated tourmaline in the matrix or minor mm-thick layers with 40% tourmaline are found throughout the unit.

Thick and continuous units of quartz-feldspar porphyry and felsic volcanic rocks (unit Awf) are differentiated in the Second Portage and Tuwawi Lake areas. These rocks are massive to layered and homogeneous, consisting of 2–5mm, euhedral quartz and plagioclase phenocrysts in an aphanitic to sugary msc-qtz-pl matrix. They have been dated at ca. 2.71 Ga (unpublished U-Pb zircon data of Davis and Sherlock) and may in part represent synvolcanic intrusions (see also Sherlock et al., 2001a, b).

Ultramafic volcanic and related rocks (unit Awkm) are present in two km-scale linear bands associated with iron-formation east of Third Portage lake, as a number of small, discontinuous lenses throughout the volcanic stratigraphy, and as generally narrow lenses at the contacts of iron-formation or quartzite. The unit consists predominantly of talc-chlorite-serpentine-magnesite, chlorite-actinolite ± magnesite or talc-tremolite-magnesite ± hornblende schists derived from variably carbonate-altered komatiitic and basaltic komatiite flows and breccia. Buff-cream dolomite-quartz-ankerite schists adjacent to the East BIF and just north of Third Portage Lake are strongly carbonated and silicified ultramafic rocks containing rare chromite, which serves to distinguish them from siliceous carbonaceous sediments. Rare primary textures include spinifex texture and polyhedral jointing. Layering on a cm- to dm-scale is defined by grain-size and proportion of amphibole and talc content.

Ultramafic to mafic volcaniclastic rocks (unit Awwk) generally form minor lenses within the ultramafic volcanic rocks, but particularly thick occurrences are found adjacent to quartzite at the north end of Third Portage Lake and between two quartzite panels extending from Third to Second Portage Lakes. These rocks consist of subrounded, 2–4 mm plagioclase crystals and quartz-feldspar porphyry lithic clasts in an ultramafic to mafic matrix of fine-grained chlorite and carbonate ± minor quartz or plagioclase. Rare m-scale beds with pebbles or granules of quartz-feldspar porphyry have been noted, suggesting the unit represents tuffaceous to reworked mafic-ultramafic volcaniclastic rocks.

Mafic volcanic flows and related rocks (unit Awm) typically form only minor lenses within the ultramafic rocks, but are particularly abundant along southwest Third Portage Lake, where they are associated with felsic-intermediate volcanic rocks. They are fine- to medium-grained, massive pl-act-chl schists or act-hbl amphibolites that locally display cm-scale amphibole layering. Coarser-grained flows can be difficult to distinguish from gabbros, as they have a coarsely recrystallized interstitial or relict diabasic texture. Primary structures are rare, although deformed pillows are noted on an island in southwest Third Portage Lake.

Metagabbroic rocks, including gabbro and peridotite, (unit Awa) are often associated with the mafic volcanic flows but also form mappable 250 m thick sills and several km-scale intrusions within the felsic-intermediate volcanic rocks. The gabbros are typically massive but locally have 0.5–1 cm, sub- to euhedral plagioclase phenocrysts and display a relict diabasic, ophitic or subophitic texture. The sill-like intrusions within unit Awvc have sharp contacts with adjacent host rocks and are interpreted as synvolcanic sills related to the mafic flows on the basis of a similar geochemical signature. A large composite mafic intrusion between southwest Tern Lake and Third Portage Lake has a crude zonation with a central peridotite core (now serpentinite) and a gabbro/diorite rim. This unit is distinguished from unit APgb on the basis of its plagioclase-phyric texture and geochemistry.

A variety of chemical and clastic sedimentary rocks are found as dm- to 10's m-scale beds or mappable units within the volcanic sequence. Oxide-facies, and lesser silicate or sulphide-facies banded iron-formation (unit Awif) are the most widespread, found as discontinuous thin (up to 10 m wide x 5 km long) interbeds throughout the sequence, but form thicker (10–100's m) units traceable for 10's of km at the contacts between ultramafic (unit Awkm) and felsic-intermediate volcanic or volcaniclastic rocks (units Awvc or Awms), quartz-muscovite schists (unit Awms) or quartzite (unit Aqz). The thickest units are known as the East, West, and Central BIF (Banded Iron-Formation) (Armitage et al., 1996), of which the latter hosts the Meadowbank deposit. Iron-formation characteristically contains cm- to m-scale chl ± grt-trm-cmm and qtz-msc-chl schist interbeds, interpreted as Fe-rich shales and volcanogenic wackes. Where less strained, beds in the iron-formation are defined by mm- to dm-thick, alternating quartz- or magnetite-grunerite-cummingtonite-rich bands. Sulphidic iron-formation, consisting of quartz-magnetite-pyrite-pyrrhotite ± arsenopyrite, is typically found where the iron-formation is gold-bearing.

A quartz-rich, muscovite ± chlorite schist (unit Awms) underlies much of the North Portage area and envelops the Central BIF. The schist is texturally homogeneous, fine-grained and only locally bedded, although well foliated. It contains rare 5mm polycrystalline quartz clasts, 3mm quartz eyes and trace tourmaline, kyanite and minor feldspar. Although superficially similar to foliated schists derived from unit Aqz, this unit is more compositionally heterogeneous, containing m- to 10's-m-scale layers and lenses of units Awvc and Awi. Given its relict volcaniclastic texture, interbedded volcanogenic units and K-enriched geochemical signature (Wilkinson, unpubl. data), it may represent an altered equivalent of units Awvc and Awi.

A thin but laterally continuous siliceous carbonate (unit Awsm) associated with iron-formation is present within laminated volcaniclastic wackes southeast of Third Portage Lake and felsic-intermediate volcanic rocks on Horsehead Peninsula. The marble consists of alternating cm-scale beds of fine-grained quartz dolomite, coarse dolomite, and fine- to medium-grained dolomite-calcite. The quartz-dolomite layers consist of a granular matrix of dolomite ± calcite and biotite with minor, rounded to subangular sand-size grains of quartz, and discrete lithic granules of 50% quartz-sand cemented by calcite-dolomite. The higher metamorphic grade horizon southeast of Third Portage Lake also contains tremolite, phlogopite, and rare humite. Zaleski et al., (2001) have interpreted the marble as a possible stromatolitic carbonate.

Greywacke-mudstones with minor quartz pebble conglomerate (unit Awg) underlie the northeast part of the map area. These rocks are fine- to medium-grained, well-bedded, chl-qtz-msc schists that are characterized by cm to dm-scale beds grading upward from greywacke to mudstone, and that locally display flame structures. Although the contact with adjacent reworked volcaniclastics is not directly observed, given the presence of dm-scale greywacke and conglomerate beds within unit Awvc, it may represent a more thoroughly reworked, equivalent facies of the volcaniclastic rocks.

Slate and phyllite (unit Awsl) forms several thin (< 100 m) but continuous horizons within the reworked and primary volcanic rocks. These chl-ctd ± ky bearing schists are bedded on a cm- to m-scale and are likely more prevalent than depicted, as they weather particularly recessively.

Amarulik wacke and associated rocks

A distinctive package of arkosic greywacke and related rocks, termed the Amarulik wackes (Zaleski et al., 1999a, b) occurs south of Third Portage Lake. The predominant unit is an arkosic greywacke consisting of < 3 mm quartz and feldspar crystal and lithic clasts in a coarse-grained bt-msc-qtz-fs matrix. The lithic clasts include quartzite, vein-quartz, quartz-plagioclase porphyry, and polycrystalline qtz-pl-fs aggregates. In contrast to unit Awg it is characterized by thick, dm- to m-scale, massive beds displaying only local grading, poor sorting, and more abundant feldspar crystal clasts. Oxide-facies iron-formation (unit Aaif) and pebble conglomerate (unit Aacg) are present as mappable, m- to km-scale interbeds.

The pebble conglomerate contains unsorted, granule- to cobble-sized lithic clasts in a massive quartzofeldspathic matrix. It is oligo- (quartz pebbles) to polymictic and bedded on a m-scale, containing clasts of (in decreasing order of abundance): quartzite, vein quartz, bt-msc-qtz-fs schist, and cherty oxide-facies iron-formation. Slate and siltstone (unit Aasl), as biotitic pelite, are interbedded with the thickest unit of iron-formation and underlie the eastern and western margins of the arkosic greywacke. Modified, cm-scale bedding is defined by varying proportions of biotite and compositional layers containing cordierite, staurolite, andalusite, garnet, or amphibole.

Immediately south of the map area the Amarulik arkosic greywacke forms a major regional package that is interpreted to be max. ca. 2.7 Ga (U-Pb data of Davis and Zaleski) but > 2.6 Ga (Roddick et al., 1992). In this area the Amarulik wackes also contain intermediate-mafic volcanic rocks not present in the Third Portage area, and are interpreted to be a reworked volcaniclastic deposit (Zaleski et al., 1999b). It is possible that it may be a facies equivalent of the 2.71 Ga volcanic succession.

Quartzite and associated rocks

The third major stratigraphic package of the Meadowbank map area consists largely of quartz-rich rocks; quartzites, slates and phyllites, and oligomictic to polymictic conglomerates that are considered the upper part of the Woodburn group. The quartzites themselves (unit APqz) are strongly recrystallized, massive to schistose, quartz arenites that locally contain heavy mineral layers with up to 7% magnetite-ilmenite-chromite. Primary beds (and rarely cross-beds) are noted at the cm- to m-scale, and compositional variations to more feldspathic arenite are locally noted. Minor muscovite is disseminated through the unit and defines bedforms but increases to 25% where quartzite is strongly sheared, particularly along the unit contacts and throughout the isolated quartzite body immediately north of Third Portage Lake.

Quartzite occurs as two major associations: 10's km thick units with interbedded slate and phyllite (unit APsl), such as that exposed northwest of Third Portage Lake, and thinner, up to 2 km thick, laterally continuous units intercalated with the volcanic rocks and Amarulik wackes (unit Aqz). Oligomictic quartz pebble conglomerate (units APqcg and Aqcg) is present in both associations; forming dm- to km-scale lenses interbedded with and underlying the thick quartzites and slates (as at Tuwawi Lake); and m- to 100's m-scale units at the basal contact of the thin quartzite units (above the Central BIF and along Tehek arm). The conglomerate consists of white quartzite and vein quartz clasts, with minor grey polycrystalline quartz and rare fuchsitic clasts in a micaceous quartz-rich matrix. Bedding is locally defined by grading in clast size and abundance.

Polymictic conglomerate associated with quartzite has been identified at several localities; between quartzite and the volcanic sequence on the northwest shore of Second Portage Lake, and between quartzite and the West BIF on the south part of Camp island. The occurrences are at shoreline and can be submerged depending on seasonal or other variations in water levels. The Second Portage conglomerate (unit APcg) contains pebbles, cobbles, and boulders up to 1m of (in decreasing order of abundance): white to buff quartzite and vein quartz, iron-formation, grey polycrystalline quartz, intermediate volcanic rocks, and granite. It is poorly sorted and locally transitional to oligomictic conglomerate, with which it can be bedded on a m- to 10's m-scale. The West BIF occurrence (unit Acg) is dominated by quartzite pebbles, polycrystalline quartz and iron-formation clasts, and occurs at an apparent conformable contact between Archean quartzite and ultramafic rocks.

The association of polymictic and oligomictic conglomerate with thick quartzite units has been interpreted as marking an unconformable contact with underlying 2.71 Ga volcanic rocks northwest of the map area (Zaleski et al., 2000). The lower contact of the conglomerate inlier at Tuwawi lake and the large thick quartzite body to the north with is similarly interpreted to be unconformable. The relative stratigraphic position of the thinner quartzite units is presently uncertain, however both structural (Tern Lake) and possibly conformable (Camp Island) contacts with the volcanic rocks are noted. As recent U-Pb geochronology has established that a thick quartzite-slate sequence at Whitehills Lake to south is Proterozoic (Pehrsson et al., 2002), the thick quartzites and related map units are designated as Archean or Proterozoic, whereas those quartzites apparently interbedded with the Archean volcanic rocks are interpreted as Archean.

PLUTONIC ROCKS

Plutonic rocks of both Archean and Paleoproterozoic age are interpreted to be present in the Meadowbank map area. The oldest unit associated with the plutonic rocks is the Tern Lake porphyry (unit Afp), a layered, quartz and K-feldspar-phyric porphyry dated at ca. 2.63 Ga (Davis and Zaleski, unpubl. data), situated just south of Tern Lake. The porphyry is characterized by embayed to broken blue quartz crystals, euhedral microcline (0.5–5mm, locally perthitic textured) and lesser plagioclase phenocrysts in a fine- to medium-grained, quartzofeldspathic matrix. Minor biotite and relict hornblende phenocrysts (to 2 mm) have been noted and ilmenite, chalcopyrite, pyrite, and magnetite are disseminated in the matrix.

The Tern Lake porphyry is variably sheared, typically more highly strained on its margins, and shows considerable textural and grain size variation along strike. At its southwestern end it is a strongly layered, 'tuffaceous' looking rock with 'beds' outlined on 0.5–10 cm scale by the abundance of phenocrysts (sparse to 25% or locally 50%). At its northeastern end it is a well foliated to locally mylonitic, K-feldspar and plagioclase porphyritic (5–15 mm), biotite-hornblende granodiorite (unit Afpg) with up to 15% disseminated sulphides. Fine-grained porphyry can be interlayered at the 10's m-scale with coarser-grained granite. Based on the overall texture and association with coarser-grained granitic rocks, the Tern Lake porphyry likely represents a deformed high-level, locally extrusive, subvolcanic intrusion.

Systematic mapping was not conducted in the large granitoid domains that underlie the eastern and western margins of the map area, consequently only an overview of these rocks is given here. Preliminary examination suggests that the domains are composite, with at least local subphases bearing fluorite. Further subdivision is likely possible. The predominant map unit is a pink, inequigranular to K-feldspar megacrystic (1–2 cm) monzogranite to granite (unit Agt) with biotite and/or hornblende, epidote, and magnetite in a variably recrystallized groundmass. In contrast to parts of the Tern Lake granodiorite (unit Afpg), the granite is relatively leucocratic, with typically less than 7% mafic phases, although these can be highly chloritized. It is commonly well foliated, with foliation defined by alignment of megacrysts, quartz grain-shapes, and mafic minerals. The margin of the western domain is characterized by a 100's m wide zone of intense, locally mylonitic, foliation and a sheeted zone of abundant m to 100's m wide, highly sheared intrusions in adjacent country rocks. The eastern domain is less strained overall along its margin, although highly deformed granite sheets are also present along its contact zone. A composite U-Pb age of 2612 ± 4 Ma (Roddick et al., 1992) has been obtained from both domains.

Minor granitic to syenitic stocks and dykes of suspected Paleoproterozoic age (units PNgt, PHgt) are found northwest and east of Third Portage Lake, where they intrude Woodburn group volcanic rocks and Archean granite, respectively. The Nueltin granites (unit PNgt) are coarse-grained, dark red, leucocratic and are vuggy textured locally, and have similar composition, accessory mineralogy, and geochemistry as the 1.76 Ga Nueltin granite suite (Petersen et al., in press). A prominent, subcircular magnetic anomaly within the eastern Archean granite just north of a unit PNgt dyke, may be a related body. Similar small stocks, also interpreted as Nueltin granites, are found just northwest of the map area at Pipedream Lake (Zaleski et al., 1997). A small body of leucocratic, inequigranular, biotite-garnet alkalic to syenitic granite (unit PHgt) occurs just north of Tuwawi Lake. This granite is similar in composition and texture to the ca. 1.835 Ga Paleoproterozoic Hudson granites (Petersen et al., in press), a suite that occurs as abundant dykes immediately south of the map area, and as large plutons farther south towards Baker Lake.

Late crosscutting lamprophyre dykes, interpreted to be related to minette flows of the Baker Lake group (Rainbird et al., 2001), are observed in drill core (Sherlock et al., 2001a, b) and are reported to occur in outcrop (Armitage, 1992), although they but were not noted in the present mapping.

Mafic to felsic intrusions of unknown age form two distinct suites in the map area. Gabbro, quartz leucogabbro and monzodiorite (unit APgb) form small stocks southwest and east of Tern Lake and a larger composite body south of Second Portage Lake. The latter can be difficult to distinguish from recrystallized amphibole-rich gneisses of the Woodburn group. These bodies are associated with localised magnetic anomalies, a relict ophitic texture and geochemically distinct from unit Awa. Northwest of Pipedream Lake, a gabbro body of similar geochemistry and texture intrudes the 2.6 Ga granite (Ashton, 1988), allowing the possibility that unit APgb is Paleoproterozoic in age.

Biotite-muscovite granite is present in the map area as dykes and sheets and several minor stocks that intrude the volcanic sequence (unit APgm). These leucocratic, inequigranular to weakly K-feldspar porphyritic intrusions are characterized by 5–7% biotite and lesser muscovite in a foliated to locally mylonitic, recrystallized groundmass. They are not observed to cut the other intrusive rocks, consequently their relative age is uncertain.

STRUCTURE

Four phases of ductile deformation affect all rocks of the Woodburn group in the Meadowbank deposit area, and are correlated with structures identified regionally (Pehrsson et al., 2000). The sequence elaborated here builds on that established by Henderson et al. (1991) and Ashton (1988).

The first recognised deformation comprises outcrop- and thin section-scale evidence for two sets of pre-S_1 isoclinal folds and foliations that are found only within the 2.7 Ga Woodburn group volcanic and related rocks. These local observations and the significantly greater intensity of overall strain in this sequence suggest that the 2.7 Ga volcanic rocks were likely deformed prior to deposition of the thick quartzite sequence.

The earliest regional deformation event, D_1, involved development of macroscopic, shallowly-plunging, isoclinal folds (F_1), bedding subparallel foliation (S_1), and mineral and clast elongation lineations (L_1). S_1, axial planar to F_1 folds of bedding and quartz veins, is defined by phyllosilicate and chlorite alignment and quartz grain shape fabrics in the supracrustal rocks, and phenocryst elongation and mineral shape fabrics in all plutonic rocks with the exception of units PNgt and PHgt dykes. S_1 can be the dominant foliation observed in outcrop (S_m) but is ubiquitously folded by mesoscale F_2 folds and variably overprinted by S_2, making delineation difficult in the field. The timing of D_1 is bracketed by intrusion of the 2.59–2.62 Ga granite suite and 1.835 Ga, the age of a post-D_2 pegmatite dyke (Roddick et al., 1992; Pehrsson et al., 2000). Variable, inhomogeneous development of S_1 in the granites (cf. Ashton, 1988; Davis and Zaleski, 1998) most likely reflects regional strain heterogeneity. If S_1 schistosity in the ca. 2.0 Ga Whitehills conglomerate (Davis and Zaleski, unpubl. data) is correlative with S_1 of the deposit area the regional D_1 deformation event may also be Paleoproterozoic in age.

The predominant structures of the Meadowbank area, those related to the second phase of regional deformation (D_2), comprise tight to isoclinal folds (F_2), axial planar foliations (S_2), crenulation or intersection lineations (L_2), and reverse faults or shear zones (Fig. 4a). S_2 varies from a differentiated crenulation to crenulation cleavage to a transposition foliation depending on host lithology and intensity of S_1 development. It locally obliterates bedding and S_1. The main regional foliation mapped in outcrop (S_m) is typically an S_1 or S_1/S_2 transposition foliation.

F_2 folds at all scales are shallowly doubly-plunging folds of S_1. North and west of Third Portage Lake they form a map-scale northwest-vergent fold set with moderately southeast-dipping axial surfaces (Fig. 4a). South of Third Portage Lake consistent S_2/S_1 asymmetry and change in S_1/S_m attitude define a regional F_2 synform, overturned to the west. North and east of Second Portage Lake F_2 axial surfaces and S_2 have variable orientations due to superposed F_4 folds (Fig. 4b,c). The regional northwest-vergent D_2 fold-fault belt deforms the Archean granites, and does not predate their intrusion as suggested by Henderson et al (1991).

A major NW-vergent D_2 thrust fault places 2.71 Ga Woodburn group volcanic rocks on the unconformably overlying quartzite sequence north of Third Portage Lake. This structure may link with a zone of D_2 high strain that extends through the deposit area and south through Second Portage Lake (Fig. 4d). This zone is characterized by increased tightness of F_2 folds, strong S_2/S_1 transposition fabrics and local mylonites, and has been termed the Third Portage fault zone by Hughes and Legault (1991). A similar zone of D_2 high strain follows the quartzite/BIF/komatiite contact at Camp Island and extends south of Third Portage Lake where a F_2 synformal fold of iron-formation is apparently truncated against its trace, suggesting the zone may be the locus of significant dislocation, and possibly a thrust fault.

The maximum age for D_2 deformation is 2.0 Ma, the maximum age of the Whitehills conglomerate (Davis and Zaleski, unpublished data), which carries the S_2 foliation. Like D_1, the lower bound for D_2 is 1835 ± 1 Ma, the age of a granite pegmatite dyke that crosscuts S_2 south of the deposit area (Roddick et al., 1992). The best estimate of the age of deformation is provided by U-Pb ages of ca. 1.8–1.9 Ga obtained from monazites within syn-D_2 staurolite porphyroblasts in rocks northwest of the map area (Berman and Pehrsson, unpubl. data). The similarity between the attitude and vergence of D_2 structures in the Woodburn group and the Amer fold-thrust belt (Patterson, 1986) has been previously noted by Ashton (1988), Davis and Zaleski (1998) and Zaleski et al., (2000).

D_3 structures consist of open to close F_3 folds of S_2/S_m with an associated L_3 crenulation lineation. F_3 chevron to similar-style folds of S_2 and L_2 have wavelengths on the order of decimetres to centimetres. The shallowly plunging to subhorizontal L_3 crenulation lineation, is locally subparallel to F_3 fold axes, is locally the dominant lineation in outcrop and crenulates minerals such as grunerite that are randomly oriented on S_2. F_3 folds define a consistently southeast-verging minor fold set with shallowly north- and northwest-dipping (< 40°) axial surfaces. These folds locally overprint structures in the deposit area and can be responsible for variable S_2 dips. S_3 foliations, defined by crenulations of S_2 and new growth of muscovite or chlorite, are typically only visible in thin section.

Mapping east of Whitehills Lake (Zaleski et al., 2000) indicates that D_3 structures overprint a suite of syeno-monzodiorite intrusions that are correlated, on the basis of composition and geochemistry, with dated, ca. 1835 Ma intrusions at MacQuoid Lake (Davis, unpublished data). This suggests that D_3 deformation postdates ca. 1840 Ma. A lower bound is provided by a ca. 1790 Ma Ar-Ar age on M_3 porphyroblastic biotite that is crenulated by S_4. Zaleski et al., (1999) have previously drawn attention to the nearly identical orientation, geometry and vergence of F_3 folds and D_3 structures in the Amer group.

D_1 through D_3 structures are folded about a map-scale, south-plunging F_4 antiform-synform pair, centred on the North Portage area (Fig. 1). Mesoscale D4 structures include upright, open to tight F_4 folds, local steep S_4 crenulation cleavage, and related L_4 crenulation lineation. F_4 folds are moderately to shallowly-plunging, with north-northeast trending, steeply dipping (> 60°) axial surfaces (Fig. 4c). Grunerite, amphibole and biotite porphyroblasts, which overgrow S_2, are locally crenulated or wrapped by S_4 and define the spaced L_4 crenulation lineation. Local quartz-carbonate veins that occur along the axial surface of F_4 folds have chloritic alteration haloes and new growth of amphibole on vein walls and S_4 cleavage.

The deposit area is dominated by type-2, mushroom fold interference patterns resulting from superposition of upright F_4 folds on inclined, tight F_2 folds. This is best expressed by the refolding pattern of the Central and East BIF's but is also noted in outcrop-scale fold interference patterns in unit Awms. Northeast-trending, southeast-dipping F_2 folds on the west limb of the major F_4 synform progressively change attitude to south-southeast-trending and southwest-dipping on the east limb. F_4 folds in the North Portage zone plunge moderately south and southwest, down the dip of F_2 limbs. In contrast, F_4 folds at the Bay zone to the south locally plunge moderately to steeply northeast, due to interference with northeast-dipping limbs of tight F_2 folds. D_4 deformation of the Meadowbank deposit area appears to only remobilize pre-existing mineralization, however, it may have facilitated mineability or exposure of the deposit due to the geometry of fold interference. The established $D_2/D_3/D_4$ succession and overprinting are consistent for the area from north Third Portage Lake to Goose and Camp Islands. Further south local outcrop evidence suggests a D_3 phase of deformation or the possibility that D_4 deformation was continuous with D_3.

The absolute upper limit on the age of D_4 deformation is ca. 1840 Ma, the age of the monzodiorite intrusive suite that is overprinted by D_3/D_4 interference folding in the Tehek Lake area (see also Zaleski et al., 2000). As noted above, biotite porphyroblasts with Ar-Ar ages of ca. 1790 Ma are crenulated by S_4. Depending on the cooling history of the area (a subject of continuing study), this may provide the maximum age of D_4. The absolute minimum age is thought to be ca. 1760 Ma, the age of the MacRae diabase dykes (Lecheminant, pers. comm.), however this type of deformation is unknown in the Baker Lake Group (Fig. 2).

METAMORPHISM

The map area preserves evidence of three distinct metamorphic events, designated M1 to M3. M1 regional metamorphism is outlined by low to mid-greenschist facies mineral assemblages that define S_1 fabrics. It occurred at unknown pressure and is constrained to predate development of D_2 structures ca. 1.8–1.9 Ga and post-date the 2.60 Ga granites that carry S_1 foliations (Fig. 2). As is the case with S_1, M1 may post-date deposition of the Paleoproterozoic Whitehills conglomerate if regional correlation of fabrics is established.

M2 regional metamorphism is characterized by mid-greenschist through amphibolite facies assemblages (summarised in Table 2) that are interpreted to be broadly coeval with D_2 deformation, ca. 1.8–1.9 Ga (Fig. 2; Berman, unpubl. data). Greenschist zone M2 phases such as chloritoid, kyanite, chlorite, and biotite are typically aligned in or wrapped by S_2, whereas amphibolite zone amphibole porphyroblasts, including staurolite, andalusite and garnet, both overgrow and are wrapped by S_2. Local carbonate and plagioclase porphyroblasts in units Awi and Awvc display curvilinear inclusion trails smoothly continuous with matrix S_2. This evidence and recrystallization of S_1 phyllosilicates that are crenulated by S_2 suggest that M2 regional metamorphism was broadly coeval with D_2 but that the thermal peak outlasted regional deformation.

Distribution of characteristic mineral assemblages defines three metamorphic mineral zones: mid-greenschist, upper greenschist or epidote amphibolite, and amphibolite (Fig. 5). The mid/upper-greenschist zone boundary trends southeast from Tuwawi Lake, across Third Portage Lake and east-northeast just north of the Third Portage deposit. The upper-greenschist/amphibolite zone boundary trends roughly east-west across Third Portage Lake, passing just north of Goose Island. The presence of kyanite after pyrophyllite in quartzite or phyllite and grunerite-chlorite in iron-formation constrain P-T conditions of the M2 mid-greenschist zone to ca. 400°C at min. 2.7 kbar. Assemblages of the upper-greenschist zone collectively suggest temperatures of ca. 450°C at low pressures. The occurrence of the assemblage bt-st-and-musc at ~120 m depth in drill core (L'Heureux, unpub. data) and quantitative thermobarometric data (Miller, unpub. report) yield amphibolite zone conditions of 3.0 kbar at ca. 549°C, for Goose Island, just south of the zone boundary. Local fo-tlc ± tr assemblages in ultramafic rocks and the maximum stability range of chloritoid-kyanite (also ca. 550°C) are consistent with these estimates. South of Third Portage Lake, humite in phlogopite marble suggests M2 temperatures locally attained 600–650°C. The trace of the upper-greenschist and amphibolite zone boundaries crosscut D_2 structural trends, possibly due to post-thickening thermal relaxation.

M3 metamorphism is defined by new growth of mineral phases that everywhere overgrow S_2 and locally S_3 fabrics, but are wrapped and/or crenulated by S_4. These phases, characteristically porphyroblastic bt, grt and cum or act, are patchily distributed across the mid-upper greenschist and amphibolite zone boundaries (Fig. 5). South of Third Portage Lake, distinct late chlorite and muscovite are widespread and similarly crenulated by S_4. Pressure-temperature estimates of ca. 450°C at 3.0 kbar have been obtained from M3 grt-chl and grt-bio assemblages in drill core at the Third Portage deposit (Armitage, 1992). Ar-Ar (Villeneuve and Kjarsgaard, unpub. data) and K-Ar (Armitage et al., 1996) ages of M3 biotite porphyroblasts in the deposit area are 1790 ± 12 and 1791 ± 32 Ma, respectively, indicating the time of cooling through the 300°C isotherm. It is possible that M3 metamorphism represents a local, thermal aureole effect related to Nueltin suite magmatism.

ECONOMIC GEOLOGY

The Meadowbank gold deposit comprises four deposit zones localised within and adjacent to iron-formation of the Central BIF (Fig. 4d). Gold occurs in iron-formation at its contact with komatiite or volcaniclastic sediments and only in those iron-formations associated with the ultramafic/volcaniclastic interface, small units within the volcaniclastics entirely or within the Amarulik wackes are uneconomic to barren. The deposits are concentrated in the noses and along the limbs of D_1 and D_2 isoclinal folds, and show multiple styles of sulfide mineralization (Pehrsson et al., 2000, Sherlock et al., 2000a, b), associated to varying degrees with prospective gold concentrations. These styles are grouped according to their relationship to the major deformation phases, pre-syn D_2, syn-D_2 and post-D_4.

Pre- to syn-D_2 mineralization most commonly comprises pyrrhotite, locally pyrite and rarely arsenopyrite and variable gold concentrated within S_1 and S_2 foliations, with or without magnetite. These sulphides are also found within or adjacent to quartz veins deformed in S_1 or S_2 foliations and veins that cut S_1 but are deformed by D_2 (Pehrsson et al., 2000). Quartz-epidote-sulphide veins from outside the deposits themselves define F_1/F_2 interference patterns at thin-section scale and are localised along shears parallel to S_2. Regional evidence for pre D_1 mineralization comes from disseminated pyrrhotite and chalcopyrite in units Awms, Awf and Awi that display grain shape fabrics elongate in S_1, and from graded pyrrhotite concentrations in volcaniclastic rocks at Goose Island. The latter texture was interpreted by Miller (1997) to represent syngenetic-mineralization. Pre-S_1/S_2 quartz-feldspar porphyry dykes in drill core at the deposit are associated with minor Cu-Au-Ag mineralization as are parts of unit Afpg (Kerswill et al., 1998), which is dated at ca. 2.63 Ga (Davis and Zaleski, unpublished data).

Syn-D_2 mineralization comprises secondary sulfides (pyrrhotite ± pyrite) and gold that occur in S_2 parallel ductile-brittle shears at the iron-formation-komatiite contact and internal to the iron-formation, and within patchy zones that cut earlier sulfide and magnetite layers in S_1 or S_2 (Pehrsson et al., 2000). The shears may be genetically related to the D_2 strain zone that follows the komatiite-bif and quartzite contact through the deposit area (Fig. 4d). The patchy textured sulphides are locally associated with in-situ brecciation, quartz veins and silicification in D_2 dilatant zones.

Regionally, minor pyrrhotite, ± pyrite and arsenopyrite are present as blebs and streaks along S_2 crenulation cleavages associated with mesoscale F_2 folds in the iron-formation and units Awms and Awvc. Minor syn-D_2 quartz-carbonate veins are present throughout the map area. These veins show two subsets; earlier veins that folded by F_2 folds but locally cut S_2, and later veins that are parallel to F_2 axial surfaces. In the komatiitic rocks they are associated with syn-M_2 coarsening of fe-chlorites, fluid infiltration, and carbonitization whereas in units Awvc, Awf and Awi they are associated with concentration/remobilization of pyrrhotite/pyrite and ilmenite/magnetite and coarsening of amphiboles. Quartz-carbonate-epidote ± allanite veins and carbonate-epidote-chlorite vienlets that are parallel to S_2/S_1 and crosscut S_1/S_m are associated with epidote and sericite alteration and concentration of pyrite that overgrows earlier deformed sulphides.

Syn-D_4 mineralization comprises coarse pyrite ± pyrrhotite are associated with quartz-carbonate veins parallel to the axial planes of minor F_4 folds and S_4 crenulation cleavages. These veins show distinct chloritic haloes and are only gold-rich where they cross the Meadowbank deposits and are barren where S_1 and S_2 foliations are gold-poor (Pehrsson et al., 2000). Remobilization of pre-D_4 sulphides parallel to S_4 is visible in thin section. Together these observations indicate that the veins and associated fluids did not introduce new sulphides or gold.

DISCUSSION

Stratigraphy and Structural Geometry

Any analysis of the structural evolution of the map area cannot be finalised without a more complete understanding of the stratigraphic relationships between the various successions. Whereas it appears that the Neoarchean volcanic and volcaniclastic rocks and thin quartzite units are likely part of a conformable stratigraphic succession, the thick quartzite-conglomerate-slate package north of Third Portage Lake is clearly deposited unconformably on the volcanic-dominated succession. This upper quartzite-dominated package is interpreted to be dated at 2.63 Ga, part of a sequence deposited unconformably on the 2.71 Ga komatiites, felsic-intermediate and related rocks of the Woodburn group (Zaleski et al., 2001). Pehrsson et al., (2002) have alternatively interpreted this as a maximum detrital age, and consider that this thick upper sequence could be Proterozoic in age, similar to the dated sequence at Whitehills Lake. Repetition of the quartzite as thinner units within the volcanic package and underlying the Amarulik wacke has been ascribed to structural interleaving by both folding and faulting (Ashton, 1988; Hughes and Legault, 1991; Henderson et al., 1991), although Ashton (1988) allowed that some quartzites could be part of the 2.71 stratigraphic package itself, an interpretation supported by Sherlock et al., (2000a, b). The Amarulik wackes themselves have previously been interpreted as volcanogenic rocks derived from and in part contemporaneous with the 2.71 volcanic sequence (Henderson and Henderson, 1994). Resolution of these stratigraphic questions awaits more detailed study.

Mineralization

Several models have been proposed for the genesis of the Meadowbank gold deposit. Miller (1997) emphasized the syngenetic nature of the sulphides for the Goose Island deposit. Armitage et al., (1996) suggested that gold mineralization was syn- to post-the 2.6 Ga granites and related to sulphidation-alteration reactions concentrated in the Central BIF during Proterozoic reactivation of Archean structures or imposition of new Proterozoic structures. Sherlock et al., (2001a, b) elaborated on this model, linking mineralization and alteration to progressive D_2/D_1 deformation and localisation of strain adjacent to the Central BIF ultramafic/iron-formation contact.

Data from this study suggests that the mineralization in the Central BIF is pre- to syn-D_2, largely ca. 1.8–1.9 Ga in age, consistent with the general models of Armitage and Sherlock. Given the evidence for high strain along the West BIF/komatiite/quartzite contact, additional influences are required to explain the localisation of economic mineralization in the Central BIF.

Inspection of the distribution of lithologies in the map area shows that unit Awms is spatially associated with the auriferous Central BIF and the Dogleg Lake synvolcanic porphyry (unit Awf), dated at ca. 2.7 Ga (Davis and Sherlock, unpublished data). Geochemical and textural evidence suggests that unit Awms is a sericite and chlorite altered rock of volcaniclastic unit Awvc protolith that was altered prior to D_1 deformation. Given this association and the local presence of anomalous tourmaline concentrations and pyrrhotite-pyrite in the unit Awf porphyry, we suggest that the distinct mineralogy of the Central BIF and distribution of the unit Awms schist may in part reflect syngenetic semi-regional alteration adjacent to 2.7 Ga porphyry. This alteration may have acted as a ground preparation event, making the Central BIF a preferential site for syn-D_2 mineralization over the West or East BIFs.

ACKNOWLEDGEMENTS

Cumberland Resources Ltd. is gratefully acknowledged for logistical support of the project, access to drill core samples and airborne geophysical data. Mike Thomas is thanked for producing vertical derivative magnetic maps. Amanda Chipoot is thanked for digitising. Ross Sherlock, Alan Armitage, Bill Davis, John Kerswill, Rob Berman, and Hamish Sandeman are thanked for stimulating discussions.

REFERENCES

Armitage, A.E.
1992: Geology and petrology of gold-bearing banded iron-formation, Keewatin District, Northwest Territories; unpublished M.Sc. Thesis Laurentian University, Thunder Bay, 162p.

Armitage, A.E., James, R.S., and Goff, S.P.
1996: Gold mineralization in Archean banded iron-formation, Third Portage Lake area, Northwest Territories, Canada, Exploration Mining Geology, v. 5, p. 1–15.

Ashton, K.E.
1981: Preliminary report on geological studies of the "Woodburn Lake Group" northwest of Tehek Lake, District of Keewatin. *In* Current Research, Part A, Geological Survey of Canada, Paper 81-1a, p. 269–273.
1982: Further geological studies of the "Woodburn Lake Group" northwest of Tehek Lake, District of Keewatin. *In* Current Research, Part A, Geological Survey of Canada, Paper 82-1A, p. 151–157.
1988: Precambrian geology of the southeastern Amer Lake area (66H/1), near Baker Lake, N.W.T.; unpublished Ph.D. Thesis, Queen's University, Kingston, 335p.

Davis, W.J. and Zaleski, E.
1998: Geochronological investigations of the Woodburn Lake group, Western Churchill Province: preliminary results; *In* Radiogenic Age and Isotopic Studies: Report 11, Geological Survey of Canada, Paper 98-2, p. 89–97.

Fraser, J.A.
1988: Geology of the Woodburn Lake map area, District of Keewatin. Geological Survey of Canada, Paper 87-11, 12 p.

Henderson, J.R. and Henderson, M.N.
1994: Geology of the Whitehills-Tehek Lakes area, District of Keewatin, Northwest Territories (parts of 56D, 66A, and 66H), Geological Survey of Canada, Open File 2923, scale 1:100 000.

Henderson, J.R., Henderson, M.N., Pryer, L., and Cresswell, D.
1991: Geology of the Whitehills-Tehek area, District of Keewatin: an Archean supracrustal belt with iron-formation-hosted gold mineralization in the central Churchill Province. *In* Current Research, Part C; Geological Survey of Canada. Paper 91-C, p. 149–156.

Heywood, W.W. and Schau, M.
1981: Geology of the Baker Lake region, District of Keewatin. *In* Current Research, Part A, Geological Survey of Canada, Paper 81-1A, p. 259–264.

Hughes, T.N.J. and Legault, M.M.
1991: Geological compilation, Meadowbank Joint Venture, District of Keewatin, N.W.T. Agnico-Eagle Group, Exploration Division, scale 1:20 000.

Kerswill, J.A., Goff, S.P., Wilkinson, L., Jenner, G.A., Kjarsgaard, B.A., Bretzlaff, R., and Samaris, C.
1998: An update on the metallogeny of the Woodburn Lake group, western Churchill Province, Northwest Territories; *in* Current Research 1998-C, Geological Survey of Canada, Paper 98-1C, p. 29–41.

Kjarsgaard, B.A., Kerswill, J.A., and Jenner, G.A.
1997: Lithostratigraphy and metallogenic implications of komatiite - banded iron-formation - felsic volcanic rocks of the Archean Woodburn Lake group, Pipedream lake, central Churchill Province, Northwest Territories; *in* Current Research 1997-C, Geological Survey of Canada, Paper 97-1C, p. 101–110.

LeCheminant, A.N. and Roddick, J.C.
1991: U-Pb zircon evidence for widespread 2.6 Ga felsic magmatism in the central District of Keewatin, N.W.T. *In* Radiogenic Age and Isotopic Studies: Report 4, Geological Survey of Canada, Paper 90-2, p. 91–99.

Miller, A.
1997: The Goose Island sulphide iron-formation-hosted fold deposit: a recrystallized syngenetic gold deposit, Woodburn Lake Group, western Churchill Province (draft); unpublished report to Cumberland Resources Ltd., June 10, 1997, 73 p.

Patterson, J.G.
1986: The Amer Belt; remnant of an Aphebian foreland fold and thrust belt. Canadian Journal of Earth Sciences, v. 23, p. 2012–2023.

Pehrsson, S.J., Wilkinson, L., Zaleski, E., Kerswill, J., and Alexander, B.
2000: Structural geometry of the Meadowbank deposit area, Woodburn Lake group implications for genesis of a major gold deposit in the Western Churchill Province. Geocanada 2000, The Millennium Geoscience Summit. CD-ROM Abstract #829 (Extended abstract).

Pehrsson, S.J., Jenner, G., and Kjarsgaard, B.K.
2002: Is the Ketyet River group correlative the Penrhyn and Piling groups of northeastern Rae?: implications for Paleoproterozoic orogenesis in the western Churchill Province. *In* Program with Abstracts, Geological Association of Canada-Mineralogical Association of Canada Annual Meeting 2002, Saskatoon, Cd-rom Abstract #1002.

Peterson, T.D., Van Breemen, O., Sandeman, H., and Cousens, B.
In press: Proterozoic (1.9–1.7 Ga) igneous suites of the western Churchill Province: granitoid and ultrapotassic magmatism in an Archean hinterland, Precambrian Research.

Rainbird, R.H., Hadlari, T., and Davis, W.
2001: Sequence stratigraphy and tectonic setting of the 1.78–1.74 Ga Wharton Group, Baker Lake and Thelon Basins, Western Churchill Province, Nunavut, Geological Association of Canada-Mineralogical Association of Canada Annual Meeting, Abstracts with Programs, v. 26, Volume 26, p. 123.

Roddick, J.A., Henderson, J.R., and Chapman, H.J.
1992: U-Pb ages from the Archean Whitehills-Tehek lakes supracrustal belt, Churchill Province, District of Keewatin, Northwest Territories, *in* Radiogenic Age and Isotopic Studies: Report 6, Geological Survey of Canada , Paper 92-2, p. 31–40.

Schau, M.
1983: Baker Lake (56D); Geological Survey of Canada, Open File 883, scale 1:100 000.

Sherlock, R.L., Alexander, R.B., March, R., Kellner, J., and Barclay, W.A.
2001a: Geological setting of the Meadowbank iron-formation-hosted gold deposits, Nunavut, Geological Survey of Canada, Current Research, no. 2001-C11, 2001; 23 pages.
2001b: Geology of the Meadowbank Iron-Formation Hosted Gold Deposit, Geological Survey of Canada, Open File 3149, scale 1:5 000.

Wilkinson, L., Pehrsson, S., Zaleski, E., Kerswill, J., and Alexander, B.
2001: Woodburn Lake group: Structural geometry of the Meadowbank deposit area implications for genesis of a major gold deposit in the Western Churchill Province. Cordilleran Round-up, Vancouver, B.C., January 22–26, 2001.

Zaleski, E., Corrigan, D., Kjarsgaard, B.A., Kerswill, J.A., Jenner, G.A., and Henderson, J.R.
1997: Geology, Woodburn Lake group, Meadowbank River to Tehek Lake(66H/1, 56E/4), District of Keewatin (Nunavut), Northwest Territories; Geological Survey of Canada, Open File 3461, scale 1:50 000.

Zaleski, E., Duke, N., L'Heureux, R., and Wilkinson, L.
1999a: Geology, Woodburn Lake group, Amarulik Lake to Tehek Lake, Kivalliq Region, Nunavut; Geological Survey of Canada, Open File 3743, scale 1:50 000.

Zaleski, E., L'Heureux, R., Duke, N., Wilkinson, L., and Davis, W.J.
1999b: Komatiites and felsic volcanic rocks overlain by quartzite, Woodburn Lake group, Meadowbank River area, Western Churchill Province, Northwest Territories (Nunavut); *in* Current Research 1999-C, Geological Survey of Canada, p. 9–18.

Zaleski, E., Pehrsson, S., Duke, N., Davis, W.J., L'Heureux, R., Greiner, E., and Kerswill, J.A.
2000: Quartzite sequences and their relationships, Woodburn Lake group, western Churchill Province, Nunavut. Geological Survey of Canada, Current Research 2000-C7, 10 p. (Online; http://www.nrcan.gc.ca/gsc/bookstore)

Zaleski, E., Davis, W.J., and Sandeman, H.A.
2001: Continental rifting, mantle magmas, and basement/cover relationships, *in* Cassidy, K., Dunphy, J.M. and Van Kranendonk, M.J., ed., International Archean Symposium, 24–28 September, Volume 2001/37: Perth, Western Australia, AGS0-Geoscience Australia, p. 374–376.




LOCATION MAP

OPEN FILE 4269
GEOLOGY
MEADOWBANK GOLD DEPOSIT AREA
NUNAVUT

Scale 1:20 000/Échelle 1/20 000
metres 500 0 500 1000 1500 2000 mètres

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2004

Projection transverse universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2004



NATIONAL TOPOGRAPHIC SYSTEM REFERENCE

LEGEND

PALEOPROTEROZOIC

PNgt — Nueltin granite suite: granitic dykes, pink to red stained hematitic, medium- to coarse-grained, leucocratic, commonly porphyritic with quartz and perthitic K-feldspar phenocrysts, biotite, magnetite, apatite, fluorite and titanite in intersitial clots, vuggy textured locally.

PHgt — Hudson granite suite: granitic dykes, pink, leucocratic, medium-grained sugary, equigranular to weakly porphyritic, minor biotite, local garnet, fluorite.

ARCHEAN OR PALEOPROTEROZOIC

(all rocks are metamorphosed at greenschist to amphibolite facies)

APgb — Gabbro, quartz leucogabbro, monzodiorite: medium to dark green, medium- to coarse-grained, ophitic, typically with hornblende oikocrysts, magnetite, minor biotite, titanite, massive to foliated.

APgm — Granite: white to pale pink, leucocratic, medium- to coarsed-grained, inequigranular to weakly K-feldspar-phyric, biotite-muscovite-bearing, local diffuse pegmatitic enclaves; as stocks and porphyritic dykes, massive to locally mylonitic.

Quartzite and associated rocks

APqz — Quartzite, minor oligomictic conglomerate, slate, quartz-muscovite schist: white, grey, blue or pink, locally green fuchsitic or reddish hematic, medium- to coarse-grained, crossbedded, massive to strongly cleaved.

APsl — Slate, phyllite: black to light grey, thinly bedded, very fine- to fine-grained commonly characterized by chlorite-muscovite-kyanite+/-chloritoid assemblages, foliated to schistose.

APqcg — Oligomictic conglomerate: white to dark grey to gossanous, orthoquartzite and vein-quartz pebble- to boulder-sized clasts in a fine- to medium-grained micaceous matrix, commonly pyritic, locally bedded, massive to cleaved.

APcg — Polymictic conglomerate: vari-coloured, rounded to subrounded pebbles, cobbles and boulders of orthoquartzite, vein quartz, quartz-feldspar porphyry, iron-formation, felsic to mafic volcanic rocks, and granite in a white quartz-rich to grey micaceous to pyritic dark or rusty matrix, locally bedded and transitional to oligomictic conglomerate, massive to cleaved to schistose.

NEOARCHEAN

(all rocks are metamorphosed at greenschist to amphibolite facies)

Agt — Syenogranite, monzogranite: pink to buff, medium- to coarse-grained, equigranular or with microline megacrysts; biotite, hornblende, epidote and/or magnetite, retrograde chlorite and muscovite common, massive to foliated, locally mylonitic.

Afp — Tern Lake porphyry: white to buff, fine-grained felsic porphyry, with quartz, K-feldspar and plagioclase phenocrysts; cm-dm scale layering defined by variations in abundance of phenocrysts; Afpg: Biotite granitodiorite and porphyry: white to buff, fine- to coarse-grained, with plagioclase and K-feldspar phenocrysts, biotite, hornblende, massive to foliated, locally mylonitic.

Amarulik wacke and associated rocks

Aakg — Arkosic greywacke, minor siltstone, mudstone and conglomerate: buff to medium grey, quartz and felspar crystal and lithic clasts in a medium- to coarse-grained bt-msc quartzofeldspathic matrix, thick massive beds common, poorly sorted, locally displaying grading; massive to foliated, recrystallized; Aacg: Pebble conglomerate; granule- to cobble-size quartzite, vein quartz, bt-schist and iron-formation clasts in a grey-buff quartzofeldspathic matrix, unsorted.

Aqz — Quartzite, minor oligomictic conglomerate, quartz-muscovite schist. White, grey, blue quartzite, locally with abundant muscovite, minor kyanite, fuchsite; medium- to coarse-grained, massive to foliated, locally schistose.

Aqcg — Oligomictic conglomerate; white to dark grey, orthoquartzite to vein quartz pebbles to boulders in a muscovite and quartz-rich matrix, locally bedded, cleaved to foliated.

Acg — Polymictic conglomerate, vari-coloured, orthoquartzite, iron-formation and polycrystalline quartz to vein quartz pebbles to boulders in a muscovite and quartz-rich matrix, locally bedded, cleaved to foliated.

Aaif — Iron-formation: dominantly oxide, local silicate facies, banded black-grey or white, typically rusty; chert, magnetite and/or silicate-rich bands or lamellae with grunerite, hornblende and garnet; typically with thin Fe-rich, garnetiferous pelitic schist interbeds, foliated.

Aasl — Slate, siltstone: medium grey to black or green; biotitic-pelite to biotite-hornblende schist; fine- to medium-grained, locally with porphyroblastic garnet-hornblende or garnet-staurolite and/or cordierite, andalusite, foliated to schistose.

WOODBURN GROUP and associated rocks (2.72–2.71 Ga)

Awv — Undifferentiated volcanic rocks and associated sediments; dominantly felsic to intermediate volcanic, volcaniclastic and volcanogenic sedimentary rocks; minor mafic or ultramafic rocks gabbro and iron-formation; rare arkose, phyllite, marble.

Awif — Iron-formation, dominantly oxide facies, local silicate or sulphide facies; banded white/grey and black, rusty; chert and magnetite bands, silicate-rich layers or lamellae with grunerite, chlorite, biotite; sulphide-rich layers with pyrite and pyrrhotite; fine- to medium-grained, typically with thin Fe-rich shale and wacke interbeds, foliated; Sulphidic iron-formation and massive to semi massive sulphide; banded white/grey and yellow metallic, rusty; pyrite and pyrrhotite-rich layers with magnetite, lesser arsenopyrite and silicate-rich layers.

Awg — Greywacke, slate, minor granule conglomerate and iron-formation; grey to brown, graded greywacke to slate couplets, chlorite and/or biotite bearing, fine- to medium-grained, well bedded, massive to cleaved or schistose.

Awsl — Slate to phyllite: black to light grey, very fine- to fine-grained commonly characterized by chlorite-muscovite-chloritoid ± kyanite assemblages, thin bedded to laminated, foliated to schistose.

Awsm — Siliceous marble: buff, tan and white, modified cm-scale bedding defined by fine-grained quartz-dolomite layers, very coarse dolomite sheaves up to 30 cm, and medium-grained granular dolomite ± calcite layers with tremolite, phlogopite, detrital quartz or rare humite; massive to foliated.

Awms — Quartz-muscovite ± chlorite schist: grey to green to pink, fine-grained, locally bedded, minor quartz eyes in a homogenous muscovite, quartz matrix with local carbonate and plagioclase; lesser quartz-plagioclase-muscovite-chlorite schists likely related to unit Awvc; Awmp: Quartz-plagioclase muscovite schist with 2–5 mm plagioclase porphyroblasts.

Awi — Intermediate volcanic rocks; crystal and lapilli tuffs and breccias, local subvolcanic intrusions and possible lava flows; white to grey or buff; euhedral blue quartz and/or plagioclase, locally amphibole phenocrysts; very fine-grained to aphanitic, commonly with abundant disseminated carbonate; foliated, locally bedded and transitional to reworked deposits; includes volcaniclastic rocks and felsic volcanics.

Awfq — Felsic volcanic rocks; massive volcanic flows and subvolcanic intrusions, minor volcaniclastic rocks; white to buff, with blue quartz and euhedral plagioclase phenocrysts, rare tourmaline, fine-grained to aphanitic, massive to foliated.

Awvc — Felsic to intermediate reworked volcaniclastic and related rocks; tuffaceous wacke, greywacke, minor siltstone and mudstone, minor primary volcaniclastic rocks; grey or pale pink or green banded, fine- to medium-grained, commonly micaceous and calcareous; quartz-feldspar-muscovite-chlorite- carbonate schists, cm- to m-scale layering, local graded beds; Awlam: laminated volcaniclastic wacke with amphibole porphyroblasts, grey to green mm- to cm-scale layering overgrown by foliated to randomly oriented coarse porphyroblasts of amphibole, locally amphibole obliterates layering, also fine-grained amphibolite.

Awm — Mafic volcanic and related rocks; mafic volcanic flows, amphibolite, subvolcanic gabbroic intrusions; black to green; fine- to medium-grained, medium- to coarse-grained, locally ophitic to subophitic textured, massive to foliated, rarely pillowed; Awa: Gabbro, peridotite, spotted amphibolite, diorite; black to dark green to grey; coarse-grained, amphibole and/or plagioclase phenocrysts or porphyroblasts, massive to foliated, may include some recrystallized, massive mafic volcanic flows; Amd: diorite.

Awwk — Ultramafic to mafic volcaniclastic rocks; dark green to black or brown; fine- to medium-grained, plagioclase, minor quartz and lithic clasts in a chlorite-rich matrix, possible tuffaceous volcaniclastic rocks with plagioclase phenocrysts, streaky layering, locally bedded.

Awkm — Ultramafic and related rocks: komatiite and basaltic komatiite flows, minor tholeiitic mafic flows; gabbro, peridotite, basaltic komatiite and mafic dykes, rare ultramafic to mafic volcaniclastic rocks; green, tan or black, massive to layered, commonly with spinifex and cumulate zones, rarely polyhedral jointing, breccia zones; carbonate alteration common, foliated to schistose; Awks: ultramafic schist characterized by talc, tremolite, serpentine, magnetite and/or Fe carbonate, basaltic komatiitic protolith.

Geological contact (approximate, inferred)
Bedding, (inclined, upright, overturned)

Structures of unspecified generation
Main foliation in outcrop (Sm)
Crenulation cleavage in outcrop
Crenulation lineation
Fold axis
Fault

Archean or Proterozoic structures
S_1 foliation
F_1 fold axial plane
L_1 mineral/stretching lineation

Proterozoic structures
S_2 foliation, crenulation to differentiated cleavage
F_2 fold axial plane
F_2 fold axis; z,u
L_2 intersection lineation
D_2 thrust fault
S_3 foliation, crenulation to differentiated cleavage
F_3 fold axial plane
L_3 crenulation lineation
F_3 fold axis; z, u
S_4 crenulation cleavage
F_4 fold axis; s, z
L_4 crenulation lineation
F_4 axial plane
Principal mineral occurrences (number refers to ID column of mineral occurrence table)
Outcrop

Geology by L. Wilkinson (1997–2000), E. Zaleski (1996–2000), J.A. Henderson (1990), D. Corrigan (1996, 1998), S. Pehrsson (1999), R. Leroux (1999), E. Grenier (1999), and W. Bleeker (2000)

Geological compilation by S. Pehrsson and L. Wilkinson, 2001–2002

Co-ordinated through the auspices of the Western Churchill NATMAP project

Digital cartography by J. Pratt and M. Proulx, Earth Sciences Sector Information Division (ESS Info)

This map was produced from processes that conform to the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001: 2000 standard

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital NTDB base data at 1:50 000 scale was not available for the map area. An approximately 50 000 scale digital base map was generated from Landsat Mapper (TM) imagery, using image analysis and GIS techniques, to identify vectorize and generalize lake outlines

Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area. Magnetic declination 2004, 2°2'W, increasing 5.9' annually



Figure 1. Simplified geology of the Rae and Hearne domains in the Western Churchill Province of the Canadian Shield (modified after Zaleski et al., (1997a)). Inset depicts the location of the map area in Northern Canada.



Figure 2. Schematic timescale illustrating the relative age of stratigraphic units, magmatism, deformation and metamorphic events in the Woodburn project area. Note that MesoArchean basement, the Whitehills conglomerate, monzodiorite-syenite and Hudson intrusive suites and Baker Lake Group are not represented in the map area itself but provide key relative time-deformation markers. Age of M2 metamorphism from unpublished data of Berman. U-Pb ages as follows: * unpublished data of Davis and Zaleski; ^Roddick et al. (1992); # Rainbird et al. (2001). Discussed in text.



Figure 3. A) Compositionally-layered, carbonate-bearing, felsic-intermediate reworked volcaniclastic unit (Awvc), one of the predominant map units of the ca. 2.71 Ga Woodburn Lake group. B) Photomicrograph of kyanite-bearing, chlorite-muscovite-chloritoid schist (Asl) associated with quartzite. Note that the kyanite porphyroblasts are wrapped by and overgrow the regional S_2 foliation. C) Close, chevron-style, southeast-vergent F_3 fold of S_2 foliation (photo looking southwest). D) Photomicrograph of F_3 crenulations overprinting S_2. Note that phyllosilicates and chlorite are bent without recovery, demonstrating that F_3 occurred after the regional thermal peak. E) Upright, open, northeast-trending F_4 folds developed on shallowly to moderately east-dipping S_2 foliation. F) Development of late, radiating grunerite on S_4 cleavage surfaces.

Facies zone	Assemblage (assemblages also contain qtz + pl with the exception of ultramafic rocks and iron-formation)	Rock Type
Lower Greenschist	bio + chl +/- musc bio +/- musc	Semi-pelite
	chl + musc + cld +/- ky	Phyllite
Middle Greenschist	chl + musc chl + bio +/- musc +/- ep	Felsic-Intermediate Volcaniclastic
	chl + act	Mafic
	chl + mgs + srp + tlc	Ultramafic
	chl + stp ^ chl + gru	Iron-Formation
	bio + musc + grt msc + chl + crd	Semi-pelite
	chl + musc + cld +/- ky	Phyllite
	bio + musc bio + grt +/- musc bio + grt + musc bio + musc + grt + st bio + musc + act	Felsic-Intermediate Volcaniclastic
Upper Greenschist	hbl + pl + ep +/- bt hbl + pl + ep + grt	Mafic
	chl + mgs + tlc + tr	Ultramafic
	cum-gru + bio + hbl	Iron-Formation
	cal + ep + tr + dol	Marble
	bio + grt +/- st bio + st + musc bio + st + grt +/- crd bio + st + and +/- grt bio + st + and +/- musc	Semi-pelite
	bio + musc + grt bio + musc + hbl +/- act bio + musc + hbl +/- grt bio + hbl +/- musc	Felsic-Intermediate Volcaniclastic
Amphibolite	hbl + pl	Mafic
	mgs + tr + srp + hbl	Ultramafic
	hbl + grt cumm + grt + hbl hbl + bio + grt	Iron-Formation
	cal + dol + tr cal + dol + hum + phl	Marble

Table 2. Metamorphic mineral assemblages of selected rock types.

ID	Commodities	Minerals	Type	Status	Name (Alternate name)	Comments
1	Au	Py-po-asp	gold	extensively drilled prospect	Third Portage zone	sulphide-bearing iron-formation
2	Au	Py-po-asp	gold	extensively drilled prospect	North Portage zone	sulphide-bearing iron-formation
3	Au	Py-po-asp	gold	extensively drilled prospect	Goose Island zone	sulphide-bearing iron-formation
4	Au	Py-po-asp	gold	extensively drilled prospect	Bay zone	sulphide-bearing iron-formation
5	Au-Pb-Zn		Polymetallic	drilled prospect		Tern Lake porphyry/granite
6	Au-Pb-Zn		Polymetallic	drilled prospect		Tern Lake porphyry/granite
7	Zn		base metal	drilled prospect		Mudstone

Table 1. Principal mineral occurrences of the Meadowbank deposit area.



Figure 4a. Sm, S_1 and F_2 structure traces for the map area. Light red lines are the outline of iron-formations for reference. F_2 axial traces are interpreted from a) the relationship between Sm/S_1/lithologic contacts and S_2 form surfaces (Figure 4b) and b) the asymmetry of minor F_2 folds.



Figure 4b. Distribution of S_2 foliations and interpreted F_4 axial surfaces. Red lines are iron-formations for reference. F_4 surfaces interpreted from the distribution of S_4, S_3 and S_2 structures (compare with Figure 4c) and asymmetry of F_4 folds.



Figure 4c. Distribution of S_3 and S_4 fabrics in the map area. Light red lines are outline of iron-formations for reference.



Figure 4d. Interpreted regional folds/faults with respective generation, based on lithology and fabric distribution, asymmetry of minor folds and plunge variations. See text for discussion.



Figure 5. Distribution of M2 mineral zones and M3 overprinting assemblages. Note that M3 post-tectonic porphyroblasts transect the M2 zones and are regional in scope, suggesting that they are not related to a deposit-centred metasomatic event as suggested by Armitage et al. (1994), but a late, low-grade regional metamorphism. Given ca.1.8 Ga $^{40}Ar/^{39}Ar$ ages for individual M3 biotite porphyroblasts (Villeneuve, pers.comm.) We suggest that M3 may be related to contact aureoles of local Paleoproterozoic granites.

OPEN FILE
DOSSIER PUBLIC
4269
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
2004

Open files are products that have not gone through the GSC formal publication process.

Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la CGC.

Recommended citation:
Pehrsson, S., Wilkinson, L., and Zaleski, E.
2004: Geology, Meadowbank gold deposit area, Nunavut; Geological Survey of Canada, Open File 4269, scale 1:20 000.

Canada