Natural Resources Canada | Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA

DESCRIPTIVE NOTES

INTRODUCTION

The Schultz Lake area (NTS 66 A) lies within the Archean Rae domain of the Western Churchill Province (Fig. 1). The area is underlain by a variety of Archean supracrustal rocks, paragneisses, granitoid gneisses and intrusive rocks that span the age range of at least 3.4 to 2.6 Ga (Table 1; Fig. 2; Bowring et al., 1989; Davis and Zaleski, 1998). This Open File report presents compilation at 1:250 000 based on sources as noted in Figure 2, and represents the geologic mapping contributions of Al Donaldson (1963–65), Tony LeCheminant (1983,1987–88), Al Miller (1980–1987), Fred Taylor (1980), Jack and Marietta Henderson (1994), Eva Zaleski (1998–2000), Ron Rainbird (1988, 1999, 2001), Thomas Hadlari (1999, 2001), Bruce Kjarsgaard (1997–1998), Tony Peterson (1998), and Sally Pehrsson (1999). Lawrence Aspler contributed significantly to a previous draft compilation. Previous relevant reports include Henderson and Henderson (1991), Zaleski et al. (1997, 1999a, b), Kerswill et al. (1999). Glacionic exposure is very good over the Hall Hills north of Whitehills Lake but is poor near Judge Sissons and Princess Mary Lakes, and particularly poor in areas underlain by the Barrensland Group. The hamlet of Baker Lake, Nunavut is situated in the southeast corner of the map sheet.

MESOARCHEAN ROCKS

The oldest rock unit in the Schultz Lake map sheet is a well-foliated, gneissic granodiorite (unit Agt), with rare included mafic banded mafic-felsic gneiss (Zaleski et al., 2001a), which underlies an area north of the Hall Hills. The locally schistose, carbonate and chlorite-altered granodiorite host has been dated at ca. 2970 Ma by Zaleski et al. (1997, 1999a, b) and at ca. 2960 Ma by Davis and Zaleski (1998). Unit Agt is cut by a set of well-differentiated west of the Thelon River, owing to the limit of reconnaissance mapping by Zaleski et al. (2001a). We note, however that sporadic outcrops of tonalitic gneiss were found in the southwest corner of the map sheet and to the west in the Aberdeen Lake map sheet, where LeCheminant et al. (1983) mapped other intermediate gneissic magmatic rocks.

NEOARCHEAN ROCKS

The Woodburn group

Outlier-and-belt preserved supracrustal rocks of the Neoarchean Woodburn group (Archean 1987) underlie much of the central and northern map area. Zaleski et al. (2000) subdivided the group in two general successions: 1) a lower succession consisting of interlayered ultramafic through felsic volcanic and volcaniclastic rocks with minor iron formation and muscovite-bearing quartzite (units Am, Amv, Aif, Awqt), all overlain by an extensive sequence of volcaniclastic to turbiditic wacke (unit Awkq); and 2) an upper succession of interbedded ortho- and heteplastic-quartzite, slate, phyllite, conglomerate, with minor mafic volcanic and reworked felsic volcanogenic rocks.

Ultramafic, mafic and felsic volcanic rocks of the lower succession underlie a belt of volcanic rocks north of Whitehills Lake that is dated at ca. 2734 ± 1 Ma (Zaleski et al., 2001b). This belt has a greater proportion of mafic pillow lavas than the typical lower succession to the north (Kjarsgaard et al., 1997). Mafic to felsic felsic volcanic and volcaniclastic rocks, also assigned to the lower succession, are exposed in a belt (NTS 66 E/16) in the southeast corner of the map area and to the southwest of the main Woodburn group belt in the Armer map sheet (NTS 66 H). Here, the volcanic rocks yield ages of ca. 2720 and 2710 Ma (Davis and Zaleski, 1998). Roddick et al. (1992) previously obtained an age of ca. 2.78 Ga for the Amarulik wackes, but Zaleski et al. (2001b) re-interpreted these rocks as largely reworked volcaniclastic sedimentary rocks of mostly ca. 2.68 Ga, with an older detrital zircon population.

Rocks of the upper succession of the Woodburn group may be present just south of Schultz Lake where thick quartzites with interspersed mafic and minor felsic volcanic rocks occur. The upper succession is interpreted to be ca. 2630 Ma (Zaleski et al., 2001b), based on ages of zircons from an interbedded reworked felsic volcanic rock in NTS 66 H. Henderson et al. (2000) reported that Re-Os systematics a maximum age for deposition of the upper succession, and the upper succession may actually represent the metamorphosed equivalents of the Paleoproterozoic Amer Group, which has strikingly similar facies associations.

Intrusive rocks

The region south of Whitehills Lake is underlain by the Akulliak gneiss (unit Ang), a foliated to gneissic granodiorite-granite-monzogranite pluton dated at ca. 2730 Ma (Davis and Zaleski, 1998). Although the pluton is not dated in the map sheet, it is intruded by inferred Archean granites and dikes that resemble it in terms of amphibolite and quartz-biotite (Zaleski et al., 2001a). The Akulliak gneiss typically has blue-grey quartz eyes, a low mafic mineral content and minor biotite typically altered to chlorite. An imprecise Archean age has been obtained from the granodiorite component (Schau., 1982) which could either represent a synvolcanic equivalent of ca. 2.7 Ga Woodburn group volcanogenic rocks or a component of the superficially similar Archean granodiorite north of Whitehills Lake. To the east, the amphibolite component defines two extensive mappable subunits; an older Archean layered amphibolite occurring as 100s of metre-thick horizons in the tonalite, and kilometre-scale metamorphosed sills or flows of uncertain Archean or Proterozoic origin.

The unfoliated Archean intrusive rock in the map area is a pink to buff, biotite ± hornblende-magnetite granite (unit Agt) that forms stocks, hosted, sheet-like bodies and northeast trend dikes. These rocks have been dated at ca. 2599–2920 Ga (Aitken, 1988; Roddick et al., 1992; Davis and Zaleski, 1998) and are variably deformed and recrystallized. In NTS 56 D, they comprise the Ishiak pluvonic complex (Schau et al., 1982). Similar, K-feldspar-phyric intrusions, between Princess Mary and Whitehills lakes, are tentatively assigned to this unit, which is correlated with the suite of 2.62–2.56 Ga granites noted throughout the Rae Province (Davis and Zaleski, 1998).

A foliated to gneissic granitoid unit (unit Agt) comprising variably deformed and metamorphosed biotite ± hornblende-bearing granodiorite, granite and tonalitic rocks, extends from the Hall Hay Hills to Princess Mary Lake. Granitic components of this unit are texturally phases that may be consistent with unit Agt. Southeast along the west shore of Baker Lake, unit Agt is commonly migmatitic and is cut by a variety of granites, by cataclasis and locally abundant metamorphosed and deformed amphibolite sheets. Mafic minerals such as biotite are typically altered to chlorite throughout this area. An unpublished age of 2894 Ma (LeCheminant and Roddick) has been obtained from a leucogranitic component north of Pitz Lake.

ARCHEAN OR PROTEROZOIC

A small body of diorite to granodiorite underlies the area north of Princess Mary Lake (unit APgd). Although roughly stock-like in character, similar to some of the Proterozoic granitoid rocks, its age is presently uncertain and it is designated Archean or Proterozoic.

PROTEROZOIC ROCKS

Ketyet River group

The Ketyet River group was informally introduced by Schau et al. (1982) to describe all metasedimentary rocks north of Baker Lake in NTS 56 D and, by extrapolation, NTS 66 A. The group was subdivided into a northern volcanic-dominated facies, and a southern sedimentary-dominated facies. The volcanic-dominated facies, which is the only facies associated represents rocks temporally and lithologically correlative with the Archean Woodburn group, whereas much of Schau's southern association represents a hitherto unrecognized Paleoproterozoic supracrustal sequence (Pehrsson et al., 2002). The informal names Ketyet River group and Woodburn group have been retained for the Proterozoic and Archean successions, respectively.

The Ketyet River group can be divided into three parts: 1) A lower succession of conglomerate (unit Pcpg), quartz-and feldspathic arenite (unit Pqz), iron-formation (unit Pif), and mafic volcanic rocks (unit Pmi); 2) a middle succession of mafic volcanic rocks (unit Pmi) and pelite (unit Pwk); and 3) an upper succession of mature quartzite (unit Pqz) and iron formation, minor marble and calc-silicate rocks. The lower succession has a maximum age of 2.45 Ga (Pehrsson et al., 2002), whereas the middle succession has a maximum age of 2.0 Ga (Davis and Zaleski, 1998) both from U-Pb detrital zircon analysis. Iron-formation occurs in both the upper and lower successions in NTS 56 D, but is noted only the upper succession in NTS 66 A. Sulphide-graphite-pyrite occur in the upper succession in drill core from the vicinity of the Kiggavik uranium deposit (A. Miller, pers. comm., 2002).

Dubawnt Supergroup

The southern edge of the map sheet overlaps the north margin of the Baker Lake sub-basin of the Dubawnt Supergroup (Fig. 1; Rainbird et al., 2003), a large intracontinental basin that straddles the Snowbird Tectonic Zone. The Dubawnt Supergroup is divided into three unconformity-bounded sequences, from oldest to youngest: 1) Baker Lake Group; 2) Wharton Group; and 3) Barrensland Group (Donaldson, 1965; Dall et al., 1992, Rainbird et al., 2003).

Baker Lake and Wharton groups

The Baker Lake Group is represented by a moderately south-dipping sequence of interbedded ultrapotassic volcanic rocks, conglomerate, and sandstone of the Christopher Island Formation (unit PtC). The sequence unconformably overlies Archean gneissic and plutonic rock of the Chesterfield Fault Zone north of Baker Lake and map unit Agt north of Princess Mary and Pitz lakes. The unconformity is marked by a several metre-thick carbonate-rich regolith (Rainbird and Hadlari, 2000). Present constraints from Ar-Ar, Pb-Pb and U-Pb age dating of flows, cogenetic intrusions, and diagenetic cements suggest that the Baker Lake Group was deposited between 1840 Ma–1790 Ma, and is thus correlative with the Dubawnt Supergroup (Rainbird et al., 2003).

The Baker Lake Group is in turn, unconformably overlain by the Wharton Group, comprising sub-arkosic sandstone of the Amarook formation (unit PA; Rainbird and Hadlari, 2000) and overlying rhyolite, conglomerate and sandstone of the Pitz formation (unit PP). The sub-Wharton Group unconformity cuts down through the Baker Lake Group in the Princess Mary Lake area where the Pitz and Amarook formations rest directly on Archean basement. U-Pb zircon ages of felsic flows indicate that the Wharton Group was deposited between 1780 and 1700 Ma (Rainbird et al., 2001).

Barrensland Group

The Barrensland Group is represented in the map area by the flat-lying to gently-dipping Thelon Formation (unit PT), a fluvial-eolian sandstone and conglomerate deposited on a significant regolith marked by pervasive quartz-hematite-illite-kaolinite, or at lower levels, chlorite-muscovite-dolomite alteration (unit PT; Chiarenzelli, 1983; Dall, 1994). This regolith is best developed on basement rocks where the Baker Lake and Wharton groups are absent. The distribution of the Thelon Formation in the Schultz and Baker lakes vicinities, presence of small outliers northwest of Pitz Lake, subcrop on the top of the Hall Hay Hills, and extensive development of regolith between indicates that the Barrensland Group was originally more extensive in the map area (Donaldson, 1965). This interpretation is estimated to have commenced by 1720 Ma, based on a U-Pb age of apatite cement in the sandstone (Miller et al., 1988).

Intense hematitic alteration produces a consistent positive total field aeromagnetic anomaly with a broad, gently sloping high that overprints or obscures the magnetic signature of underlying lithologies. The extensive regolith developed between Schultz and Judge Sissons lakes and formation outliers indicate that the sub-Thelon unconformity surface lay very close to the present day topographic surface. Regolith alteration is associated with uranium mineralization (U), including the Kiggavik (Lone Gull) deposit (Miller, 1983; Fuchs and Hilger, 1989).

Intrusive rocks

The oldest known Proterozoic intrusive rocks in the map area are deformed, bouldined metadiabase dykes and lenses (unit Pdb) noted west of Thelon River that have yielded a U-Pb baddeleyite age of 2153 ± 4 Ma (LeCheminant and Roddick, unpublished data). These dykes locally occur in unit Agt near Whitehills and Baker lakes, and are similar in age to the extensive Macquoid dykes of NTS 56 M to the south.

Several small bodies of massive to weakly foliated, leucocratic, pink monzogranite dated at ca. 1835 Ma (Roddick et al., 1992) occur north of Whitehills Lake (unit Pgt). Minor dykes, sheets and sills of the suite occur throughout the map area and as intrusions into metamorphosed Archean rock units. They form part of the 1.84–1.80 Ga Hudson granitoid suite (Hoffman and Bowring, 1984; Hoffman and Chamberlain, 1988; and van Breemen, 1998), and are known from the Aberdeen Lake area immediately west (LeCheminant et al., 1983).

The ca. 1810 Ma Martell syenites (unit Pse; Schau et al., 1982; Rainbird et al., 2002) underlie several small islands in Whitehills Lake. These brown to red, hypalsyenal syenites have distinctive hornblende contact aureoles with surrounding Ketyet River group wackes and are interpreted to be the cogenetic with alkalic Christopher Island Formation volcanics. Related syenite and phlogopite-lamprophyre dykes are numerous in the Whitehills area, but have not been differentiated at the scale of this compilation.

Two sets of unmetamorphosed diabase dykes transect the area. The older, northeast-trending set has been dated at ca. 1590 Ma (Rainbird et al., 2001) and is correlative with the Mackenzie dyke swarm (LeCheminant et al., 1987) and part of the 1750 Ma Mouffon dyke swarm in NTS 66 B, which is interpreted to feed amygdaloidal basalts of the Pitz Formation (LeCheminant et al., 1983), however it is not differentiated from unit Pdb, owing to insufficient detail at this scale. The second, northwest-trending set includes dykes up to 50 m wide that crosscut all units of the map area (unit Pedb). These are considered to be part of the 1267 Ma Mackenzie dyke swarm (LeCheminant and Heaman, 1989).

STRUCTURE AND METAMORPHISM

All rocks of the Schultz Lake map area are deformed and metamorphosed with the exception of the Dubawnt Supergroup, Martell syenites and younger diabase dyke swarms. The regional foliation compiled on this Open File is typically an S_1 or S_0 fabric, as established by more detailed mapping in the Whitehills Lake area (Zaleski et al., 2001a). These authors distinguished four main phases of regional deformation (D_1–D_4), within the middle Ketyet River group, thus constraining the regional tectonothermal events as younger than 2.0 Ga. These structures include west-northwest to northeast-trending, tight to recumbent to isoclinal F_1 folds, and north-west vergent mapscale F_2 folds and related faults (Zaleski et al., 2001a); minor, open to tight folds outcrop-scale, southeast-vergent F_3 folds, and open to tight, northerly-trending, upright, F_4 folds, such as the prominent fold in the Ketyet River group at Whitehills Lake. Zaleski et al. (2001a) also noted a locally penetrative, northeast-trending fabric (S_3) that is locally folded by F_4 (Zaleski, unpublished data), suggesting significant heterogeneity of later Proterozoic deformation.

Regional metamorphism varies from greenschist facies over the Whitehills greenstone belt and Ketyet River group to middle or upper amphibolite facies over Archean units across the map area. Low-pressure amphibolite-granulite facies assemblages are known from wackes of the Ketyet River group north and south Whitehills Lake. High-pressure garnet-amphibolite-clinopyroxene assemblages locally occur in unit Agnlt, near the community of Baker Lake.

The contact between a large package of rocks assigned to the Ketyet River group and underlying Archean rocks south of Schultz Lake, is interpreted to be a south-vergent thrust (A. Miller, pers. comm.). The character of the contacts of adjacent smaller occurrences of Ketyet River group are unknown and are assumed to be stratigraphic, as is the contact between the Ketyet River group and Archean rocks in the Whitehills Lake greenstone belt.

Several northeast-vergent, post-metamorphic, low-angle thrust faults juxtapose rocks of greenschist and amphibolite facies at Whitehills Lake and/or place older Archean units atop the Ketyet River group (Zaleski et al., 2001a). They are defined by high strain zones,metres to 10s of metres wide and locally by carbonate-chlorite alteration zones. These cannot be traced with certainty west of the Thelon River.

Schau et al. (1982) interpreted a multistage Neoarchean and ca. 1.9 Ga deformation history for the Chesterfield Fault Zone, which may extend into the east part of the map sheet. Preliminary thermobarometric data (Berman and Peterson, unpub. data) suggests that this crystic zone is the locus of a significant Proterozoic paleopressure discontinuity. The relationship between reworking of the Chesterfield Fault Zone and the various northwest-vergent faults is presently unknown.

It is to be expected that Archean deformation and/or metamorphism occurred in the Woodburn group and older rocks, however, the penetrative nature of Paleoproterozoic deformation and metamorphism has made it impossible to differentiate these, except locally at the outcrop or thin-section scale. The variable state of deformation within the both Archean and Proterozoic granite suites largely reflects heterogenous Proterozoic strain partitioning (Ashton (1988), Davis and Zaleski (1998), Zaleski et al. (2002)).

The Baker Lake and Wharton groups are both crosscut by east-striking normal faults, and northeast striking strike-slip faults (Rainbird and Hadlari, 2000; Hadlari and Rainbird, 2001). Principle expansion and tilting postdate deposition of the sub-horizontal overlying ca. 1720 Ma Thelon Formation (unit PT), and occurred during and after deposition of the 1.84–1.80 Ga Hudson suite (Fig. 2). Eta north-northwest-trending faults were subsequently re-activated and crosscut the sub-Thelon regolith (e.g. Kiggavik) and Thelon Formation with relatively minor offset.

ECONOMIC GEOLOGY

Rocks of the Schultz Lake area are host to the Kiggavik uranium deposit and related occurrences south of Schultz Lake (Fuchs and Hilger, 1989; Miller et al., 1988). Uranium occurs at the base contact of the Thelon Formation and in Archean rocks. Uranium is associated with hematite-illite-kaolinite alteration. Other types of occurrences (Cu-Pb-Zn-Ag, Mo and U) are known from the Ketyet River Group in the Whitehills Lake area (Pehrsson et al., 2002; Zaleski et al., 2001a). Minor uranium occurrences are also present in NTS 56 D (Schau et al., 1982). Taylor (1980) reports gahnite occurrences in a Mesoarchean(?) metagabbro sill, and a number of gahnite-bearing veins were also found southwest of Whitehills Lake in unit Ag.

Base metal and polymetallic occurrences are known from the Amarulik wackes and Whitehills greenstone belt (Zaleski et al., 2001a). Gold is present in iron-formation of the Woodburn Lake group immediately north of the map area at Third Portage Lake (Meadowbank gold deposit), and within volcanic rocks of the Whitehills greenstone belt and Ni-Cr occurrences are noted in the komatiitic rocks of the Woodburn group (Zaleski et al., 2001a).

ACKNOWLEDGMENTS

The production of this map evolved over the past 20 years, involving the contributions of many individuals as cited in the introduction. We particularly thank A.H. Miller, J.A. Donaldson, and A.N. LeCheminant, for discussions on the geology of the area. This compilation benefited from previous interpretations of the geology suggested by L.B. Aspler. Dr. Steven Barrie is thanked for her thorough and timely review. We also thank Marcelle Hauseux for some helpful additions.

REFERENCES

Ashton, K.E.

1981: Preliminary report on geological studies of the "Woodburn Lake Group" northwest of Tehek Lake, District of Keewatin; in Current Research, Part A, Geological Survey of Canada, Paper 81-1A, p. 269–274.

1988: Precambrian geology of the southeastern Amer Lake area (66 H/1), near Baker Lake, N.W.T.; unpublished Ph.D. Thesis, Queen's University, Kingston, 335 p.

Chiarenzelli, J.R.

1983: Mid-Proterozoic chemical weathering, regolith, and silcrete in the Thelon Basin, Northwest Territories. Unpublished Msc. Thesis, Carleton University, Ottwa, Ont., p. 336.

Davis, W.J. and Zaleski, E.

1998: Geochronological investigations of the Woodburn Lake Group, western Churchill Province, Northwest Territories: preliminary results; in Radiogenic Age and Isotopic Studies: Report 11; Geological Survey of Canada, Current Research 1998-F, p. 89–97.

Donaldson, J.A.

1965: The Dubawnt Group, District of Keewatin and Mackenzie. Geological Survey of Canada, Paper 64-20, 11 p.

1966: Geology, Schultz Lake, District of Keewatin; Geological Survey of Canada, Preliminary Map 7-1966, scale 1:250 000.

Fuchs, H.D. and Hilger, W.

1989: Kiggavik (Lone Gull): an uncontrmity-related uranium deposit in the Thelon Basin, Northwest Territories, Canada; in Möllerfahrte, E. (Ed.), Uranium Resources and Geology of North America. International Atomic Energy Agency, Vienna, pp. 429–454. IAEA-TECDOC-500.

Gall, Q.

1994: The Proterozoic Thelon paleosol, Northwest Territories, Canada. Precambrian Research, 68, p. 115–137.

Gall, Q., Peterson, T.D., and Donaldson, J.A.

1992: A proposed revision of early Proterozoic stratigraphy of the Thelon and Baker Lake basins, Northwest Territories; in Current Research, Part C; Geological Survey of Canada, Paper 92-1C, p. 129–137.

Hadlari, T. and Rainbird, R.H.

2001: Volcano-sedimentary correlation and fault relationships in the Baker Lake sub-basin, Thirty Mile Lake area, Nunavut; in Current Research, Geological Survey of Canada, 2001-C10, p. 9.

Henderson, J.R. and Henderson, M.N.

1991: Geology of the Whitehills-Schultz area, District of Keewatin: an Archean supracrustal belt with iron-formation and felsic metavolcanics in the central Churchill Province; in Current Research, Part C; Geological Survey of Canada, Paper 91-1C, p. 149–156.

Kerswill, J.A., Kjarsgaard, B.A., Bretzlaff, R., Jenner, G.A., and Semenes, C.

1999: Metallogeny and geology of the Hall Hay Hills area, central Churchill Province, Northwest Territories (Nunavut); in Current Research 1999-C; Geological Survey of Canada, p. 459–41.

LeCheminant, A.N. and Heaman, L.M.

1989: Mackenzie igneous events, Canada: middle Proterozoic hotspot magmatism associated with ocean opening. Earth and Planetary Science Letters, 96: 38–48.

LeCheminant, A.N. and Roddick, J.C.

1991: U-Pb zircon evidence for widespread 2.6 Ga felsic magmatism in the central District of Keewatin, N.W.T.; in Radiogenic Age and Isotopic Studies: Report 4, Geological Survey of Canada, Paper 90-2, p. 91–99.

LeCheminant, A.N., Ashton, K.E., Chiarenzelli, J., Donaldson, J.A., Rust, M.A., Tella, S., and Thompson, D.S.

1983, Geology of Aberdeen Lake map area, District of Keewatin: preliminary report; in Current Research, Part A., Geological Survey of Canada, Paper 83-1A, 1983; pages 437–448.

Miller, A.R.

1983: A Progress Report: Uranium-Phosphorous Association in the Helikian Thelon Formation and Sub-Thelon Saprolite, Central District of Keewatin; in Current Research, Part A, Geological Survey of Canada, Paper 83-01A, 1983, p. 459–464.

Miller, A.R., Cumming, G.L., and Krstic, D.

1989: U-Pb, Pb-Pb and K-Ar isotope study of the uraniferous phosphate-bearing rocks in the Thelon Formation, Dubawnt Group, Northwest Territories, Canada; Canadian Journal of Earth Sciences, v. 26, p. 867–880.

Pehrsson, S.J., Jenner, G., Kjarsgaard, B.A.

2002: The Ketyet River group: correlation with Paleoproterozoic supracrustal sequences of northeastern Rae and implications for Proterozoic orogenesis in the western Churchill Province. GAC-MAC annual meeting, Program with Abstracts, 99.

Peterson, T.D. and Van Breeman, O.

1999: Review and progress report of Proterozoic granitoid rocks of the western Churchill Province, Northwest Territories (Nunavut); in Current Research 1999-C; Geological Survey of Canada, p. 110–127.

Rainbird, R.H., Davis, W.J., Stern, R.A., Hadlari, T., and Donaldson, J.A.

2002: Integrated geochronology of the late Paleoproterozoic Baker Lake Group (Dubawnt Supergroup), Baker Lake Basin. GAC-MAC annual meeting, Program with Abstracts, p. 99.

Rainbird, R.H. and Hadlari, T.

2000: Revised stratigraphy and sedimentology of the Paleoproterozoic Dubawnt Supergroup at the northern margin of Baker Lake Basin, Nunavut. Geological Survey of Canada, Current Research 2000-C6, 5 p.

Rainbird, R.H., Hadlari, T., Aspler, L.B., Donaldson, J.A., LeCheminant, A.N., and Peterson, T.D.

2003: Sequence stratigraphy and evolution of the Paleoproterozoic intracontinental Baker Lake and Thelon basins, western Churchill Province, Nunavut, Canada. Precambrian Research, 125, p. 21–53.

Rainbird, R.H., Hadlari, T., and Davis, W.

2001: Sequence stratigraphy and tectonic setting of the 1.78–1.74 Ga Wharton Group, Baker Lake and Thelon Basins, Western Churchill Province, Nunavut; in Geological Association of Canada-Mineralogical Association of Canada Annual Meeting, Abstracts with Programs, v. 25, p. 123

Roddick, J.C., Henderson, J.R., and Chapman, H.J.

1992: U-Pb ages from the Archean Whitehills–Ishiak lakes supracrustal belt, Churchill Province, District of Keewatin, Northwest Territories; in Radiogenic Age and Isotopic Studies: Report 6; Geological Survey of Canada, Paper 92-2, p. 31–40.

Sandeman, H.A., Cousens, B.L., and Hemmingway, C.J.

2003: Continental tholeiitic mafic rocks of the Paleoproterozoic Hurwitz Group, Central Hearne sub-domain, Nunavut: insight into the evolution of the Hearne sub-continental lithosphere. Canadian Journal of Earth Sciences, 40, p. 1219–1237.

Schau, M.

1980: Zircon ages from a granite/ultramafite complex and a layered gneiss complex northeast of Baker Lake, District of Keewatin; in Loveridge, W.D., Rubidium-Strontium and Uranium-Lead Isotopic Age Studies, Report 3; Current Research, Part C; Geological Survey of Canada, Paper 80-1C, p. 237–238.

Schau, M., Tremblay, F., and Christopher, A.

1982: Geology of Baker Lake map area, District of Keewatin: a progress report; in Current Research, Part A, Geological Survey of Canada, Paper 82-1A, p. 143–150.

Taylor, F.C.

1980: Precambrian geology of the Hall Hay Hills area, District of Keewatin. Geological Survey of Canada, Memoir 415, 19 p.

Zaleski, E., Corrigan, D., Kjarsgaard, B.A., Jenner, G.A., Kerswill, J.A., and Henderson, J.R.

1997: Preliminary results of mapping and structural interpretation from the Woodburn project, western Churchill Province, Northwest Territories; in Current Research 1997-C, Geological Survey of Canada, p. 91–100.

Zaleski, E. and Davis, W.J.

2002: Mantle-derived magmatism and continental rift deposits of the Neoarchean Woodburn Lake group, Western Churchill Province. GAC-MAC annual meeting, Program with Abstracts, p. 138.

Zaleski, E., Davis, W.J., and Sandeman, H.A.

2001b: Continental extension, mantle magmas and basement/cover relationships, Cassidy, K.M., Dunphy, J.M., and Van Kranendonk, M.J., (eds.), Fourth International Archean Symposium, v. 2001:57. Perth, Australia, AGSO-Geoscience Australia, p. 374–376.

Zaleski, E., Duke, N., L'Heureux, R.L., Davis, W.J., and Wilkinson, L.

1999a: Clastic and chemical sedimentary sequences, Woodburn Lake group, Amarulik Lake to Tehek Lake, western Churchill Province, Northwest Territories (Nunavut); in Current Research 1999-C; Geological Survey of Canada, p. 19–28.

Zaleski, E., L'Heureux, R.L., Duke, N., Wilkinson, L., and Davis, W.J.

1999b: Komatiitic and felsic volcanic rocks overlain by quartzite, Woodburn Lake group, Meadowbank River area, western Churchill Province, Northwest Territories (Nunavut); in Current Research 1999-C; Geological Survey of Canada, p. 9–18.

Zaleski, E., Pehrsson, S.J., Davis, W.J., Greiner, E., L'Heureux, R., Duke, N., and Kerswill, J.A.

2001a: Geology, Hall Hay Hills to Whitehills Lake area, Nunavut; Geological Survey of Canada, Open File 3697, scale 1:50 000.

Zaleski, E., Pehrsson, S.J., Duke, N., Davis, W.J., L'Heureux, R., Greiner, E., and Kerswill, J.A.

2000: Quartzite sequences and their relationships, Woodburn Lake group, western Churchill Province, Nunavut; in Current Research 2000; Geological Survey of Canada, 10 p.



Figure 1. Regional geology highlighting the location of Schultz Lake (NTS 66 A) in relation to Thelon and Baker Lake basins and to older Proterozoic supracrustal rocks.



1 Taylor, 1986, Map 1602A
2 Zaleski et al., 2001a, OF3697
3 Henderson and Henderson, 1994, OF2923
4 Zaleski et al., 1999, OF3743
5 Donaldson, 1966, Map 7-1966

Figure 2. Map references.



LOCATION MAP

OPEN FILE 1839

GEOLOGY

SCHULTZ LAKE

NUNAVUT

Scale 1:250 000/Échelle 1/250 000

kilometres 5 0 5 10 15 20 kilometres

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2004

Projection transverse universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2004



NATIONAL TOPOGRAPHIC SYSTEM REFERENCE AND INDEX TO ADJOINING GEOLOGICAL SURVEY OF CANADA MAPS

LEGEND

MESOPROTEROZOIC

Pmdb — *Mackenzie diabase dykes.*

PALEOPROTEROZOIC

DUBAWNT SUPERGROUP (units PC–P1)

BARRENSLAND GROUP (units P1–P1')

PT — *THELON FORMATION; quartz arenite; cross-stratified fluvial sandstone; very large-scale cross-stratified eolian sandstone.*

PTr — *SUB-THELON FORMATION; regolith; silcrete; pervasive quartz-hematite-illite-kaolinite or chlorite-muscovite-dolomite alteration (e.g. Chiarenzelli, 1983).*

unconformity

WHARTON GROUP (units PA–PP)

PP — *PITZ FORMATION; rhyolite and interbedded conglomerate and sandstone.*

PA — *AMAROOK FORMATION; very large-scale cross-stratified, subarkosic arenite (eolian facies); conglomerate and pebbly sandstone (fluvial facies).*

unconformity

BAKER LAKE GROUP (units Pgt–Pwk)

Pse — *Martell Syenite; brown to red, fine- to coarse-grained equigranular or porphyritic syenite.*

PCi — *CHRISTOPHER ISLAND FORMATION; massive minette flows; interbedded lithic arenite and cobble- to boulder-grade, clast-supported conglomerate (alluvial fan and fluvial facies).*

unconformity

Pgt — *Hudson Granite; pink leucocratic, medium grained equigranular to weakly porphyritic, minor biotite; 1835 ± 1 Ma (Roddick et al., 1992).*

intrusive contact

KETYET RIVER GROUP (informal) (units Pmi–Pwk)

Pwk — *Metawacke; buff, grey to pale-brown weathering, typically flaggy to schistose, pervasive carbonate, rare granular to pebbly interbeds.*

Pif — *Iron-formation; sulphide-silicate facies; hematite, pyrrhotite, grunerite, biotite.*

Pcg — *Polymictic boulder, cobble and pebble conglomerate with wacke and slate interbeds; matrix- and clast-supported; < 2.0 Ga (Pehrsson et al., 2002).*

Pdb — *Undifferentiated diabase or metagabbro dykes.*

Pqz — *Quartzite, locally crossbedded and rippled; white, but locally hematitic; minor metamorphic kyanite; fuchsite; minor oligomictic conglomerate; < 2.45 Ga (Pehrsson et al., 2002).*

Pqcg — *Oligomictic conglomerate; locally with chert and quartzite clasts.*

Pmi — *Mafic to intermediate volcanic rocks; dark to medium green, fine- to medium-grained, rarely porphyritic to subophitic; commonly plagioclase porphyritic and locally megascystic (1–3 cm), variably foliated, minor volcaniclastic rocks.*

ARCHEAN OR PROTEROZOIC

APgd — *Granodiorite, diorite; age uncertain.*

NEOARCHEAN

Agt — *Granite; medium- to coarse-grained; biotite, amphibole, and/or magnetite-bearing; massive to variably foliated.*

Ag, Agnlt — *Un-subdivided granitoid rocks: granite, granodiorite, tonalite, minor diorite and gabbro; biotite ± hornblende; medium- to coarse grained; foliated to gneissic; 2695 ± 4 Ma (LeCheminant and Roddick, unpublished data); locally mylonitic, e.g. extension of Chesterfield Fault Zone (Agnlt) of Schau et al. (1982).*

Atng — *Porphyritic to inequigranular granodiorite, tonalite, and amphibolite; gneissic to layered (extension of Akulliak Gneiss from Schau et al., 1982).*

WOODBURN GROUP (informal) (units Am–Awkq)

Awkq — *Arkosic metawacke; buff to medium grey; poorly sorted, massive to graded beds; with interbedded iron-formation, volcaniclastic and minor mafic to felsic volcanic rocks, psammite to rare pelite.*

Afvc — *Felsic to intermediate volcaniclastic rocks and volcanic wacke; fine-grained to gritty to pebbly with abundant quartz and plagioclase crystals; muscovite, carbonate and chlorite.*

Aqz — *Quartzite; massive to schistose; local muscovite, biotite, sillimanite; local kyanite-chloritoid schist.*

Aif — *Iron-formation; oxide, carbonate, and silicate facies; predominately white, also grey, blue, or pink.*

Am — *Mafic volcanic rocks; minor intermediate to felsic volcanic and volcaniclastic rocks, also unidifferentiated metasedimentary rocks; locally schistose; minor komatiite flows (K).*

MESOARCHEAN

Agd — *Granodiorite; foliated to schistose, typically with strong fabrics; chlorite-biotite bearing; medium- to coarse-grained; white weathering.*

Geological contact (defined, inferred)

Limit of mapping

Thrust (defined, inferred)

Fault (defined, inferred)

Barrensland Group/Wharton Group unconformity (defined, inferred)

Wharton Group/Baker Lake Group unconformity (defined, inferred)

Baker Lake Group/Archean unconformity (defined, inferred)

Bedding, inclined; top unknown, top known, overturned

Crenulation cleavage

Foliation, inclined

Lineation, with plunge

Crenulation lineation

Mesoscopic fold axis

Antiform

Axial trace (F_2); antiform, synform

Axial trace (F_3); antiform, synform

Axial trace (F_4)

Geochronology location (see Table 1)

Local uranium mineralization U×

Table 1. Geochronology.

MAP #	AGE/ERROR	METHOD/MINERAL	ROCK TYPE	REFERENCE
8	1835 ± 1 Ma	U-Pb monazite	massive granite	Roddick et al., 1992
7	< 2.0 Ga	U-Pb detrital zircon	feldspathic arenite	Pehrsson et al.,