



### LEGEND

Note: Map units listed below occur within one metre of the surface. Where organic or eolian sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that could contain placer gold. In order to accentuate these potentially exploitable deposits, a compound map unit is presented, e.g. CbA<sup>1</sup>. This means that colluvial complex sediments overlie alluvial terrace sediments thought to be late Tertiary in age. The legend is part of a larger regional study hence coloured boxes indicate units that appear on the map. In addition, not all symbols in the legend are represented on this map.

**CENOZOIC**

**QUATERNARY**

**HOLOCENE**

**m** Made Land: placer mines, roads, and airstrip

**ORGANIC DEPOSITS:** peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, fens, and swamps situated on valley bottoms; peatmoat commonly encountered within 1 m of the surface. Thermoclast collapse is common.

**Organic Blanket:** undrained; thickness > 1 m to 5 m

**Organic Veneer:** blanket bog generally < 1 m thick

**ALLUVIAL DEPOSITS:** gravel to silt size sediments, well stratified, deposited by streams

**Ap** Floodplain Sediments: gravel, cobble to pebbles; massive to well stratified, capped by sand and silt; flat lying, includes lacustrine and organic deposits in abandoned channels and backswamp areas; subject to periodic inundation and reworking by floods; thickness 1 to 5 m

**Af** Alluvial Fan Sediments: gravel, sand, silt, and diatomite, massive to well stratified; sediments form fan-shaped landforms or complexes of coalesced fan-shaped landform at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

**Ax** Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale

**HOLOCENE AND PLEISTOCENE (UNDIVIDED)**

**COLLUVIAL DEPOSITS:** stony diatomite resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits may contain reworked glacial/alluvial and moraine sediments within the limits of pre-Red River ice cover and reworked eolian sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating

**Cb-v** Colluvial Blanket and Veneer Sediments: diatomite, stony with a sandy matrix; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous over bedrock

**Ca** Colluvial Apron Sediments: locally diatomite and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope low complex of small steep debris flow and solifluction deposits; thickness is < 1 m at the upper and lower slope limit to up to 5 m or more in the thickest part of the apron

**Cl** Landslide Sediments: silt loam to boulders, poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly

**Cx** Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fan, colluvial blanket, landslide sediments and colluvial drift within the limits of glaciation; the unit commonly occurs along the lower slopes of valley margins

**CEA<sup>1</sup>** Colluvial/Eolian Apron (muck): primary deposits of eolian fine sand and silt reworked and interbedded with organic silt, and detrital, alluvial fan gravel and sand and variable amounts of stony colluvial diatomite; forms aprons along valley bottoms through reworking of eolian sediments from valley sides to valley floor; commonly preserved on north-facing slopes; thickness 1 to 20 m; commonly contains segregated bodies of ice and buried ice wedges

**MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)**

**AI<sup>1</sup>** Alluvial Terrace Sediments: gravel, cobble to pebbles with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m

**AI<sup>2</sup>** Alluvial Fan Sediments: single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments delineated by cryoturbation; thickness up to 10 m

**ACx<sup>1</sup>** Alluvial/Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; thin to thick bedded, interstratified with colluvial diatomite; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid-valley locations

**EOLIAN DEPOSITS:** well sorted medium sand to silt initially transported and deposited by wind action during glaciations and commonly reworked through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout low lying areas

**EB<sup>1</sup>** Eolian Blanket: fine sand and silt, well sorted, massive, may form crescent shape and linear dunes and features or gently undulating inter-dune eolian plains; thickness 1 to 5 m

**EB<sup>2</sup>** Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m

**LATE PLEISTOCENE - MCCONNELL GLACIATION**

**GLACIOFLUVIAL DEPOSITS:** gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

**GM** Glaciofluvial Terrace Sediments: gravel and sand, unweathered, forming one or more terraces

**MIDDLE PLEISTOCENE - REID GLACIATION**

**GLACIOFLUVIAL DEPOSITS:** gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

**GR** Glaciofluvial Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces

**LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - PRE-REID GLACIATIONS (UNDIVIDED)**

**GLACIOLASTRINE DEPOSITS:** well stratified sand, silt, clay, deposited in lakes ponded by glacial ice

**LUPR** Glacioastrine Undivided: sand, silt, and clay, undifferentiated at this scale of mapping

**GLACIOFLUVIAL DEPOSITS:** gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

**GI<sup>1</sup>** Glaciofluvial Terrace Sediments: gravel and sand, deeply weathered; incised into flights of terraces; thickness 1 to > 5 m

**MORAINAL DEPOSITS (TILL):** glacial diatomite, mainly till, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

**TYP<sup>1</sup>** Till Blanket: diatomite, stony, silty sand matrix; massive; conforms to underlying topography; thickness > 1 m; extensively colluviated on slopes

**TYP<sup>2</sup>** Till Veneer: diatomite, stony, silty sand matrix; massive; discontinuous and may contain extensive areas of thin (< 1 m) colluvium

**AI<sup>1</sup>** Alluvial Terrace Sediments: sandy pebbles and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Red River glacial or glacial drainage; thickness 1 to 5 m

**UNDIFFERENTIATED DRIFT:** diatomite, gravel, sand, silt and clay deposited from glacial ice, glacial streams, and glacially diamict lakes; extensive weathering, poor exposure and peatmoat make differentiation into component glacial sediments difficult; thicknesses commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and integrated with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

**D<sup>1</sup>** Drift: flat to gently sloping

**D<sup>1</sup>** Drift Modified by Landsliding: drift translated along failure plains into irregular steps and sub parallel escarpments

**D<sup>2</sup>** Fluvially Incised Drift: formerly extensive areas of drift incised by closely spaced stream valleys

**LATE PLEISTOCENE**

**VT** Basalt: columnar alkalic olivine basalt and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Redoubt Creek; thickness 10 m

**FLUCCINE AND LATE MIOCENE**

**ALLUVIAL DEPOSITS:** pingraic gravel and sand, highly dissected and deeply weathered

**PT** Pediment and Bajada Sediments: inclined fluvial surfaces which are found at a mesoscale position in unglaciated drainage systems; usually thicker than 5 m; formed as a result of limited aggradation of stream gravel and significant colluviation

**AI<sup>1</sup>** High Level Terrace Sediments (includes White Channel Gravel and equivalent sediments): weathered pebbles to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may be remnant features from the southward flowing paleo-Yukon River drainage system

**PALEOZOIC AND MESOZOIC**

**R** Bedrock: schist, gneiss, ultramafics, granodiorite, monzonite, marble, and basalt; includes areas of thin colluvial cover, blockfields, and sorted stone polygons in alpine areas

### SYMBOLS

Geologic contact: defined, approximate, inferred

Open system pingo, collapsed open system pingo

Thermoclast collapse activity

Landslide movement direction in bedrock and colluvium

Terraces created by widespread landslide movement in drift

Slope scars (flicks on stepped side)

Degraded Cirque: active during pre-Red Glaciations

Degraded Arête: active during pre-Red Glaciations

Meltwater channel: flow direction, unknown flow direction

Large meltwater channel

All time (pre-Red) glacial limit: defined, inferred

Cryoturbation terrace

Tor

Landform Streamlined by glacial ice

Vertebrate fossil locality

Stratigraphic section

Fault trace

Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock slopes

Abandoned valley: paleoflow defined

Abandoned valley: paleoflow undefined

Paleoflow, suspected buried valley

Rock glacier

### DESCRIPTIVE NOTES

The Ladue Creek map area is predominantly underlain by Paleozoic schist and gneiss that is intruded by Mesozoic felsic plutonic rocks (Templeman-Kiul, 1974). Its physiography is dominated by the valley of White River that reaches widths of 4 km. It and the more isolated Ladue River and Ladue Creek represent long established drainage courses that have incised into the Koonkia Plateau. The White River follows relict linear courses. It appears to follow lithological changes such as the margin of a mesozoic plutonic complex above the confluence of Ladue River and Ladue Creek, and the margin of a Mesozoic plutonic complex above the confluence of Ladue River and Ladue Creek. The Ladue River follows a linear course that is parallel to the margin of a Mesozoic plutonic complex. The Ladue River and Ladue Creek are thought to be the result of a Mesozoic plutonic complex. The Ladue River and Ladue Creek are thought to be the result of a Mesozoic plutonic complex. The Ladue River and Ladue Creek are thought to be the result of a Mesozoic plutonic complex.

Natural exposures of surficial deposits are rare in the map area. Dense vegetation covers much of the area. Permanently frozen ground is common in the form of talus and soil. Consequently, surficial sediments have been largely mapped from the interpretation of air photographs. Reconnaissance of the map area is made by field parties to the east where surficial sediments and their stratigraphies are better exposed and understood.

The Ladue Creek map area has never been glaciated with the exception of uplands within the southeastern quadrant that once supported cirque glaciers. Colluvium is the dominant sediment. It is formed by the breakdown of bedrock into silt that is transported down slope by gravitational processes such as solifluction and landsliding. Fluvial deposits are confined to valley bottoms. Stony wedges during glacial periods and the Holocene have deposited extensive deposits of eolian sand along White River and extensive deposits of loess in sheltered steep-side valleys. These have been reworked and mixed with organic sediments to form thick accumulations of muck. They commonly contain extensive bodies of reworked ice.

Organic deposits, formed in bogs and fens, are extensive in valley bottom settings and commonly contain massive lenses of ice. Terraces and valley bottom gravels may locally contain economically viable amounts of placer gold. However, the placer potential of most of the map area is presently unknown.

### REFERENCES

Templeman-Kiul, D.J., 1974. Reconnaissance geology of Ashikh, Snag and part of Stewart River map areas, west-central Yukon; Geological Survey of Canada, Paper 73-41, 87p.

Geology by L.E. Jackson, Jr. (1999 - 2002)

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Digital cartography K. Shimamura, Terrain Sciences Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by Pam Dhas, Earth Sciences Sector Information Division (ESS Info)

Magnetic declination 2005, 25°08' E, decreasing 18.2' annually

Elevations in feet above mean sea level

Contour interval 100 feet

OPEN FILE 4573  
SURFICIAL GEOLOGY  
**LADUE CREEK**  
YUKON TERRITORY

Scale 1:50 000 / Échelle 1/50 000

Universal Transverse Mercator Projection  
North American Datum 1983  
© Her Majesty the Queen in Right of Canada 2005

Projection transversale universelle de Mercator  
Système de référence géodésique nord-américain, 1983  
© Sa Majesté la Reine du chef du Canada 2005

115 N03 115 N08 115 013 115 014 115 015 115 016  
OF4579 OF4580 OF4581 OF4582 OF4583 OF4584

115 N10 115 N09 115 012 115 011 115 010 115 009  
OF4578 OF4577 OF4588 OF4587 OF4586 OF4585

115 N07 115 N08 115 005 115 008 115 007 115 006  
OF4576 OF4575 OF4582 OF4583 OF4584 OF4585

115 N02 115 N01 115 004 115 003 115 002 115 001  
OF4574 OF4573 OF4581 OF4580 OF4579 OF4578

115 K15 115 K16 115 J13 115 J14 115 J15 115 J16  
OF4544 OF4545 OF4546 OF4547

Geology by L.E. Jackson, Jr. (1999 - 2002)

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Digital cartography K. Shimamura, Terrain Sciences Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by Pam Dhas, Earth Sciences Sector Information Division (ESS Info)

Magnetic declination 2005, 25°08' E, decreasing 18.2' annually

Elevations in feet above mean sea level

Contour interval 100 feet