



LEGEND

Note: Map units listed below occur within one metre of the surface. Where organic or colluvial sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that could contain placer gold. In order to accentuate these potentially erodible deposits, a compound map unit is presented, e.g., CEA⁺. This means that colluvial complex sediments overlie alluvial terrace sediments thought to be late Tertiary in age. The legend is part of a larger regional study hence coloured boxes indicate units that appear on this map. In addition, not all symbols in the legend are represented on this map.

CENOZOIC

QUATERNARY

HOLOCENE

Made Land: placer mines, roads, and airstrip

ORGANIC DEPOSITS: peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, ferns, and swamps situated on valley bottoms; permafrost is commonly encountered within 1 m of the surface. Thermokarst collapse is common.

Organic Blanket: unsorted; thickness > 1 m to 5 m

Organic Veneer: blanket bog generally < 1 m thick

ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by streams

Floodplain Sediments: gravel, cobble to pebble; massive to well stratified, capped by sand and silt; flat lying, includes lacustrine and organic deposits in abandoned channels and backswamp areas; subject to periodic inundation and reworking by streams; thickness 1 to 5 m

Alluvial Fan Sediments: gravel, sand, silt, and diatomite, massive to well stratified; sediments form fan-shaped landforms or complexes of coalesced fan-shaped landform at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale

HOLOCENE AND PLEISTOCENE (UNDIVIDED)

COLLUVIAL DEPOSITS: stony diatomite resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and windblowing; colluvial deposits may contain reworked glaciofluvial and moraine sediments within the limits of pre-Holocene cover and reworked older sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating

Colluvial Blanket and Veneer Sediments: diatomite, stony with a sandy matrix, massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous

Colluvial Apron Sediments: bouldery diatomite and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope complex of small steep slope flow and solifluction deposits; thickness < 1 m at the upper and lower slope limit to up to 5 m or more in the thickest part of the apron

Landslide Sediments: silt loam to boulders, poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly

Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fan, colluvial blanket, landside sediments and colluvial drift within the limits of calculation; the unit commonly occurs along the lower slopes of valley margins

Colluvial Eolian Apron (muck): primary deposits of eolian fine sand and silt reworked and transported by wind; organic silt, and detrital, alluvial fan gravel and sand and variable amounts of stony colluvial diatomite; forms aprons along valley bottoms through reintergradation of eolian sediments from valley sides to valley floor; commonly preserved on north-facing slopes; thickness 1 to 30 m; commonly contains segregated bodies of ice and buried ice wedges

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)

ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: gravel, cobble to pebble with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m

Alluvial Fan Sediments: single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m

Alluvial-Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; thin to thick bedded, interstratified with colluvial diatomite; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid-valley locations

EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited by wind action during glaciations and commonly reworked through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout low lying areas

Eolian Blanket: fine sand and silt, well sorted, massive, may form crescent-shaped and linear dunes and features or gently undulating near-stone eolian plains; thickness 1 to 5 m

Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m

LATE PLEISTOCENE - MCCONNELL GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

Glaciofluvial Terrace Sediments: gravel and sand, unweathered, forming one or more terraces

MIDDLE PLEISTOCENE - REID GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

Glaciofluvial Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces

LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - PRE-REID GLACIATIONS (UNDIVIDED)

GLACIOFLUVIAL DEPOSITS: well stratified sand, silt, clay, deposited in lakes ponded by glacial ice

Glaciolacustrine Undivided: sand, silt, and clay, undifferentiated at this scale of mapping

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Glaciofluvial Terrace Sediments: gravel and sand, deeply weathered; incised into flights of terraces; thickness 1 m to > 5 m

MORAINAL DEPOSITS (TILL): glacial diatomite, mainly ill, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Till Blanket: diatomite, stony, silty sand matrix; massive; conforms to underlying topography; thickness > 1 m; extensively colluviated on slopes

Till Veneer: diatomite, stony, silty sand matrix; massive; discontinuous and may contain extensive areas of thin (< 1 m) colluvium

ALLUVIAL DEPOSITS: Gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: sandy pebble and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Holocene or glacial drainage; thickness 1 to 5 m

UNDIFFERENTIATED DRIFT: diatomite, gravel, sand, silt and clay deposited from glacial ice; glacial streams, and glacially dammed lakes; extensive weathering; poor exposure and permafrost make differentiation into component glacial sediments difficult; thicknesses commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and integrated with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Drift: flat to gently sloping

Drift Modified by Landsliding: drift translated along failure plains into irregular steps and sub parallel scarps

Fluvially Incised Drift: formerly extensive areas of drift incised by closely spaced stream valleys

LATE PLEISTOCENE

Basalt: columnar alkali olivine basalt and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Rosebud Creek; thickness 10 m

PLIOCENE AND LATE MIOCENE

ALLUVIAL DEPOSITS: preglacial gravel and sand; highly dissected and deeply weathered

Podium and Bajada Sediments: incised fluvial surfaces which are found at a mid-slope position in unglaciated drainage systems; usually thicker than 5 m; formed as a result of limited aggradation of stream gravel and significant colluviation; composed of thin, poorly sorted gravel that contains both locally derived subangular stream gravel deposits and angular bedrock fragments

High Level Terrace Sediments (includes White Channel Drift and equivalent sediments): weathered pebble to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may have remnant features from the southward-flowing paleo-Yukon River drainage system

PALEOZOIC AND MESOZOIC

Bedrock: schist, gneiss, ultramafics, granitoids, monzonite, marble, and basalt; includes areas of thin colluvial cover, blockfields, and sorted stone polygons in alpine areas

SYMBOLS

Geologic contact, defined, approximate, inferred

Open system pingo, collapsed open system pingo

Thermokarst collapse activity

Landslide movement direction in bedrock and colluvium

Scarp created by widespread landslide movement in drift

Terrace scarp (flicks on sloped side)

Degraded Cirque: active during pre-Holocene Glaciations

Degraded Arête: active during pre-Holocene Glaciations

Meltwater channel: flow direction, collapsed open system pingo

Large meltwater channel

At time (pre-Holocene) glacial limit, defined, inferred

Cryoturbation terrace

Tor

Landform streamlined by glacial ice

Vertebrate fossil locality

Stratigraphic section

Fault trace

Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock slopes

Abandoned valley: paleoflow defined

Abandoned valley: paleoflow undefined

Paleoflow, suspected buried valley

Rock glacier

DESCRIPTIVE NOTES

Ladue River map area is dominated by V-shaped valleys incised up to 300 m into the Klondike Plateau. Bedrock is dominated by Paleozoic schist (Templeman-Kull, 1974). Drainage patterns are rectangular. This pattern likely reflects neotectonic patterns of faulting or other fractures within the plateau complex.

Natural exposures of surficial deposits are rare in the map area. Dense vegetation covering lower slopes and valley bottoms and permanently frozen ground within a few feet of the surface make creation of exposures with hard tools extremely difficult. Consequently, surficial sediments have been largely mapped from the interpretation of air photographs. Reconstruction of late Cenozoic history is made with reference to map areas to the east where surficial sediments and their stratigraphic histories are better exposed and understood.

With the exception of unnamed uplands in the southeastern and southwestern portions of the map area, the region has never been glaciated. Colluvium is the dominant sediment. It is formed by the breakdown of bedrock into regolith that is transported down-slope by gravitational processes such as solifluction and landsliding. Exploration trenches cut into gentle bedrock slopes reveal that rock has been locally disintegrated and finely weathered to one to two feet of depth or more. Degraded cirques occur in the upland mentioned above. These fall within the extensive glaciated area of the Pleistocene glaciations recognized by Nelson and Jackson (2002).

Organic deposits, formed in bogs and ferns, and muck, consisting of reworked eolian silt (loess), are widespread in valley bottom settings and commonly contain massive lenses of ice. They overlie most fluvial deposits in the map area.

Valley bottom and terrace gravel have been extensively mined for placer gold along the southern and of Moosehorn Range. With the exception of this area, the placer potential of the Ladue River map area is unknown.

REFERENCES

Nelson, F.E.H. and Jackson, L.E., Jr.
2002: Cirque forms and slope glaciation during the Pleistocene, west-central Yukon, in Yukon Exploration and Geology 2002, D.S. Edmond and L.L. Lewis (eds.), Exploration and Geological Services Division, Yukon Region, Yukon and Northern Affairs, Canada, p. 185-198.

Templeman-Kull, D.J.
1974: Reconnaissance geology of Ashcroft, Snag and part of Stewart River map areas, west-central Yukon; Geological Survey of Canada, Paper 73-41, 97p.

Geology by L.E. Jackson, Jr. (1999 - 2002)

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Digital cartography by K. Shimamura, Terrain Sciences Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by Pam Ohas, Earth Sciences Sector Information Division (ESS Info)

Magnetic declination 2005, 25°01' E, decreasing 18.1' annually

Elevations in feet above mean sea level

Contour interval 100 feet

OPEN FILE 4574
SURFICIAL GEOLOGY
LADUE RIVER
YUKON TERRITORY
Scale 1:50 000/Echelle 1/50 000

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2005

Projection transversale universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2005

115 N15	115 N18	115 O13	115 O14	115 O15	115 O16
OF4578	OF4580	OF4582	OF4584	OF4586	OF4588
115 N19	115 N18	115 O12	115 O11	115 O10	115 O9
OF4579	OF4577	OF4588	OF4587	OF4587	OF4588
115 N7	115 N8	115 O5	115 O6	115 O7	115 O8
OF4576	OF4578	OF4582	OF4583	OF4584	OF4585
115 N2	115 N1	115 O4	115 O3	115 O2	115 O1
OF4574	OF4573	OF4581	OF4589	OF4588	OF4587
115 N15	115 N16	115 J13	115 J14	115 J15	115 J16
OF4544	OF4545	OF4544	OF4545	OF4546	OF4546