



LEGEND

Note: Map units listed below occur within one metre of the surface. Where organic or eolian sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that could contain placer gold. In order to accentuate these potentially exploitable deposits, a compound map unit is presented, e.g., CwA1T. This means that colluvial complex sediments overlie alluvial terrace sediments thought to be late Tertiary in age. This legend is part of a larger regional study hence coloured boxes indicate units that appear on this map. In addition, not all symbols in the legend are represented on this map.

CENOZOIC

QUATERNARY HOLOCENE

Made Land: placer mines, roads, and airstrip

ORGANIC DEPOSITS: peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, ferns, and swamps situated on valley bottoms; permafrost is commonly encountered within 1 m of the surface. Thermokarst collapse is common.

Organic Blanket: undrained; thickness > 1 m to 5 m

Organic Veneer: blanket bog generally < 1 m thick

ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by streams

Floodplain Sediments: gravel, cobble to pebbles; massive to well stratified, capped by sand and silt; flat lying; includes lacustrine and organic deposits in abandoned channels and backswamp areas; subject to periodic inundation and reworking by floods; thickness 1 to 5 m

Alluvial Fan Sediments: gravel, sand, silt and diatomite; massive to well stratified; sediments form fan-shaped landforms or complexes of coalesced fan-shaped landform at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

Alluvial Complex Sediments: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale

HOLOCENE AND PLEISTOCENE (UNDIVIDED)

COLLUVIAL DEPOSITS: stony diatomite resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits may contain reworked glacial and moraine sediments within the limits of pre-Ried ice cover and reworked eolian sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating

Colluvial Blanket and Veneer Sediments: diatomite, stony with a sandy matrix; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous over bedrock

Colluvial Apron Sediments: bouldery diatomite and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope free of small steep debris flow and solifluction deposits; thickness is < 1 m at the upper and lower slope limit to up to 5 m or more in the thickest part of the apron

Landslide Sediments: silt loam to boulders; poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly

Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fan, colluvial blanket, landslide sediments and colluvial drift within the limits of glaciation; the unit commonly occurs along the lower slopes of valley margins

Colluvial/Eolian Apron (mud): primary deposits of eolian fine sand and silt reworked into massive; sediments form a wedge-like slope free of small steep debris flow and solifluction deposits; thickness is 1 to 20 m; commonly contains segregated bodies of ice and buried ice wedges

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)

ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: gravel, cobble to pebbles with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m

Alluvial Fan Sediments: single fans or aprons of coalesced fans formed of gravel and sand, poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m

Alluvial/Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; thin to thick bedded; interstratified with colluvial diatomite; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid valley locations

EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited by wind action during glaciations and commonly reworked through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout low lying areas

Eolian Blanket: fine sand and silt; well sorted; massive; may form crescent shape and linear dunes and features or gently undulating fine-grained eolian plains; thickness 1 to 5 m

Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m

LATE PLEISTOCENE - MCCONNELL GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

Glacifluvial Terrace Sediments: gravel and sand, unweathered, forming one or more terraces

MIDDLE PLEISTOCENE - REID GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

Glacifluvial Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces

LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - PRE-RIED GLACIATIONS (UNDIVIDED)

GLACIOCLASTIC DEPOSITS: well stratified sand, silt, clay, undifferentiated at this scale of mapping

Glaciolacustrine Undivided: sand, silt, and clay, undifferentiated at this scale of mapping

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Glacifluvial Terrace Sediments: gravel and sand, deeply weathered; incised into flights of terraces; thickness 1 to > 5 m

MORAINAL DEPOSITS (TLL): glacial diatomite, mainly silt, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

TLLPR: diatomite, stony, silt-sand matrix; massive; conforms to underlying topography; thickness > 1 m; extensively colluviated on slopes

TLLVPR: diatomite, stony, silt-sand matrix; massive; discontinuous and may contain extensive areas of thin (< 1 m) colluvium

ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

AI^{PR}: Alluvial Terrace Sediments: sandy pebbles and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Ried glaciers or glacial drainage; thickness 1 to 5 m

UNDIFFERENTIATED DRIFT: diatomite, gravel, sand, silt and clay deposited from glacial ice, glacial streams, and glacially derived lakes; extensive weathering, poor exposure and permafrost make differentiation into component glacial sediments difficult; thicknesses commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and interbedded with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

D^{PR}: Drift: flat to gently sloping

D^{IR}: Drift Modified by Landsliding: drift translated along failure plains into irregular steps and sub parallel segments

De^{PR}: Fluvially Incised Drift: formerly extensive areas of drift incised by closely spaced stream valleys

LATE PLEISTOCENE

Basalt: columnar alkaline olivine basalt and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Rossbud Creek; thickness 10 m

PLEISTOCENE AND LATE PLEISTOCENE

ALLUVIAL DEPOSITS: pegmatite gravel and sand; highly dissected and deeply weathered

P^T: Pediment and Bajada Sediments: inclined fluvial surfaces which are found at a midstage position in unglaciated drainage systems; usually thinner than 5 m; formed as a result of limited aggradation of stream gravel and significant colluviation; composed of thin, poorly sorted gravel that contains both locally derived subangular stream gravel deposits and angular bedrock fragments

AI^T: High Level Terrace Sediments (includes White Channel Gravel and equivalent sediments): weathered pebbles to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may be remnant features from the southward-flowing paleo-Yukon River drainage system

PALEOZOIC AND MESOZOIC

R: Bedrock: schist, gneiss, ultramafics, granodiorite, monzonite, marble, and basalt; includes areas of thin colluvial cover, bioturbation, and sorted stone polygons in alpine areas

SYMBOLS

Geologic contact, defined, approximate, inferred

Open system pingo, collapsed open system pingo

Thermokarst collapse activity

Landslide movement direction in bedrock and colluvium

Scarp created by widespread landslide movement in drift

Terrace scarp (flicks on sloped side)

Degraded Arête: active during pre-Ried Glaciations

Meltwater channel: flow direction, unknown flow direction

Large meltwater channel

At time (pre-Ried) glacial limit; defined, inferred

Cryoturbation terrace

Tor

Landsform Streamlined by glacial ice

Vertebrate fossil locality

Stratigraphic section

Fault trace

Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock slopes

Abandoned valley: paleoflow defined

Abandoned valley: paleoflow undefined

Paleoflow, suspected buried valley

Rock glacier

DESCRIPTIVE NOTES

Rice Creek map area is dominated by V-shaped valleys incised up to 300 m into the Klondike Plateau. Bedrock is dominated by early Mesozoic felsic plutons (Dempsey-Kluke, 1974). Geopatterns are intricate. The pattern likely reflects nonlinear patterns of faulting or other treatments within the plutonic complex.

Natural exposures of surficial deposits are rare in the map area. Dense vegetation covering lower slopes and valley bottoms and permanently frozen ground within a few tens of cm below the surface make creation of exposures with hard tools extremely difficult. Consequently, surficial sedimentary features have been largely mapped from the interpretation of air photographs. Reconnaissance of late Cenozoic history is made with reference to map areas to the east where surficial sediments and their stratigraphies are better exposed and understood.

The region has never been glaciated with the exception of the unnamed upland that trends north-south in the central portion of the map area. It is bounded by a low ridge and small valley glaciers. Colluvium in the upland is formed by the breakdown of bedrock into regolith that is transported down-slope by gravitational processes such as solifluction and landsliding. The degraded cirque occur in the upland, medially down slope that fall within the drainage associated with early Pleistocene glaciations recognized in west-central Yukon by Nelson and Jackson (2002). This glacial limit features excellent evidence of glacial erosion in the form of a 10 m high rim crest of a 10 m high rim crest within the alpine parts of this upland. The surfaces of these features are covered with sorted stone polygons, individual blocks of granodiorite within these features are generally 1 m. These form under periglacial conditions but the specific processes that create them are poorly understood.

Organic deposits in bogs and ferns and musk, consisting of reworked silt and clay, are extensive in valley bottom settings and commonly contain massive lenses of ice. They overlie most fluvial deposits in the map area.

With the exception of Mason Creek, the placer potential of Rice Creek map area is unproven.

REFERENCES

Nelson, P.E.N. and Jackson, L.E. Jr.
2002: Cirque forms and alpine glaciation during the Pleistocene, west-central Yukon: in Yukon Exploration and Geology 2002, D.S. Emond and L.L. Laine (eds.), Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs, Canada, p. 183-186.

Templeman-Kluit, D.J.
1974: Reconnaissance geology of Aishihik, Inigok and part of Stewart River map areas, west-central Yukon; Geological Survey of Canada, Paper 73-41, 9p.



OPEN FILE 4575
SURFICIAL GEOLOGY
RICE CREEK
YUKON TERRITORY
Scale 1:50 000/Échelle 1/50 000

Mètres 1 0 1 2 3 4 Kilomètres

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2005

Projection transverse universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2005

115 N13	115 N16	115 013	115-014	115-015	115-016
OF479	OF480	OF490	OF491	OF492	OF488
115 N10	115 N8	115-012	115-011	115-010	115-09
OF478	OF477	OF489	OF488	OF487	OF486
115 N7	115 N8	115-05	115-08	115-07	115-08
OF475	OF476	OF482	OF483	OF484	OF485
115 N0	115 N1	115-04	115-03	115-02	115-01
OF474	OF473	OF481	OF480	OF479	OF478
115 N15	115 N16	115-013	115-014	115-015	115-016
OF479	OF480	OF490	OF491	OF492	OF488

Geology by L.E. Jackson, Jr. (1999 - 2002)

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Digital cartography: K. Shimamura, Terrain Sciences Division

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatica Canada, modified by Pam Dhais, Earth Sciences Sector Information Division (ESS Info)

Magnetic declination 2005, 25°10' E, decreasing 18.3' annually

Elevations in feet above mean sea level

Contour interval 100 feet

OPEN FILE
DOSSIER PUBLIC
4575

Les dossiers publics ont été publiés par le Service géologique du Canada.

2005

Recommended citation:
Jackson, L.E. Jr.
2005: Surficial Geology, RICE CREEK, Yukon Territory, Geological Survey of Canada, Open File 4575, scale 1:50 000.