



LEGEND
Note: Map units listed below occur within one metre of the surface. Where organic or eolian sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit.

- CENOZOIC
QUATERNARY
HOLOCENE
Made Land: placer mines, roads, and airstrip
Organic Blanket: undrained; thickness > 1 m to 5 m
Organic Veneer: drained bog generally < 1 m thick
ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by streams
Floodplain Sediments: gravel, cobble to pebbles; massive to well stratified, capped by sand and silt
Alluvial Fan Sediments: gravel, sand, silt, and diatomite, massive to well stratified, deposited from fan-shaped landforms or complexes of coalesced fan-shaped landform
Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale
HOLOCENE AND PLEISTOCENE (UNDIVIDED)
COLLUVIAL DEPOSITS: stony diamictum resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits contain weathered glaciofluvial and moraine sediments within the limits of pre-Filly ice cover and reworked eolian sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating
Alluvial Blanket and Veneer Sediments: diamictum, stony with a sandy matrix; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous over bedrock
Colluvial Apron Sediments: bouldery diamictum and bouldery sandy gravel, poorly sorted; massive to poorly stratified; colluvial blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous over bedrock
Landslide Sediments: silt loam to boulders, poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly
Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial and alluvial fans, colluvial blankets, landslide sediments and colluvial aprons within the limits of glaciation; the unit commonly occurs along the lower slopes of valley margins
Colluvial/Eolian Apron (muck): primary deposits of eolian fine sand and silt reworked and interstratified with organic silt, and clastic, alluvial fan gravel and sand in variable amounts of former colluvial diamictum; forms aprons along valley bottoms through reworking of eolian sediments from valley sides to valley floor; commonly preserved on north-facing slopes; thicknesses 1 to 20 m; commonly contains segregated bodies of ice and buried ice wedges
MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)
Alluvial Terrace Sediments: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain glacial pebbles
Alluvial Fan Sediments: single fans or aprons of coalesced fans formed of gravel and sand; poorly to moderately sorted, now isolated from water and debris flows due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m
Alluvial/Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately sorted; thin to thick bedded; interstratified with colluvial diamictum; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in irregular locations
EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited by wind action during glacial periods and commonly reworked through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout the lying area
Eolian Blanket: fine sand and silt, well sorted; massive; may form crescent-shaped and linear dunes and features of gently undulating inter-dune eolian plains; thickness 1 to 5 m
Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas; thickness < 1 m
LATE PLEISTOCENE - McCONNELL GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation
Glaciofluvial Terrace Sediments: gravel and sand, unweathered, forming one or more terraces
MIDDLE PLEISTOCENE - REID GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m
Glaciofluvial Terrace Sediments: gravel and sand, moderately weathered, forming one or more terraces
LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - pre-Reid GLACIATIONS (UNDIVIDED)
GLACIOCLASTIC UNDEPOSITED: sand, silt, and clay; undifferentiated at the scale of mapping
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorphs and sand wedges), and strong chemical weathering
Glaciofluvial Terrace Sediments: gravel and sand, deeply weathered, incised into flights of terraces; thickness 1 to > 5 m

HOLOCENE AND PLEISTOCENE (UNDIVIDED)
COLLUVIAL DEPOSITS: stony diamictum resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits contain weathered glaciofluvial and moraine sediments within the limits of pre-Filly ice cover and reworked eolian sediments; colluvial deposits are products of formation and reworking over a significant part of the Pleistocene and Holocene epochs; surface is commonly hummocky or undulating

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED)
Alluvial Terrace Sediments: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain glacial pebbles

LATE PLEISTOCENE - McCONNELL GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

MIDDLE PLEISTOCENE - REID GLACIATION
GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m

LATE PLEISTOCENE TO MIDDLE PLEISTOCENE - pre-Reid GLACIATIONS (UNDIVIDED)
GLACIOCLASTIC UNDEPOSITED: sand, silt, and clay; undifferentiated at the scale of mapping

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorphs and sand wedges), and strong chemical weathering

GLACIOFLUVIAL TERRACE SEDIMENTS: gravel and sand, deeply weathered, incised into flights of terraces; thickness 1 to > 5 m

MORAINAL DEPOSITS (TILL): glacial diamictum, mainly till, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorphs and sand wedges), and strong chemical weathering

Till Veneer: diamictum, stony, silt-sand matrix; massive; conforms to underlying topography; thickness > 1 m, extensively colluviated on slopes

Till Veneer: diamictum, stony, silt-sand matrix; massive; conforms to underlying topography; thickness > 1 m, extensively colluviated on slopes

ALLUVIAL DEPOSITS: Gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain glacial pebbles

ALLUVIAL TERRACE SEDIMENTS: sandy pebbles and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Filly glaciers or glacial drainage; thickness 1 to 5 m

UNDIFFERENTIATED DRIFT: diamictum, gravel, sand, silt and clay deposited from glacial ice, glacial streams, and glacially dammed lakes; extensive weathering; poor exposure and preservation; may have experienced several cycles of alluviation and erosion; thicknesses commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and intergraded with colluvial surface soils; may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorphs and sand wedges), and strong chemical weathering

Drift: flat to gently sloping

Drift Modified by Landsliding: drift translocated along failure planes into irregular steps and sub parallel scarps

Fluvially Incised Drift: formerly extensive areas of drift incised by closely spaced stream valleys

LATE PLEISTOCENE
Basalt: columnar, alkaline olivine basalt and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Flossbeck Creek; thickness 10 m

PLEISTOCENE AND LATE MOGENSE
ALLUVIAL DEPOSITS: preglacial gravel and sand; highly dissected and deeply weathered

Pediment and Rapids Sediments: inclined fluvial surfaces which are found at a midstage position in unglaciated drainage systems; usually thinner than 5 m; formed as a result of limited aggradation of stream gravel and significant colluviation; composed of thin, poorly sorted gravel that contains both locally derived subangular stream gravel deposits and angular bedrock fragments

High Level Terrace Sediments (includes White Channel Gravel and equivalent sediments): weathered pebbles to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts; frequent signs of cryoturbation (ice wedge pseudomorphs and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may be remnant features from the southward-flowing paleo-Yukon River drainage system

PALEOZOIC AND MESOZOIC
Bedrock: schist, gneiss, ultramafic, granodiorite, metabasite, mafic, and basalt; igneous areas of thin colluvial cover, blockfields, and sorted stone polygons in alpine areas

SYMBOLS
Geologic contact: defined, approximate, inferred
Open system pingo, collapsed open system pingo
Thermokarst collapse activity
Landslide movement direction in bedrock and colluvium
Scarps created by widespread landslide movement in drift
Terrace scarp (folds on sloped side)
Degraded Active: active during pre-Reid Glaciations
Degraded Inactive: active during pre-Reid Glaciations
Meltwater channel: flow direction, unknown flow direction
Large meltwater channel
All time (pre-Reid) glacial limit; defined, inferred
Cryoturbation terrace
Tor
Landform Streamlined by glacial ice
Vestibular fossil locality
Stratigraphic section
Fault trace
Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock slopes
Abandoned valley: paleoflow defined
Paleoflow, suspected buried valley
Rock glacier

DESCRIPTIVE NOTES
The physiography of the Crag Mountain map area is dominated by 'V' and 'U' shaped valleys incised up to 300 m into the Kootenai plateau. Bedrock is dominated by Paleozoic gneiss and Mesozoic felsic plutons (Mortensen 1998). Drainage ridges from north-south to east-west.

REFERENCES
Lewy, G.W.
2002: Glaciation, gravel and gold in the Filly Mile Creek area, west-central Yukon; in Yukon Exploration and Geology 1999, D.S. Emond and L.H. Weston (eds.), Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs, Canada, p. 189-200.
Nelson, F.E.H. and Jackson, L.E. Jr.
2002: Cirque forms and alpine glaciation during the Pleistocene, west-central Yukon; in Yukon Exploration and Geology 2002, D.S. Emond, L.L. Lewis (eds.), Exploration and Geological Services Division, Yukon Region, Indian and Northern Affairs, Canada, p. 183-188.
Mortensen, J.K.
1996: Geological compilation maps of the northern Stewart River map area, Kootenai and Elktonia districts (153N/15, 16, 150N/13, 14, and parts of 150N/15, 16; Indian and Northern Affairs Canada, Northern Affairs: Yukon region; map scale 1:50 000.

OPEN FILE 4579
SURFICIAL GEOLOGY
CRAG MOUNTAIN
YUKON TERRITORY
Scale 1:50 000/Echelle 1/50 000
Geology by L.E. Jackson, Jr. (1999-2002)
Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP
Digital cartography by Shimamura, Terrain Services Division
Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada
Magnetic declination 2005, 25°28' E, decreasing 18.7' annually
Elevations in feet above mean sea level
Contour Interval 100 feet



UNIVERSAL TRANSVERSE MERCATOR PROJECTION
North American Datum 1983
Système de coordonnées géographiques nord-américain, 1983
© Her Majesty the Queen in Right of Canada 2005

Canada logo

Grid coordinate table with columns for UTM Easting and Northing coordinates.

OPEN FILE DOSSIER PUBLIC 4579
Geological Survey of Canada
2005

Recommended citation:
Jackson, L.E. Jr.
2005: Surficial Geology, CRAG MOUNTAIN, Yukon Territory; Geological Survey of Canada, Open File 4579, scale 1:50 000.