GEOLOGICAL SURVEY OF CANADA CARTNAT Natural Resources Ressources naturelles Canada Canada's National Geoscience Mapping Program Le Programme national de cartographie géoscientifique du Canada 139°30' 140°00' Bench gravels mined to underlying bedrock strath; upper (north) end of pit Thinly bedded (up to 5 cm) to laminated stony silt and stony sand; thin discontinuous layer of unidentified tephra 1.2 m below surface; stones up to 3 cm; angular, locally derived from adjacent slope Thinly bedded fine to medium sand; base not seen; bedrock contact estimated to be at 5 m depth Road cut through high terrace gravel Massive, poorly sorted, matrix-supported sandy pebble gravel; clasts rounded to sub-angular; angular clasts present appear to be frost shattered; maximum clast size 15 cm; gravel 10YR 5/6; blebs of aeolian silt within gravel apparently mixed in by cryoturbation (7.5YR 4/4); chert and chert pebble conglomerate present many clasts have 7.5YR 4/4 coloured clay skins; appears that a Wounded Moose-type paleosol has been overturned and mixed by cryoturbation; some schist clasts have been reduced to sand but most are quite fresh Vertical exposure on east side of roughly N-S trench in a placer mine Section 3 4 m Laminated to thinly bedded pebbly silt and pebbly sand; scattered lenses of very fine pebble gravel; clasts angular, locally derived Cb-v Blebs and thin beds of Gold Run tephra interbedded bedded with silt and very fine sand; Gold Run tephra dated at 740 + /- 60 ka (John Westgate, University of Toronto, 2004 personal communication) Crudely laminated to laminated and cross-laminated sandy silt with scattered pebbles and granules; dark greyish brown (10YR 3/2) Crudely laminated silt, silty sand and pebbly sandy silt; very dark greyish brown (10YR 3/2); locally organic rich; pebbles are angular; contains at least two very thin unidentified tephra laminae; abrupt basal contact Oxidized, crudely stratified sandy pebble gravel; dark yellowish brown (10YR 4/6); pebbles rounded to subangular; clasts up to 15 cm; average clast size 5 cm; appears to be imbricated; locally derived schist make up angular clasts Cliff-top trench (5 m wide) approximately 76 m above Yukon River; cliff appears to intersect former channels cut in strath Ranges from pebbly coarse sand to sandy cobbly pebble gravel over exposure; clasts up to 15 cm (rare); most clasts 10 cm; rounded to subrounded; contains angular blocks of schist ripped up from underlying bedrock; lower contact rregular; pebbly fabric indicates paleoflow from north Weathered schist bedrock STRATIGRAPHIC LEGEND Note: Not all stratigraphic units from the legend are present on stratigraphic logs HOLOCENE (< 0.010 Ma) LATE PLIOCENE AND EARLY PLEISTOCENE (2.7 - 0.780 Ma) Paleosol developed in late Pliocene to early Pleistocene glaciofluvial and non-glacial sediments Stratified sand and gravel; glaciofluvial sediments deposited Stratified sand and gravel; alluvial sediments during older pre-Reid glaciations LATE PLEISTOCENE TO EARLY HOLOCENE (< 0.125 Ma) Stratified to massive diamicton; till deposited during one of Silt (organic rich), peat, and organic detritus, and extensive several older pre-Reid glaciations interstratified and segregated ice; collectively called muck Fine sand, silt and clay; lacustrine or slack water fluvial Massive to stratified diamicton; colluvial sediments (may locally Stratified to massive sand and gravel; non-glaciofluvial sediments, may be graded to pre-Reid outwash Massive to stratified silt and fine sand; eolian sediments PLIOCENE (pre-glacial, 5 - 2.7 Ma) Stratified silt and sand; resedimented eolian sediments locally Basalt and basalt breccia interstratified with alluvial fan sediments Stratified to massive gravel and sand; White Channel Gravel Stratified sand and gravel; alluvial sediments and equivalent clastic units predating regional glaciation, includes late Tertiary pediment sediments MIDDLE PLEISTOCENE (0.780 - 0.125 Ma) PRE-PLIOCENE (> 5 Ma) Paleosol developed in Reid and younger pre-Reid glaciofluvial Mesozoic and Paleozoic bedrock sediments Stratified sand and gravel; glaciofluvial sediments deposited Tephra - identification and age, if known, described during Reid glaciation on stratigraphic log Stratified sand and gravel; glaciofluvial sediments deposited during younger pre-Reid glaciations Ice-wedge pseudomorph or sand wedge Massive to stratified silt and fine sand; primary and resedimented eolian sediments 139°30' **OPEN FILE 4589** SURFICIAL GEOLOGY **OGILVIE** 115 N/15 | 115 N/16 | 115-0/13 | 115-0/14 | 115-0/15 | 115-0/16 OF4579 OF4580 OF4590 OF4591 OF4592 OF4593 YUKON TERRITORY 115 N/10 115 N/9 115-0/12 115-0/11 115-0/10 115-0/9 Scale 1:50 000/Échelle 1/50 000 OF4578 OF4577 OF4589 OF4588 OF4587 OF4586 115 N/7 | 115 N/8 | 115-0/5 | 115-0/6 | 115-0/7 | 115-0/8 OF4575 | OF4576 | OF4582 | OF4583 | OF4584 | OF4585 Universal Transverse Mercator Projection North American Datum 1983 Système de référence géodésique nord-américain, 1983 © Her Majesty the Queen in Right of Canada 2005 © Sa Majesté la Reine du chef du Canada 2005 115 N/2 115 N/1 115-0/4 115-0/3 115-0/2 115-0/1 OF4574 OF4573 OF4581 OF4349 OF4348 OF4347 115 K/15 | 115 K/16 | 115 J/13 | 115 J/14 | 115 J/15 | 115 J/16

LEGEND

Note: Map units listed below occur within one metre of the surface. Where organic or eolian sediments < 1 m thick overlie these, a pattern is overlaid upon the map unit. Along some valleys, colluvial or alluvial sediments > 1 m thick overlie older alluvial gravels that could contain placer gold. In order to accentuate these potentially exploitable deposits, a compound map unit is presented, e.g., Cx/AtT. This means that colluvial complex sediments overlie alluvial terrace sediments thought to be late Tertiary in age. This legend is part of a larger regional study hence coloured boxes indicate units that appear on this map. In addition, not all symbols in the legend are represented on this map.

HOLOCENE

Made Land: placer mines, roads, and airstrip

ORGANIC DEPOSITS: peat and organic silt formed predominantly by the accumulation of vegetative material in bogs, fens, and swamps situated on valley bottoms; permafrost is commonly encountered within 1 m of the surface. Thermokarst

Organic Blanket: undivided; thickness > 1 m to 5 m

Organic Veneer: blanket bog generally < 1 m thick ALLUVIAL DEPOSITS: gravel to silt size sediments, well stratified, deposited by

Floodplain Sediments: gravel, cobble to pebble; massive to well stratified, capped by Ap sand and silt; flat lying; includes lacustrine and organic deposits in abandoned channels and backswamp areas; subject to periodic inundation and reworking by

floods; thickness 1 to 5 m Alluvial Fan Sediments: gravel, sand, silt, and diamicton, massive to well stratified; Af sediments form fan-shaped landforms or complexes of coalesced fan-shape landform at the confluence of tributary streams; may be subject to flooding accompanied by sudden stream migration and inundation; thickness up to 10 m

> Alluvial Sediments Complex: sediments forming floodplains, fans, and terraces that cannot be subdivided at this map scale

> HOLOCENE AND PLEISTOCENE (UNDIVIDED) COLLUVIAL DEPOSITS: stony diamicton resulting from the physical and chemical breakdown of bedrock and subsequent reworking and transportation by creep, solifluction, and landsliding; colluvial deposits may contain reworked glaciofluvial and morainal sediments within the limits of pre-Reid ice-cover and reworked eolian sediments; colluvial deposits are products of formation and reworking over a significant part of the Pielstocene and Holocene epochs; surface is commonly hummocky or undulating

Colluvial Blanket and Veneer Sediments: diamicton, stony with a sandy matrix; massive to poorly stratified; colluviated blankets generally conform to underlying bedrock and exceed 1 m in thickness; veneers are < 1 m in thickness and are commonly discontinuous over bedrock

Colluvial Apron Sediments: bouldery diamicton and bouldery sandy gravel, poorly sorted; massive; sediments form a wedge-like slope-toe complex of small steep debris flow and solifluction deposits; thickness is < 1 m at the upper and lower slope limit to up to 5 m or more in the thickest part of the apron

Landslide Sediments: silt loam to boulders, poorly sorted to unsorted; massive; clasts are subangular to angular and are locally derived; thickness varies greatly Colluvial Complex Sediments: areas of intergrading colluvial and alluvial sediments which are too complex to subdivide at the scale of mapping; unit may include colluvial

and alluvial fan, colluvial blanket, landslide sediments and colluviated drift within the

limits of glaciation; the unit commonly occurs along the lower slopes of valley margins Colluvial/Eolian Apron (muck): primary deposits of eolian fine sand and silt CEaP resedimented and interstratified with organic silt, and detritus, alluvial fan gravel and sand and variable amounts of stony colluvial diamicton; forms aprons along valley bottoms through resedimentation of eolian sediments from valley sides to valley floor, commonly preserved on north-facing slopes; thickness 1 to 20 m; commonly contains segregated bodies of ice and buried ice wedges

MIDDLE TO LATE PLEISTOCENE (UNDIVIDED) ALLUVIAL DEPOSITS: gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: gravel, cobble to pebble with a sandy matrix; massive to well stratified; capped by sand and silt; sediments are of flood plain origin now isolated from flooding by stream incision; thickness 1 m to 10 m Alluvial Fan Sediments: single fans or aprons of coalesced fans formed of gravel and

AfP sand, poorly to moderately sorted, now isolated from water and debris floods due to fluvial incision; sediments disturbed by cryoturbation; thickness up to 10 m Alluvial/Colluvial Complex Sediments: silt, sand and gravel, poorly to moderately ACXP sorted; thin to thick bedded, interstratified with colluvial diamicton; sediments underlie the floors and margins of narrow upland valleys and grade laterally up slope into colluvial blankets; sediments may represent several depositional cycles; thickness may exceed 10 m in mid-valley locations EOLIAN DEPOSITS: well sorted medium sand to silt initially transported and deposited

by wind action during glaciations and commonly resedimented through fluvial and colluvial processes; deposits of very fine sand and coarse silt < 1 m thick are distributed discontinuously throughout low lying areas Eolian Blanket: fine sand and silt, well sorted; massive; may form crescent-shape and Eb^P linear dunes and featureless or gently undulating inter-dune eolian plains; thickness 1

Eolian Veneer: thin deposits of very fine sand and coarse silt distributed discontinuously throughout low lying areas, thickness < 1 m LATE PLEISTOCENE - McCONNELL GLACIATION GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display poor soil development with rare cryoturbation

> Glaciofluvial Terrace Sediments: gravel and sand, unweathered, forming one or more MIDDLE PLEISTOCENE - REID GLACIATION

GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice; deposits display moderate soil development with signs of cryoturbation; soil thickness < 0.5 m Glaciofluvial Terrace Sediments: gravel and sand, moderately weathered, forming one

LATE PLIOCENE TO MIDDLE PLEISTOCENE - pre-Reid GLACIATIONS (UNDIVIDED) GLACIOLACUSTRINE DEPOSITS: well stratified sand, silt, clay, deposited in lakes ponded by glacial ice

> Glaciolacustrine Undivided: sand, silt, and clay; undifferentiated at this scale of GLACIOFLUVIAL DEPOSITS: gravel and sand deposited by streams flowing away from glacial ice in meltwater channels and outwash plains; massive to well stratified.

Surface soils may extend to 2 m depth with well developed clay skins on clasts,

frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and

Glaciofluvial Terrace Sediments: gravel and sand, deeply weathered; incised into flights of terraces; thickness 1 to > 5 m

strong chemical weathering

NATIONAL TOPOGRAPHIC SYSTEM REFERENCE AND INDE TO ADJOINING GEOLOGICAL SURVEY OF CANADA MAPS

Geology by L.E. Jackson, Jr. (1999 - 2002), S.R. Morison and C. Mougeot (1998)

Digital cartography K. Shimamura, Terrain Sciences Division

Co-ordinated through the auspices of the Ancient Pacific Margin NATMAP

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by Parm Dhesi, Earth Sciences Sector Information Division (ESS Info)

Magnetic declination 2005, 25°33' E, decreasing 18.8' annually Elevations in feet above mean sea level

Contour interval 100 feet

MORAINAL DEPOSITS (TILL): glacial diamicton, mainly till, generally consisting of a matrix ranging from sand to clay that supports clasts ranging from boulders to pebbles in size; deposited either directly from glacial ice or by gravity flow from glacial ice; surface soils may extend to 2 m depth with well developed clay skins on clasts, frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Till Blanket: diamicton, stony, silty-sand matrix; massive; conforms to underlying topography, thickness > 1 m; extensively colluviated on slopes

Till Veneer: diamicton, stony, silty-sand matrix; massive; discontinuous and may contain extensive areas of thin (< 1 m) colluvium

ALLUVIAL DEPOSITS: Gravel and sand deposited by streams that were not fed by glacial meltwater; sediments may have experienced several cycles of alluviation and erosion, but are now inactive due to burial or fluvial incision; basal gravels within these sediments commonly contain placer gold

Alluvial Terrace Sediments: sandy pebble and cobble gravel deposited by streams having a fluvial source but graded to the margins of pre-Reid glaciers or glacial drainage; thickness 1 to 5 m

> UNDIFFERENTIATED DRIFT: diamicton, gravel, sand, silt and clay deposited from glacial ice, glacial streams, and glacially damned lakes; extensive weathering, poor exposure and permafrost make differentiation into component glacial sediments difficult; thicknesses commonly exceed 10 m and mask underlying bedrock topography; commonly colluviated and intergraded with colluvium; surface soils may extend to 2 m depth with well developed clay skins on clasts, frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering

Drift: flat to gently sloping

Drift Modified by Landsliding: drift translated along failure plains into irregular steps

and sub parallel scarps

Fluvially Incised Drift: formerly extensive areas of drift incised by closely spaced stream

Basalt: columnar alkaline olivine basalt and flow breccia; erosional remnants of formerly valley filling flows underlying terraces along lower Rosebud Creek; thickness

PLIOCENE AND LATE MIOCENE ALLUVIAL DEPOSITS: preglacial gravel and sand; highly dissected and deeply

Pediment and Bajada Sediments: inclined fluvial surfaces which are found at a midslope position in unglaciated drainage systems; usually thinner than 5 m; formed as a result of limited agradation of stream gravel and significant colluviation; composed of thin, poorly sorted gravel that contains both locally derived subangular stream gravel deposits and angular bedrock fragments

sediments): weathered pebble to cobble gravel > 1 m thick; surface soils may extend to 2 m depth with well developed clay skins on clasts, frequent signs of cryoturbation (ice wedge pseudomorph and sand wedges), and strong chemical weathering; within the Yukon River valley, terraces above the 500 m contour may be remnant features from the southward-flowing paleo-Yukon River drainage system

High Level Terrace Sediments (includes White Channel Gravel and equivalent

PALEOZOIC AND MESOZOIC Bedrock: schist, gneiss, ultramafics, granodiorite, monzonite, marble, and basalt: includes areas of thin colluvial cover, blockfields, and sorted stone polygons in alpine

SYMBOLS

SYMBOLS
Geologic contact; defined, approximate, inferred
Open system pingo, collapsed open system pingo
Thermokarst collapse activity
Landslide movement direction in bedrock and colluvium
Scarps created by widespread landslide movement in drift
Terrace scarp (ticks on sloped side)
Degraded Cirque: active during pre-Reid Glaciations
Degraded Arête: active during pre-Reid Glaciations
Meltwater channel: flow direction, unknown flow direction
Large meltwater channel
All time (pre-Reid) glacial limit; defined, inferred
Cryoplanation terrace
Tor
Landform Streamlined by glacial ice
Vertebrate fossil locality
Stratigraphic section
Fault trace
Lineaments (fault, fracture, joint system) defined by linear drainage courses, aligned gaps in ridges, or aligned breaks in bedrock slopes
Abandoned valley: paleoflow defined
Abandoned valley: paleoflow undefined
Paleoflow, suspected buried valley
Rock glacier

DESCRIPTIVE NOTES

The Ogilvie map area lies within the Klondike Plateau, an incised rolling upland predominantly underlain by Paleozoic schist and gneiss (Bostock 1942). Natural exposures of surficial deposits are confined to cliffs along Yukon River and artificial exposures created by placer mining activities along Sixty Mile River and Ten Mile Creek. Dense vegetation covers much of the area. Permanently frozen ground is often only a few tens of centimeters below the surface making digging with hand tools difficult. Consequently, surficial sediments have been largely mapped from the interpretation of airphotographs. The Ogilvie map area has never been glaciated with the exception of features interpreted as eroded cirques on the highest summits in the northwest quadrant of the map area. Consequently, colluvium, covering slopes and ridges, is the dominant sediment. It is formed by the breakdown of bedrock into regolith that is transported down-slope by Fluvial deposits are confined to terraces and valley bottoms. The oldest (At¹) consist of gravel and sand that cap the highest terraces along Yukon River and high terraces along tributaries such as Sixty Mile River, Rosebute, Reindeer, and Lucky Joe creeks. Measurements of clast imbrications within gravel capping a terrace 77 m above the west bank of Yukon River immediately south of the mouth of Sixty Mile River (section 4) indicate that the Yukon River originally flowed south in this area. Reversal of Yukon River to its present flow direction is postulated to have occurred as a result of the first regional glaciation of southern and central Yukon Territory during the Late Pliocene Epoch ca. 3.1 to 2.6 million years ago (Froese et al. 2000, 2001; Duk-Rodkin et al. 2001) Following the reversal of Yukon River, a succession of terraces were cut and a succession of overlying terrace gravels deposited (At^r). Fission track dating of tephra present within colluvial sediments overlying a terrace 18 m above the present Sixty Mile River flood plain indicates that this terrace was cut prior to about 700 ka. Strong winds during glacial periods and the Holocene have deposited extensive loess and sand on terraces and sediments to form thick accumulations called muck. They commonly contain extensive bodies of segregated ice. Organic deposits in bogs and fens are extensive in valley bottom settings and commonly also contain massive lenses Placer gold has been mined from gravel underlying Ten Mile Creek and from terrace gravel along Sixty Mile River below the mouth of Ten Mile Creek. However, the placer potential of most of the map area is presently unproven.

REFERENCES

1942: Ogilvie; Geological Survey of Canada map 711A, 1:253 440.

Duk-Rodkin, A. Barendregt, R.W., White, J.M., and Singhroy, V.H.

2001: Geologic evolution of the Yukon River: implications for placer gold; Quaternary International, v. 82, p. 5-31. Froese, D.G., Barendregt, R.W., Enkin, R.J., and Baker, J.

2000: Paleomagnetic evidence for multiple late Pliocene-early Pleistocene glaciations in the Klondike area, Yukon Territory; Canadian Journal of Earth Sciences, v. 37, p.863-877. Froese, D.G. Ager, T, Duk-Rodkin, A., Westgate, J. and White, J. and Smith, D. 2001: Drainage reversal and integration of the Pliocene upper Yukon River (3.1-2.6 Ma): key new evidence. Abstracts, Arctic Workshop, Amherst, Massachusetts

OPEN FILE

Open files are products DOSSIER PUBLIC that have not gone through the GSC formal publication process. Les dossiers publics sont GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA

Das été soumis au 2005 publication de la CGC.

Recommended citation: Jackson, L.E., Jr., Morison, S.R., and Mougeot, C. 2005: Surficial Geology, OGILVIE, Yukon Territory; Geological Survey of Canada, Open File 4589, scale 1:50 000.