



LEGEND

This legend is common to Open File 4684 and 4701. Coloured legend blocks indicate map units that appear on this map. Not all map symbols shown in the legend necessarily appear on this map.

SURFICIAL DEPOSITS

QUATERNARY

HOLOCENE

COLLUVIUM: block and rubble accumulations, 1–50 m thick.

Ca Talus: active block and rubble accumulations as much as 50 m thick forming talus (scree) aprons and fans below cliffs resulting from rock falls and debris flows; commonly crossed by debris flow channels and levees.

Cr Rock glacial debris: talus, generally 10–50 m thick, deformed by active flow of interstitial or buried ice to form rock (felsen) glaciers with transverse ridges and furrows, and pits, and with steep, unstable sides and fronts.

FLUVIAL SEDIMENTS: alluvium: gravel and sand, 2–20 m thick.

Ap Alluvial plains: active braided floodplains; includes active proglacial outwash.

At Alluvial terraces: gravel and sand, 2–20 m thick.

Af Alluvial fans: gravel and sand, 2–20 m thick.

MARINE AND GLACIAL MARINE SEDIMENTS: gravel, sand, silt, and clay, 1–20 m thick, deposited in deltaic and beach environments during regression of the postglacial sea.

Mr Beach sediments: gravel and sand, 1–5 m thick, forming ridges and swales.

Mt Deltaic sediments: clay, silt, sand, and gravel, 5–20 m thick, forming coarsening upward sequences under dissected terraces.

Mv Deepwater proglacial silt veneers: silt, clay silt, and fine sand with dropstones, 1–2 m thick.

Mb Deepwater proglacial silt blankets: silt, clay silt, and fine sand with dropstones and minor gravel, 2–10 m thick.

GLACIAL LACUSTRINE SEDIMENTS: clay, silt, sand, and gravel deposited in glacial dammed lakes in deepwater, beach, and deltaic environments.

Lt Deltaic sediments: clay, silt, sand, and gravel, 5–20 m thick, forming coarsening upward sequences under dissected terraces.

Lv Deepwater proglacial silt veneers: silt, clay silt, and fine sand with dropstones, 1–2 m thick.

Lb Deepwater proglacial silt blankets: silt, clay silt, and fine sand with dropstones, 2–5 m thick.

GLACIOFLUVIAL SEDIMENTS: gravel and sand, 1–10 m thick, deposited behind, at, and in front of the ice margin.

Gp,lf Proglacial outwash: gravel and sand, 1–10 m thick, forming broad floodplains. Gp: terraces, Gf: fans, Gf.

Gr,h Ice contact stratified drift: gravel and sand, 1–5 m thick, forming eskers, G; and kames, Gh.

EARLY HOLOCENE AND WISCONSINAN

TILL: nonsorted stony muds, 0.5–60 m thick, deposited in subglacial and ice marginal environments; lithic composition generally reflects underlying bedrock.

Tm End moraines: 5–60 m high, composed of or mented by till, extensively kettled in places; large features mainly covered by debris-rich relict glacier ice.

Tv Till veneer: 0.5–2 m thick and discontinuous.

Tvw Washed till veneer: 0.5–2 m thick, surface armoured by stones due to washing by subglacial meltwater.

Tb Till blanket: 2–10 m thick forming an undulating blanket with drumlins and ribbed moraines in places.

Tbr Ribbed till blanket: 2–10 m thick forming ribbed (Rogen) moraines.

BEDROCK

PRE-QUATERNARY

R ROCK: rock of various compositions and ages (Jackson and Sangster, 1987) variously modified by glacial erosion during the Quaternary and with patchy till cover; hilly and hummocky surfaces, ice moulded in places, with late basins in subglacially scoured regions; smooth surfaces exhibiting little or no sign of glacial erosion in peninsular interiors (Dyke, 1993); cliffs resulting from glacial over-steepening; in places veneered by thin till, commonly boundary.

Geological boundary (defined, assumed) ...

Areas covered by periglacial icefields during the Little Ice Age (indicated by a white pattern) ...

Glacial lake spillway ...

Glacial lake limit ...

Marine limit ...

Marine limit elevation in metres ... 60

Weakly developed strandline ...

Cliff in bedrock ...

Lateral meltwater channel; barb on upstope side ...

Subglacial and proglacial meltwater channel (large, small) ...

Esker ...

Kame ...

Ice contact face ...

Ribbed moraine ...

Lateral moraine ...

End moraine ...

Margin of glacial dispersal train; teeth toward silt, steep side of teeth face down ice ...

Lateral sliding boundary; teeth on sliding side ...

Ice-based ice on other side; steep side of teeth face down ice ...

Iceberg scour ...

Drumlinoid hill ...

Crag-and-tail ...

Ice moulded bedrock ...

Striae (ice flow direction known, unknown) ...

Crossed striae (numbers indicate relative age, 1 being the oldest) ...

Field observation site: boundary distinction (bd), boundary gravel (bg), damiction (d), gravel (g), gravelly sand (gs), mud (m), muddy sand (ms), rock (r), sand (s), sandy gravel (sg), stony mud (sm), silt (t) ...

Field observation site: material as above near rock outcrop ...

Radiocarbon date ...

REFERENCES

Dyke, A.S., 1983: Landscapes of cold-centred Late Wisconsinan ice caps, Canadian Arctic; Progress in Physical Geography, v. 7, p. 225–247.

Jackson, G.D. and Sangster, D.F., 1987: Geology and resource potential of a proposed national park, Bylot Island and northwest Baffin Island, Northwest Territories; Geological Survey of Canada, Paper 87-17, 87 p.



OPEN FILE 4684
SURFICIAL GEOLOGY
LOWER RAVN RIVER
BAFFIN ISLAND
NUNAVUT

Author: A.S. Dyke
Geology by A.S. Dyke, 2004
Field data provided by De Beers Canada Corporation, 2003
Digital cartography by M.M. Proulx, Earth Sciences Sector Information Division (ESS Info)

This map was produced from processes that conform to the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001:2000 standard
Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by ESS Info
Locational accuracy of the base appears to be ±100 m based on plotting of GPS measured field site locations
Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area
Mean magnetic declination 2005, 42°39'W, decreasing 42.7' annually
Elevations in metres above mean sea level
Contour interval 20 m
Field altimetry and the placement and trend of raised shorelines may conflict significantly with the contours

Scale 1:50 000/Échelle 1/50 000
Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2005

37 113	37 114	37 115	37 116
OF4685	OF4686	OF4687	OF4688
37 117	37 118	37 119	37 120
OF4689	OF4690	OF4691	OF4692
37 121	37 122	37 123	37 124
OF4693	OF4694	OF4695	OF4696
37 125	37 126	37 127	37 128
OF4697	OF4698	OF4699	OF4700
37 129	37 130	37 131	37 132
OF4701	OF4702	OF4703	OF4704
37 133	37 134	37 135	37 136
OF4705	OF4706	OF4707	OF4708

OPEN FILE DOSSIER PUBLIC 4684

Les données de ce produit ne sont pas protégées par le droit de la propriété intellectuelle. Elles sont disponibles en français et en anglais.

2005

Recommended citation:
Dyke, A.S., 2005: Surficial geology, Lower Ravn River, Baffin Island, Nunavut; Geological Survey of Canada, Open File 4684, scale 1:50 000.