



- LEGEND**
- This legend is common to Open Files 4685 to 4701.  
Coloured legend blocks indicate map units that appear on this map.  
Not all map symbols shown in the legend necessarily appear on this map.
- SURFICIAL DEPOSITS**
- QUATERNARY**
- HOLOCENE**
- COLLUVIUM:** block and rubble accumulations, 1–50 m thick.
- Talus:** active block and rubble accumulations as much as 50 m thick forming talus (scree) aprons and fans below cliffs resulting from rock falls and debris flows; commonly crossed by debris flow channels and levees.
- FLUVIAL SEDIMENTS:** alluvium; gravel and sand, 2–20 m thick.
- Alluvial plains:** active braided floodplains; includes active proglacial outwash.
- Alluvial terraces:** gravel and sand, 2–20 m thick.
- Alluvial fans:** gravel and sand, 2–20 m thick.
- MARINE AND GLACIAL MARINE SEDIMENTS:** gravel, sand, silt, and clay, 1–20 m thick, deposited in deltaic and beach environments during regression of the postglacial sea.
- Beach sediments:** gravel and sand, 1–5 m thick, forming ridges and swales.
- Deltaic sediments:** clay, silt, sand, and gravel, 5–20 m thick, forming coarsening upward sequences under dissected terraces.
- Deepwater proglacial silt veneers:** silt, clay silt, and fine sand with dropstones, 1–2 m thick.
- Deepwater proglacial silt blankets:** silt, clay silt, and fine sand with dropstones and minor gravel, 2–10 m thick.
- GLACIAL LACUSTRINE SEDIMENTS:** clay, silt, sand, and gravel deposited in glacier dammed lakes in deepwater, beach, and deltaic environments.
- Deltaic sediments:** clay, silt, sand, and gravel, 5–20 m thick, forming coarsening upward sequences under dissected terraces.
- Deepwater proglacial silt veneers:** silt, clay silt, and fine sand with dropstones, 1–2 m thick.
- Deepwater proglacial silt blankets:** silt, clay silt, and fine sand with dropstones, 2–5 m thick.
- GLACIOFLUVIAL SEDIMENTS:** gravel and sand, 1–10 m thick, deposited behind, at, and in front of the ice margin.
- Proglacial outwash:** gravel and sand, 1–10 m thick, forming braided floodplains. Gp: terraces, Gt: and fans, Gt.
- Ice contact stratified drift:** gravel and sand, 1–5 m thick, forming eskers, Gt; and kames, Gh.
- EARLY HOLOCENE AND WISCONSINAN**
- TILL:** nonsorted stony muds, 0.5–60 m thick, deposited in subglacial and ice marginal environments; lithic composition generally reflects underlying bedrock.
- End moraines:** 5–60 m high, composed of or mantled by till, extensively kettled in places; large features mainly corred by debris-rich relict glacier ice.
- Till veneer:** 0.5–2 m thick and discontinuous.
- Washed till veneer:** 0.5–2 m thick, surface armoured by stones due to washing by subglacial meltwater.
- Till blanket:** 2–10 m thick forming an undulating blanket with drumlins and ribbed moraines in places.
- Ribbed till blanket:** 2–10 m thick forming ribbed (Rogen) moraines.
- BEDROCK**
- PRE-QUATERNARY**
- ROCK:** rock of various compositions and ages (Jackson and Sangster, 1987) variously modified by glacial erosion during the Quaternary and with patchy till cover; hilly and hummocky surfaces, ice moulded in places, with lake basins in subglacially scoured regions; smooth surfaces exhibiting little or no sign of glacial erosion in peninsular interiors (Dyke, 1983); cliffs resulting from glacial over-steepening; in places veneered by thin till, commonly bouldery.

- Geological boundary (defined, assumed)
- Areas covered by periglacial icefields during the Little Ice Age (indicated by a white pattern)
- Glacial lake spillway
- Glacial lake limit
- Marine limit
- Marine limit elevation in metres
- Weakly developed strandline
- Cliff in bedrock
- Lateral meltwater channel; bars on upslope side
- Subglacial and proglacial meltwater channel (large, small)
- Esker
- Kame
- Ice contact face
- Ribbed moraine
- Lateral moraine
- End moraine
- Margin of glacial dispersal train; teeth toward axis, steep side of teeth face down ice
- Lateral sliding boundary; teeth on sliding side
- Lateral sliding boundary; teeth on sliding side
- Iceberg scour
- Drumlinoid hill
- Crag-and-tail
- Ice moulded bedrock
- Striae (ice flow direction known, unknown)
- Crossed striae (numbers indicate relative age, 1 being the oldest)
- Field observation site: bouldery diamict (bd), bouldery gravel (bg), diamict (d), gravel (g), gravelly sand (gs), mud (m), muddy sand (ms), rock (r), sand (s), sandy gravel (sg), stony mud (sm), till (t)
- Field observation site: material as above near rock outcrop
- Radiocarbon date
- Date
- Material
- Lab no.
- Elevation (m)

REFERENCES

Dyke, A.S., 1993: Landscapes of cold-centred Late Wisconsinan ice caps, Canadian Arctic: Progress in Physical Geography v.17, p.223-247.

Jackson, G.D. and Sangster, D.F., 1987: Geology and resource potential of a proposed national park, Bylt Island and northwest Baffin Island, Northwest Territories, Geological Survey of Canada, Paper 87-17, 31 p.

OPEN FILE  
DOSSIER PUBLIC  
4695

Geological Survey of Canada  
Commission géologique du Canada  
2005

Recommended citation:  
Dyke, A.S., 2005: Surficial geology, Nina Bang Lake, Baffin Island, Nunavut: Geological Survey of Canada, Open File 4695, scale 1:50 000.



LOCATION MAP

Author: A.S. Dyke

Geology by A.S. Dyke, 2004

Field data provided by De Beers Canada Corporation, 2003

Digital cartography by M.M. Proulx, Earth Sciences Sector Information Division (ESS Info)

This map was produced from processes that conform to the ESS Info Publishing Services Subdivision Quality Management System, registered to the ISO 9001:2000 standard

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

OPEN FILE 4695  
SURFICIAL GEOLOGY  
NINA BANG LAKE  
BAFFIN ISLAND  
NUNAVUT

Scale 1:50 000/Échelle 1/50 000

kilometres 1 0 1 2 3 4 kilometres

Universal Transverse Mercator Projection  
North American Datum 1983  
© Her Majesty the Queen in Right of Canada 2005

Projection transverse universelle de Mercator  
Système de référence géodésique nord-américain, 1983  
© Sa Majesté la Reine du chef du Canada 2005

Digital base map from data compiled by Geomatics Canada, modified by ESS Info

Locational accuracy of the base appears to be ±100 m based on plotting of GPS measured field site locations

Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area  
Mean magnetic declination 2005, 43°30'W, decreasing 43.7° annually

Elevations in metres above mean sea level  
Contour interval 20 m

Field altimetry and the placement and trend of raised shorelines may conflict significantly with the contours

37 F103 OF4695	37 F104 OF4696	37 F105 OF4697	37 F106 OF4698
37 F107 OF4699	37 F108 OF4700	37 F109 OF4701	37 F110 OF4702
37 F111 OF4703	37 F112 OF4704	37 F113 OF4705	37 F114 OF4706
37 F115 OF4707	37 F116 OF4708	37 F117 OF4709	37 F118 OF4710
37 F119 OF4711	37 F120 OF4712	37 F121 OF4713	37 F122 OF4714
37 F123 OF4715	37 F124 OF4716	37 F125 OF4717	37 F126 OF4718
37 F127 OF4719	37 F128 OF4720	37 F129 OF4721	37 F130 OF4722
37 F131 OF4723	37 F132 OF4724	37 F133 OF4725	37 F134 OF4726
37 F135 OF4727	37 F136 OF4728	37 F137 OF4729	37 F138 OF4730
37 F139 OF4731	37 F140 OF4732	37 F141 OF4733	37 F142 OF4734
37 F143 OF4735	37 F144 OF4736	37 F145 OF4737	37 F146 OF4738
37 F147 OF4739	37 F148 OF4740	37 F149 OF4741	37 F150 OF4742

NATIONAL TOPONYMIC SYSTEM REFERENCE AND INDEX  
TO ALTERNATE GEOLOGICAL SURVEY OF CANADA MAPS