

# Buried valley aquifers: New data collection for municipal water supply and watershed management, Caledon East, Ontario




Figure 1. Location map of southern Ontario showing the location of the Caledon East study area (red dot). Geology modified from Barnett et al., 1991.

**Recommended citation:**  
 Russell, H.A.J., Pullan, S.E., Hunter, J.A., Sharpe, D.R., and Hohy, S. 2006. Buried valley aquifers: New data collection for municipal water supply and watershed management, Caledon East, Ontario. Geological Survey of Canada, Open File 5275.

**The map is available from:**  
 Bookstore, Geological Survey of Canada  
 611 Booth Street, Ottawa, Ontario K1S 5B6  
 Tel: 1-800-252-4371; Fax: (613) 953-6068  
 E-mail: gsc\_bookstore@geoscan.gc.ca  
 Web Site: <http://www5.gov.gc.ca/gsc/bookstore/>

**Web Publication:**  
 Products of the Southern Ontario Hydrogeology Project are also available on the project website: <http://www.southernhydrogeology.ca>

**Viewing Only:**  
 Ontario Geological Survey Publications  
 600 St. Mary Street  
 Toronto, Ontario, M7A 1C3

## Introduction

- Buried-valley aquifers are an important source of water supply in southern Ontario. They are significant to water resource managers as the need increases for source water protection, security of supply, and for constraining estimates of watershed-scale water balances.
- Prospecting methods for this aquifer type have seldom used modern exploration techniques to discover, target, and assess the reservoir potential and flow-system properties. This work outlines the exploration strategy employed near Caledon East in an ~141 km<sup>2</sup> area of the Credit River and Humber River watersheds, southern Ontario.

## Regional Geological Setting

- The lithostratigraphic framework of the study area (Karrow, 1967) has been reinterpreted using basin analysis principles and event stratigraphic concepts (Figs. 1, 2; Sharpe et al., 1996). A key result is the mapping of a regional unconformity that is defined by drumlinized Newmarket Till and tunnel channels (Barnett et al., 1996).
- A key result is the mapping of a regional unconformity that is defined by drumlinized Newmarket Till and tunnel channels (Barnett et al., 1996; Sharpe et al., 2004).
- To permit mapping of the regional stratigraphy using archival data, the lithostratigraphic framework has been simplified to five principal units. They are, stratigraphically upward: 1) Paleozoic bedrock, 2) Lower sediment, 3) Newmarket Till, 4) Oak Ridges Moraine and channel sediment, and 5) Halton Till. Lower sediment (LS) comprises 10 poorly exposed formations representing middle Wisconsinian and older sediment (Fig. 2) described mainly from Scarborough Bluffs (Karrow, 1967; Eyles et al., 1985; Sharpe et al., 1996).

## Geological Model




Figure 2. a) Lithostratigraphic framework and b) conceptual geological model of the ORM region.

## Study Area Geology





Figure 3. Geological context of study site. a) Digital elevation model (DEM) with a surficial geology drape of the area. Geology from Russell and White (1996). Seismic sections (e.g. S-1) and key borehole sites (CVC-1, IWA (Interim Waste Authority sites) are indicated.

## Data

- A continuous core borehole (CVC-1) was drilled to a depth of 180 m and logged with downhole geophysics (Fig. 4). Sampling was completed for grain-size, total organic carbon, and water content (see Russell et al., 2005).
- Reflection seismic data were collected along 3 profiles, at a spacing of 4-6 km, for a total of ~12 line-kms (Figs. 5).
- Seismic data were collected using a 120 channel array of 40 Hz geophones at 5 m spacing, and an 1/4 m (mivib) 17M vibratory seismic source (30-300 Hz). The output here is shown as a variable density display (see Pullan et al., 2004, for additional info).
- Geophysical logs obtained in the cased borehole include natural gamma, conductivity, magnetic susceptibility, spectral gamma, temperature and seismic velocity measurements (Fig. 6).

## Sediment Log



- Bedrock is intercepted at 178 m and is Georgian Bay Formation.
- The 178 m sediment log fines upward from basal gravel and sand to mud recording a decrease in flow energy in the fill succession and a change in depositional environments.
- Secondary, smaller fining-upward cycles (S-6) record temporal variations in depositional processes (Fig. 7) and control various scales of aquifer heterogeneity.
- The coarsest basal sediment is within the buried valley (below 100 m depth, Fig. 5) and has not been previously recorded in water well records from the area.
- Detailed sediment descriptions allow for improved hydrogeological correlation, characterization and groundwater modeling.

## Seismic Stratigraphy




Figure 5. Reflection seismic data displayed as south to north profiles. Location of respective lines is shown in figure 3. Seismic reflection patterns are correlated with sediment facies identified in borehole CVC-1 (Fig. 4). Five main seismic facies are identified. Horizontal scale of lines is in kilometres, vertical scale is in milliseconds (left side) and metres below ground surface (right side).

## Borehole Geophysics




Figure 6. Downhole geophysics for borehole CVC-1 showing geophysical logs and preliminary interpretation. The pronounced change in gamma and magnetic susceptibility data at 98 m depth corresponds to a distinct upward change in core character from sand to silt.

## Correlation of Seismic and Sediment Facies




Figure 7. Correlation of the four principal seismic facies with sediment facies and illustration of representative seismic signature and core photos. The fieldwork and project management were carried out under the direction of the Conservation Authorities Moraine Coalition. Steve Davies oversaw the drilling, core collection and initial field logging of the borehole. George Cornell provided further detailed logging of the BH in the laboratory. Seismic data were collected and processed by GAPS (Geophysical Applications Processing Services), Ouellet, Ontario. Funding for the drilling of CVC-1 was provided through the Ministry of the Environment's Municipal Groundwater Study Program. Funding for the seismic survey was provided through the Ministry of Northern Development and Mines via the Ontario Geological Survey. Graphic layout by Tracy Barry.

## Bedrock Valley




Figure 8. Interpolated bedrock surface from archival data (a) and including new data (b). Note interpolative artifacts between seismic lines.




Figure 9. Graph of interpolated valley cross-section at position of CVC-1. Valley geometry from (a) archival data, (b) addition of CVC-1, and (c) addition of seismic profile S-1.

## Hydrogeological Implications




Figure 10. Schematic cross-section extending from the Niagara Escarpment toward the southeast along a continuation of the Caledon East seismic line 2.

- New seismic reflection data and the CVC-1 borehole data has increased the known volume of the previously poorly defined buried valley at Caledon by ~40%.
- Locally the buried valley fill may consist of up to 100 m of interbedded sand and gravel with hydraulic conductivity values in the range of 10<sup>-2</sup> to 10<sup>-5</sup> m s<sup>-1</sup> (values from Freeze and Cherry, 1979).
- New data provides a framework for strategic hydraulic testing of this buried-valley aquifer.
- Buried valleys (bedrock or sediment hosted) are important aquifers regionally and currently supply water to municipalities at Aurora, Newmarket, Uxbridge, Nobleton, Inglewood, and Cheltenham.
- The Caledon East area illustrates the need for new, high-quality, subsurface data (i.e. seismic profiles and core boreholes) to discover and assess previously unknown aquifer potential.
- This buried valley east of the Forks of the Credit is one example of a possible relationship that may exist between re-entrants along the Niagara Escarpment and buried valleys to the east (e.g. Hookley, Sheldon, Limehouse).
- Geoscience expertise and methods available from geological surveys provides an important complement to the water resource management skills of conservation authorities, and municipal governments.

## Summary

- New borehole and seismic data collected within a buried valley has increased the known valley volume by 40%.
- Seismic facies analysis indicates that the valley fill may contain up to 100 m of sand and gravel.
- 60% of the valley fill (prior to data collection) was interpreted to be Oak Ridges Moraine (ORM) sediment. New data collection indicates a greater percentage of the fill is ORM sediment.
- New data indicates that the buried valley probably contains a coarse grained aquifer that has not been previously intercepted by water well records.
- The thickness and lateral continuity of the seismic facies interpreted as sand and gravel suggests an aquifer that could possibly provide municipal-scale water supply.

## References

Barnett, P.J., Cowan, W.R. and Henry, A.P. 1991. Quaternary Geology of Ontario, southern sheet. Ontario Geological Survey, map 2556.

Barnett, P.J., Sharpe, D.R., Russell, H.A.J., Brennan, T.A., Kenny, F.M., and Pugh, A., 1998. On the origin of the Oak Ridges Moraine. Canadian Journal of Earth Sciences, v. 35, p. 1152-1167.

Dames and Moore, Canada, 1992. Five phase groundwater study in the town of Caledon, Peel Region of Peel. Project 89-1600.

Dames and Moore, Canada, 1998. Groundwater quantification study Bolton, Caledon East, and Paisley; volume 1 and 2. Text, Figures, and Appendices. Peel Region of Peel, Project 1684-021.

Eyles, N., Clark, B.M., Kaye, B.G., Howard, K.W.F., and Eyles, C.H., 1965. The application of basin analysis techniques to glacialized terrain: an example from the Lake Ontario Basin, Canada. Geoscience Canada, v. 12, p. 22-32.

Freeze, R.A. and Cherry, J.A., 1979. Groundwater. Englewood Cliffs, Prentice-Hall.

Gornell and Associates, 2004. Internal report submitted to the Conservation Authorities Moraine Coalition.

Karrow, P.F., 1967. Pleistocene Geology of the Scarborough Area. Toronto, Ontario Ministry of Natural Resources, Report 46, 108 p.

Pugh, A., Pullan, S.E., and Sharpe, D.R., 1999. Seismic facies and regional architecture of the Oak Ridges Moraine area, southern Ontario. Canadian Journal of Earth Sciences, v. 36, p. 409-432.

Pullan, S.E., Hunter, J.A., and Good, R.L., 2002. Using Downhole Geophysical Logs to Provide Detailed Lithology and Stratigraphic Assignment, Oak Ridges Moraine, southern Ontario. Current Research 2002-08. Ottawa, Geological Survey of Canada, p. 1-12.

Pullan, S.E., Hunter, J.A., Russell, H.A.J., and Sharpe, D.R., 2004. Delineating buried-valley aquifers using shallow seismic reflection profiling and downhole geophysical logs: an example from southern Ontario, Canada. International Conference on Environment and Engineering Geophysics (ICEEG-2004), Wuhan, China, June 6-11, 2004.

Russell, H.A.J., and Stacey, P. (compilers) 2001. Bedrock Topography and Sediment Thickness DEMs of the Greater Toronto and Oak Ridges Moraine Areas, southern Ontario. Geological Survey of Canada, Open File 3699, 1 October, 2001.

Russell, H.A.J., Arnold, R.V.C., and Sharpe, D.R., 2003. Evidence for rapid sedimentation in a tunnel channel, Oak Ridges Moraine, southern Ontario, Canada. Sedimentary Geology, v. 161, p. 33-58.

Russell, H.A.J., Davis, S., Gornell, G., Hunter, J.A., and Sharpe, D.R., 2005. Sedimentology and Geophysical summary of the CVC-1 borehole and core at Caledon East, southern Ontario. Geological Survey of Canada, Open File (in preparation).

Russell, H.A.J., Pullan, S.E., Hunter, J.A., Sharpe, D.R., and Hohy, S., 2004. Buried-valley Aquifers: Delineation and Characterization from Seismic Reflections and Core Data at Caledon East, Ontario. In: Berg, H., Russell, H., and Thornton, H. (editors) Geological Models for Groundwater Flow Modeling, Workshop Extended Abstracts, May 15, 2004, GAC/MAC St. Catharines, Illinois State Geological Survey, Open File 2004-4.

Sharpe, D.R., Dyke, L.D., Hinton, M.J., Pullan, S.E., Russell, H.A.J., Brennan, T.A., Barnett, P.J., and Pugh, A., 1996. Groundwater prospects in the Oak Ridges Moraine area, southern Ontario: application of regional geological models. Current Research 1996-E. Ottawa, Geological Survey of Canada, p. 181-190.

Sharpe, D.R., Pugh, A., Pullan, S.E., and Gornell, G., 2002. Application of seismic stratigraphy and sedimentology to regional investigations: an example from Oak Ridges Moraine, southern Ontario, Canada. Canadian Geotechnical Journal, v. 40, p. 711-730.

Sharpe, D.R., Pugh, A., Pullan, S.E., and Shaw, J., 2004. Regional unconformities and the sedimentary architecture of the Oak Ridges Moraine area, southern Ontario. Canadian Journal of Earth Sciences, v. 41, p. 183-198.

## Acknowledgments

The fieldwork and project management were carried out under the direction of the Conservation Authorities Moraine Coalition. Steve Davies oversaw the drilling, core collection and initial field logging of the borehole. George Cornell provided further detailed logging of the BH in the laboratory. Seismic data were collected and processed by GAPS (Geophysical Applications Processing Services), Ouellet, Ontario. Funding for the drilling of CVC-1 was provided through the Ministry of the Environment's Municipal Groundwater Study Program. Funding for the seismic survey was provided through the Ministry of Northern Development and Mines via the Ontario Geological Survey. Graphic layout by Tracy Barry.

## Citation

Russell, H.A.J., Pullan, S.E., Hunter, J.A., Sharpe, D.R., and Hohy, S. 2006. Buried valley aquifers: New data collection for municipal water supply and watershed management, Caledon East, Ontario. Geological Survey of Canada, Open File 5275.

