

# Geological and Geochemical Data from the Canadian Arctic Islands.

## Part II: Rock Eval/TOC data

Mark Obermajer, K. Ross Stewart, and Keith Dewing

Geological Survey of Canada

### **Abstract**

A total of 4716 Rock Eval analyses are given for samples from 109 oil and gas wells (or deviated legs) and 1655 outcrop samples from the Arctic Islands. Each sample is assigned to a stratigraphic unit.

### **Introduction**

The Arctic Archipelago includes a land area of 780,000 km<sup>2</sup> covering much of the Northwest Territories and Nunavut. It contains three of the 10 largest islands in the world (Baffin, Victoria and Ellesmere) plus twenty-two others of appreciable size. There was an oil and gas exploration boom in the Arctic Islands between 1960 and 1985 that resulted in 191 exploration boreholes and the discovery of 20 major petroleum fields. Estimates of the hydrocarbon resource made by the Geological Survey of Canada predict 45 to 50 Tcf of in-place natural gas in the Sverdrup Basin, whereas the National Energy Board (NEB) predicts 45 Tcf of marketable gas in the whole Arctic Islands area.

The Geological Survey of Canada is undertaking an update of the geological and geochemical datasets for the Arctic Islands. This includes Rock Eval data (herein) and the depth to the top of each stratigraphic unit in the Arctic Islands hydrocarbon exploration boreholes (Dewing and Embry, 2007).

The stratigraphic nomenclature is presented in Figure 1. Formation descriptions are available from the Lexicon of Canadian Geological Names ([http://gdr.nrcan.gc.ca/index\\_e.php](http://gdr.nrcan.gc.ca/index_e.php)). These include age and lithological information, as well as the location of the type section and references. Excellent summaries of the geology of the Canadian Arctic are found in Trettin (1991).

## **Experimental**

Rock-Eval/TOC analysis provides fast and reliable characterization of the quantity and quality of sedimentary organic matter, as well as its thermal maturity. Pyrolysis experiments were conducted using Delsi Rock-Eval II and Rock-Eval VI units equipped with a Total Organic Carbon analysis module.

A typical Rock-Eval II experiment was initiated with heating of a pulverized rock sample at 300°C for 3 min in helium atmosphere, when naturally occurring hydrocarbons (free and adsorbed) are volatilized. During the next stage, the oven temperature is steadily increased to 600°C at a rate of 25°C/min and decomposition of kerogen occurs. The final stage involves oxidation and combustion of the residual organic matter at 600°C. The amount of hydrocarbons volatilized at 300°C and evolved from kerogen at 300°C to 600°C is quantitatively determined by a flame ionization detector, and recorded as the S1 and S2 peaks, respectively. The temperature measured at the maximum of the S2 peak is referred to as Tmax. The quantity of organic CO<sub>2</sub> generated from 300°C to 390°C, determined by a thermal conductivity detector, comprises the S3 peak. The percentage of carbon in CO<sub>2</sub> formed during oxidation at 600°C and in the hydrocarbon peaks S1 and S2 is used to define the total organic carbon content (TOC), expressed as a weight percentage. The determination of the quality of organic matter is based upon the calculation of Hydrogen (HI) and Oxygen (OI) indices (HI=S2/TOCx100, OI=S3/TOCx100) which are related to the atomic H/C and O/C ratios (Espitalie et al., 1977).

The OI versus HI cross plots ("pseudo van Krevelen diagrams") can be used as an organic matter type indicator at low and moderate maturities. The Tmax is an indicator of relative thermal maturity. According to Espitalie et al. (1985) the oil window is defined by the following Tmax ranges: 440°-448°C (Type I), 430°-455°C (Type II) and 430°-470°C (Type III). A cross plot of Tmax versus HI is used to constrain estimations of organic matter type and its thermal maturity, while the Production Index (PI=S1/[S1+S2]) is used to indicate staining of a sample or as an additional maturity parameter.

## **Guidelines for Interpretation**

Rock-Eval results correlate to other techniques (Espitalie et al., 1985; Tissot and Welte, 1978, Peters, 1986). Source rock potential is sensitive to lithology, TOC and S2 values. It is common practice to rate carbonate rocks with lower TOC comparable with richer clastic rocks. Extractable HC yields from leaner carbonate rocks are comparable to richer clastic rocks (Tissot and Welte, 1978, p. 430; Gehman, 1962). The organic matter associated with carbonate rocks is often more hydrogen-rich and thermally labile than that in fine-grained clastic rocks. As a result, more TOC in carbonate rocks may be transformed into bitumen compared with average clastic source rocks of comparable maturity.

Rock-Eval/TOC parameters have significance only above threshold TOC, S1 and S2 values. If TOC is less than less than or equal to 0.3% then all parameters have questionable significance and the experiment suggests no potential. Oxygen Index (OI), S3/TOC, has questionable significance if TOC is less than or equal to 0.5%. Both Tmax and Production Index ( $PI = S1/(S1+S2)$ ), have questionable significance if S1 and S2 values are less than or equal to about 0.35 mg HC/g rock. Results can be affected by mineral matrix effects. These either retain generated compounds, generally lowering the S1 or S2 peaks, while increasing Tmax, or by liberating inorganic CO<sub>2</sub> and increasing S3 and OI. These effects are important if TOC, S1 and S2 are low, an effect not significant where sources have TOC values greater than 5%. OI values greater than 150 mg/g TOC suggest either low TOC or a mineral matrix CO<sub>2</sub> contribution during pyrolysis. Note that TOC and Hydrogen Index decrease with increasing thermal maturity due to hydrocarbon generation.

## **Arctic Islands borehole dataset**

This report contains files (\*.xls format) with Rock Eval data for samples from the Arctic Island boreholes and outcrop. These data are archived at GSC-Calgary and include data collected by Panarctic Oils Ltd and stored in the library collections at GSC-Calgary (Leythaeuser and Stewart, 1986), data collected by the GSC as part of an economic assessment of Bathurst Island (Stewart, 1999), Thomas Gentzis' PhD thesis (Gentzis, 1991), as well as samples run by the GSC as part of published (e.g., Brooks et al., 1992; Fowler, 1991; Goodarzi et al., 1987; Stasiuk and Fowler, 1994) and unpublished regional or stratigraphic studies. About 700 samples were run in 2005-06 to augment the previous datasets. An additional 1060 samples of the Kanguk Formation, previously released as GSC Open File 2727 (Nunez-Betelu, 1993), are included.

Figure 2 shows the TOC distribution and an HI vs OI plot (pseudo-van Krevelen diagram) for each of the main sedimentary successions.

The \*.xls files include the following data fields:

1. SAMP\_ID – either the UWI or a GSC-Calgary curation C-number
2. LOC\_ID – Location ID/Unique Well Identifier
3. LOC\_NAME – Section name/Well name
4. SAMP\_LAT – Surface Latitude
5. SAMP\_LONG – Surface Longitude (negative is west)
6. DEPTH\_FROM – Depth (m) of sample top from Kelly Bushing. May sometimes be sample midpoint.
7. DEPTH\_TO – Depth (m) of sample base
8. OFROM – Depth of sample top in units submitted to lab
9. OTO – Depth of sample base in units submitted to lab
10. DEPTH\_UNIT – Feet or Metres
11. EQUIP\_TYPE – Rockeval II or Rockeval 6

### **Rockeval Parameters:**

12. TOC - Total Organic Carbon (weight percent)
13. TMAX - maximum temperature at S2 peak (°C)
14. S1 - hydrocarbons volatilized at 300°C (mg HC/g rock)
15. S2 - hydrocarbons evolved from kerogen at 300°C to 600°C (mg HC/g rock)
16. S3 - organic CO<sub>2</sub> generated from 300°C to 390°C (mg/g rock)
17. RC – residual carbon
18. PC – pyrolysable carbon
19. MINC – mineral carbon
20. HI - Hydrogen Index (mg hydrocarbons/g organic carbon)
21. OICO - Oxygen Index CO (mg CO/g organic carbon)
22. OI - Oxygen Index CO<sub>2</sub> (mg CO<sub>2</sub>/g organic carbon)
23. TPKS2 – maximum temperature of surface S2
24. S3CO – CO from organic and mineral sources
25. PI - Production Index (S1/[S1+S2])
26. QUANTITY – Quantity of material analysed
27. REFERENCE – Source of data (if other than GSC lab or if previously published).
28. UNIT – Stratigraphic Unit, based on GSC Open File 5542

## References

- Brooks, P.W., Embry, A.F., Goodarzi, F. and Stewart, R.
- 1992: Organic geochemistry and biological marker geochemistry of Schei Point Group (Triassic) and recovered oils from the Sverdrup Basin (Arctic Islands, Canada). *Bulletin of Canadian Petroleum Geology*, v. 40, p. 173-187.
- Dewing, K. and Embry, A.F.
- 2007: Geological and Geochemical Data from the Canadian Arctic Islands. Part I: Stratigraphic Tops from Arctic Islands Oil and Gas Exploration Boreholes. *Geological Survey of Canada Open File 5442*. 1 CD.
- Espitalie, J., Laporte, J.L., Madec, M., Marquis, F., Leplat, P., Paulet, A. and Bouteleau, J.
- 1977: Methode rapide de characterisaion des roches meres de leur potential petrolier et de leur degre d'evolution. *Revue de l'Institut Francais du Petrole* v. 32, p. 23-42.
- Espitalie, J., Deroo, G. and Marquis, F.
- 1985: Rock Eval Pyrolysis and Its Applications. Preprint; Institut Francais du Petrole, Geologie No. 27299, 72 p. English translation of, *La pyrolyse Rock-Eval et ses applications*, Premiere, Deuxieme et Troisieme Parties, in *Revue de l'Institut Francais du Petrole*, v. 40, p. 563-579 and 755-784; v. 41, p. 73-89.
- Fowler, M.G., Goodarzi, F., Gentzis, T., and Brooks, P.W.
- 1991: Hydrocarbon potential of Middle and Upper Devonian coals from Melville Island, Arctic Canada. *Organic Geochemistry*, v. 17, p. 681-694.
- Gehman, H. M.
- 1962: Organic matter in limestones; *Geochimica et Cosmochimica Acta*, v. 26, p. 885-897.
- Gentzis, T.
- 1991: Regional maturity and source-rock potential of Paleozoic and Mesozoic strata, Melville Island, Arctic Canada. Unpublished PhD thesis, The University of Newcastle-upon-Tyne.
- Goodarzi, F., Nassichuk, W.W., Snowdon, L.R., and Davies, G.R.
- 1987: Organic petrology and Rockeval analysis of the Lower Carboniferous Emma Fiord Formation in the Sverdrup Basin, Arctic Archipelago. *Marine and Petroleum Geology*, v. 4, p. 132-145.

Leythaeuser, D. and Stewart, K.R.

1986: Generation, migration and accumulation of hydrocarbons in the Sverdrup Basin, Canada. A geochemical basin study. Volume II – tables and figures. Unpublished report to Panarctic Oils. GSC Catalogue No. QE196.9 L4 1986 v.2.

Nunez-Betelu, L.K.

1993: Rock- eval/ TOC pyrolysis data from the Kanguk Formation (Upper Cretaceous), Axel Heiberg and Ellesmere Islands, Canadian Arctic. Geological Survey of Canada, Open File 2727, 29 p.

Peters, K. E.

1986: Guidelines for evaluating source rock using programmed pyrolysis; Bulletin of American Association of Petroleum Geologists, v. 70, p. 318-329.

Snowdon, L. R.

1995: Rock-Eval Tmax suppression: documentation and amelioration. Bulletin of American Association of Petroleum Geologists, v. 79, p. 1337-1348.

Stasiuk, L.D. and Fowler, M.G.

1994: Paleozoic hydrocarbon source rocks, Truro Island, Arctic Canada: organic petrology, organic geochemistry and thermal maturity. Bulletin of Canadian Petroleum Geology, v. 42, p. 419-431.

Stewart, K.R.

1999: Appendix One: Rock-eval analyses. In: C.D. Anglin and J.C. Harrison (eds.) Mineral and energy resource assessment of Bathurst Island area, Nunavut. Geological Survey of Canada, Open File 3714, p. F55-F88.

Tissot, B. P., and Welte, D. H.

1978: Petroleum formation and occurrence; Springer-Verlag, Berlin, 538 p.

Trettin, H.P. (ed.)

1991: Geology of the Innuitian Orogen and Arctic Platform of Canada and Greenland: Geological Survey of Canada, Geology of Canada no. 3 (also Geological Society of America, Geology of North America v. E), 569 p.

### **Selected References for Figure 1**

- Beauchamp, B., Harrison, J.C., Utting, J., Brent, T.A., and Pinard, S.  
2001: Carboniferous and Permian subsurface stratigraphy, Prince Patrick Island, Northwest Territories, Canadian Arctic; Geological Survey of Canada, Bulletin 565, 93 p.
- de Freitas, T., Harrison, J.C., and Mayr, U.  
1997: Sequence stratigraphic correlation charts of the lower Paleozoic Franklinian succession, Canadian Arctic and parts of North Greenland. Geological Survey of Canada Open File 3410, 3 charts.
- Dewing, K. and Nowlan, G.S.  
2004: Correlation chart of Cambrian and Ordovician stratigraphy, Arctic Islands, Nunavut, Canada; Geological Survey of Canada, Open File 1837, 1 chart.
- Embry, A.F.  
1983a: Stratigraphic subdivision of the Heiberg Formation, eastern and central Sverdrup Basin, Arctic Islands. Geological Survey of Canada, Paper 83-1B, p. 205-213.  
1983b: The Heiberg Group, western Sverdrup Basin, Arctic Islands. Geological Survey of Canada, Paper 83-1B, p. 381-389.  
1984a: Stratigraphic subdivision of the Roche Point, Hoyle Bay and Barrow formations (Schei Point Group), western Sverdrup Basin, Arctic Islands. Geological Survey of Canada, Paper 84-1B, p. 275-283.  
1984b: The Wilkie Point Group (Lower-Upper Jurassic), Sverdrup Basin, Arctic Islands. Geological Survey of Canada, Paper 84-1B, p. 299-308.  
1984c: The Schei Point and Blaa Mountain Groups (Middle-Upper Triassic), Sverdrup Basin, Canadian Arctic Archipelago. Geological Survey of Canada, Paper 84-1B, p. 327-336.  
1985a: Stratigraphic subdivision of the Isachsen and Christopher formations (Lower Cretaceous), Arctic Islands. Geological Survey of Canada, Paper 85-1B, p. 239-246.  
1985b: New stratigraphic units, Middle Jurassic to lowermost Cretaceous succession, Arctic Islands. Geological Survey of Canada, Paper 85-1B, p. 269-276.  
1986a: Stratigraphic subdivision of the Blind Fiord and Bjorne formations (Lower Triassic), Sverdrup Basin, Arctic Canada. Geological Survey of Canada, Paper 86-1B, p. 329-340.

1986b: Stratigraphic subdivision of the Awingak Formation (Upper Jurassic) and revision of the Hicles Cove Formation (Middle Jurassic), Sverdrup Basin, Arctic Canada. Geological Survey of Canada, Paper 86-1B, p. 341-349.

1994: Uppermost Triassic, Jurassic, and lowermost Cretaceous stratigraphy, Melville Island area, Arctic Canada. In: R.L. Christie and N.J. McMillan (eds.) The Geology of Melville Island, Arctic Canada. Geological Survey of Canada, Bulletin 450, p. 139-159.

Embry, A.F. and Klovan, J.E.

1976: The Middle-Upper Devonian clastic wedge of the Franklinian geosynclinal; Bulletin of the Canadian Society of Petroleum Geologists, v. 24, p. 485-639.

Harrison, J.C.

1995: Melville Island's salt-based fold belt, Arctic Canada; Geological Survey of Canada, Bulletin 472, 331 p.

Harrison, J.C. and Brent, T.A.

2005: Basins and fold belts of Prince Patrick Island and adjacent areas, Canadian Arctic Islands; Geological Survey of Canada, Bulletin 560, 197 p.

Mayr, U.

1978: Stratigraphy and correlation of lower Paleozoic formations, subsurface of Cornwallis, Devon, Somerset, and Russell islands, Canadian Arctic Archipelago. Geological Survey of Canada, Bulletin 276, 55 p.

1980: Stratigraphy and correlation of lower Paleozoic formations, subsurface of Bathurst Island and adjacent smaller islands, Canadian Arctic Archipelago. Geological Survey of Canada, Bulletin 306, 52 p.

Mayr, U., Brent, T.A., de Freitas, T., Frisch, T., Nowlan, G.S., and Okulitch, A.V.

2004: Geology of eastern Prince of Wales Island and adjacent smaller islands, Nunavut; Geological Survey of Canada, Bulletin 574, 88 p.

Thorsteinsson, R. and Mayr, U.


1987: The sedimentary rocks of Devon Island, Canadian Arctic Archipelago; Geological Survey of Canada, Memoir 411, 182 p.

Uyeno T.T

1990: Biostratigraphy and conodont faunas of Upper Ordovician though Middle Devonian rocks, eastern Arctic Archipelago. Geological Survey of Canada, Bulletin 401, 210 p.




Natural Resources  
Canada Ressources naturelles  
Canada



*Late Cretaceous Kanguk succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Early-Late Cretaceous Hassel succession*



Natural Resources  
Canada Ressources naturelles  
Canada




*Early Cretaceous Christopher succession*



Natural Resources  
Canada


Ressources naturelles  
Canada



*Early Cretaceous Isachsen succession*




Natural Resources  
Canada Ressources naturelles  
Canada



Mid-Late Jurassic McConnell Island-Rignes-Deer Bay succession




Natural Resources  
Canada Ressources naturelles  
Canada



*Early-Mid Jurassic Jameson Bay succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Late Triassic-Early Jurassic Barrow-Heiberg succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Mid to Late Triassic Schei Point Group*




Natural Resources  
Canada Ressources naturelles  
Canada



*Early Triassic Blind Fiord-Bjorne succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Carboniferous-Permian clastic succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Carboniferous-Permian basinal carbonate succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Carboniferous lacustrine succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Devonian clastic wedge*




Natural Resources  
Canada Ressources naturelles  
Canada



*Devonian carbonate succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Devonian Boothia-related succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Silurian basinal succession*




Natural Resources  
Canada Ressources naturelles  
Canada



*Silurian carbonate succession*



Natural Resources  
Canada Ressources naturelles  
Canada



*Cambrian-Ordovician carbonate succession*