

LEGEND

This legend is common to GSC Open File maps produced for NTS sheets
84 L, 84 M, 94-I, and 94 P. Not all map units in the
common legend appear on this map

Note: In areas where the surficial cover forms a complex pattern, the area is coloured according to the dominant unit and labelled in descending order of cover (e.g., O-Tr). Where buried aggregate deposits (sand and gravel - commonly associated with Gt or Gd surficial units) are known, or suspected, areas are coloured according to the overlying unit and labelled in the following manner: Lv/Gd.

QUATERNARY
SURFICIAL DEPOSITS
POST LAST GLACIATION

NONGLACIAL ENVIRONMENTS

AN	ANTHROPOGENIC DEPOSITS: culturally-made or modified geological materials such that their physical properties (e.g., structure, cohesion, compaction) have been drastically altered; >2 m thick.
----	--

ORGANIC DEPOSITS: peat and muck; 1 to 3 m thick on average; formed by the accumulation of plant material in various stages of decomposition; generally occurs as flat, wet terrain (swamps and bogs) over poorly drained substrates.

O ¹	Bog peat: sphagnum or forest peat formed in an ombrotrophic environment; wet terrain; may be treed or treeless; O ¹ h, hummocky, mounds and plateaus; area may be underlain by ground ice or shallow permafrost conditions; O ¹ k, thermokarst terrain related to melting ground ice.
O ²	Fen peat: peat derived from sedges and partially decayed shrubs in a eutrophic environment; forms relatively open peatlands with a mineral-rich water table that persists seasonally near the surface; generally covered with low shrubs and an occasional sparse layer of trees.
O	Undifferentiated bog and fen deposits: Oh, undifferentiated hummocky bog and fen deposits; area may be underlain by ground ice or shallow permafrost conditions; Ok, undifferentiated bog and fen deposits with thermokarst terrain related to melting of ground ice; Oc, undifferentiated bog and fen deposits cut by numerous subparallel channels on gentle slopes.

COLLUVIAL DEPOSITS: mass wasting debris; poorly sorted, massive to stratified debris deposited by direct, gravity-induced movement; composition dependant on source material.

Ch	Landslide and slump debris: active and inactive landslides; hummocky topography; diamicton, generally 1 to 10 m thick, but may exceed 10 m near the toe of large landslides.
Cv	Colluvial veneer: thin and discontinuous cover of slumped and/or soliflucted material <1 m thick; overlies bedrock or till.
C	Undifferentiated colluvial deposits.

ALLUVIAL DEPOSITS: sorted gravel, sand, minor silt and organic detritus deposited by streams; commonly stratified.

Ap	Floodplain deposits: sorted gravel, sand, silt and organic detritus >1 m thick; forming active floodplains close to river level with meander channels and scroll marks.
At	Fluvial terrace deposits: inactive terraces above modern floodplain; >2 m thick; represents a potential aggregate source.
Ad	Deltaic sediments: stratified sand and gravel underlain by silt and clay; generally 2 to 15 m thick; occuring at the mouths of streams entering lakes.
Af	Alluvial fan deposits: poorly sorted gravel, sand and organic detritus >1 m thick.
Av	Alluvium veneer: < 1 m thick; primarily as uniform sheets of slope wash on gentle slopes.
A	Undifferentiated fluvial deposits.
L ¹	LACUSTRINE DEPOSITS: sand, silt and minor clay deposited in a former lake; >1 m thick; generally overlain by organic deposits; exposed by recent fluctuations in lake levels.

POSTGLACIAL OR LATE WISCONSINAN

PROGLACIAL AND GLACIAL ENVIRONMENTS

GLACIOLACUSTRINE DEPOSITS: fine sand, silt, and clay, with minor debris-flow diamicton, deposited in glacier-dammed lakes in valleys and along the margin of the retreating Laurentide Ice Sheet; usually overlain by organic deposits in lowlands.

Lb	Glaciolacustrine blanket: >1 m thick.
Lv	Glaciolacustrine veneer: thin and discontinuous; <1 m thick.

GLACIOFLUVIAL DEPOSITS: well to poorly stratified sand and gravel; minor diamicton; deposited behind, at or in front of the ice margin by glacial meltwater; represents a potential aggregate source.

G	Proglacial outwash: cross-stratified gravel and sand deposited in front of the ice margin; Gp, outwash plain deposits, generally 1 to 5 m thick, generally mantle valley floors and surfaces adjacent to glacial meltwater channel margins; Gt, outwash terrace deposits, often associated with meltwater channels and canyons; 1 to 10 m thick; Gd, glaciofluvial delta deposits; 1 to >30 m thick.
Gi	Ice-contact stratified drift: poorly-sorted sand and gravel with minor diamictons; deposited in contact with the retreating glacier; 1 to >20 m thick; Gi ^h , hummocky topography relating to melting of underlying ice; Gi ^k , surface marked by kettle holes; Gi ^r , esker ridges; Gi ^t , kame terraces; Gi ^d , ice-contact glaciofluvial delta deposits; 1 to >30 m thick, surface marked by kettles.

TILL: diamicton deposited directly by the Laurentide Ice Sheet; sandy to clayey matrix with striated clasts of various lithologies, including many Canadian Shield, carbonate and sandstone erratics; clast content is typically low (<10 %).

Tb	Till blanket: >1 m thick, continous till cover forming undulating topography that locally obscures underlying units.
Ts	Streamlined and fluted till: >1 m thick, till surface marked by streamlined landforms including flutes and drumlins.
Th	Hummocky till: >1m thick; hummocky till surface.
Tr	Ridged till deposits: >1 m thick, moraines or crevasse fillings forming a ridged topography.
Tv	Till veneer: <1 m thick, discontinuous till cover, underlying bedrock topography is discernible.

PRE-QUATERNARY
BEDROCK

R	Sedimentary bedrock, Cretaceous Fort St. John Group shales (including the Shaftesbury Formation) and Dunvegan Formation sandstone exposed in highlands and along meltwater channel and canyon walls.
---	--

Geological boundary (defined)	
Meltwater channel or underfit channel, small (paleoflow direction known, unknown)	
Major moraine	
Minor moraine and crevasse filling	
Fluting parallel to ice flow (direction unknown)	
Drumlin parallel to ice flow (flow direction unknown)	
Iceberg scour	
Thermokarst	
Till clast fabric	
Spring	
Field observation site (with, without sample)	