Natural Resources Canada | Ressources naturelles Canada

GEOLOGICAL SURVEY OF CANADA | COMMISSION GÉOLOGIQUE DU CANADA

DESCRIPTIVE NOTES

INTRODUCTION

Rocks underlying Smith Island and the adjoining peninsula on the Quebec mainland form the western end of the Cape Smith Belt segment of the Circum-Superior Belt of Paleoproterozoic (Aphebian) age (Baragar and Scoates, 1981). Relationships within the Cape Smith Belt have been well documented by the mapping of the Québec Ministère de l'Énergie et des Ressources (Hynes and Francis, 1982; Hynes and Rochette, 1990) and by the detailed stratigraphic and petrochemical studies of Hynes and Francis (1982), Francis et al. (1983), Baragar and Scoates (1987), and Picard et al. (1990). Much of this work is now embodied in the recent compilation by St-Onge et al. (2006). Previous work within the present map area include the author's preliminary maps (Baragar and Pichè, 1982; Baragar, 1983) and a map of the peninsula on the mainland by Togola (1992). Only a part of the Cape Smith Belt is represented here, but its context is readily interpretable by reference to well studied counterparts to the east as noted above.

The stratigraphy of the Cape Smith Belt, in general, comprises basal Povungnituk and overlying Chukotat groups, together deformed by southward folding and thrusting. The Povungnituk Group rests unconformably upon Archean basement rocks and is dominantly composed of mafic to ultramafic volcanic rocks with subordinate sedimentary rocks and carbonates. Upward in the section and outward from the craton (i.e. toward the north) the sediments change to deeper water facies (shale, semipelite, greywacke) with abundant, subaqueous, tholeiitic basalt units extensively intruded by diabase sills. The Chukotat Group is predominantly volcanic and ranges in composition, broadly comparable, from komatiite to essentially tholeiitic basalt, all parts of a consanguineous suite. The relationship between the groups is unclear: St-Onge et al. (1992) (and this study) interpreted the Chukotat as a thrust fault, whereas Hynes and Francis (1982) regarded it as a conformable boundary. Certainly, the Chukotat Group is younger, and if in thrust relationship with the Povungnituk Group, is unlikely to have undergone major displacement as sills of Chukotat Group affinity are present at several levels within the Povungnituk Group.

The Cape Smith Belt is cut by a suite of gabbroic sills (Parent). The Povungnituk Group underlies Te Gobin and marginal parts of the adjoining mainland and south coast of Smith Island. The Chukotat Group forms the bulk of Smith Island and extends along strike into the eastward-continuing sequence of the mainland. Only the upper 2000 m of the Povungnituk Group is exposed and shale, representing its deeper water facies, is present only at the Chukotat Group is limited to its lower 7000 m of komatiitic compositions (cf. Baragar and Scoates, 1987; Picard et al., 1990). A composite straight cross-section of both Povungnituk and Chukotat groups is shown in Figure 1. The Povungnituk Group section strikes northward through Te Gobin to the mainland and the Chukotat Group section transects Smith Island at about the midpoint of its east-west axis.

POVUNGNITUK GROUP

GENERAL GEOLOGY

The Povungnituk Group in this area is formed predominantly of pillowed and massive lava flows with rare, thin (centimetres-thick), black shale interlayers, commonly containing pyrrhotite. Overlying the main body of volcanic rocks is a thicker zone of thin-bedded black to grey shale and siltstone intruded by several dolerite and gabbro sills. On Te Gobin the pair of sills that give form to the northeast-plunging folds at its eastern end are seemingly felsiphatic, pyroxene-hornblende gabbro that are interpreted as Povungnituk Group sills on the basis of lithological similarity to Povungnituk Group dolerite observed farther east (Baragar and Scoates, 1987); however, no chemical data is available and it is entirely possible that they could be of Chukotat Group origin. The much thicker, folded sills intruding Povungnituk Group sedimentary rocks at a higher stratigraphic level on the mainland have a different aspect and, as will be shown later, do have a Chukotat Group chemistry.

VOLCANIC PETROGRAPHY AND COMPOSITION

The volcanic rocks are megascopically aphyric, but in thin section most can be seen to be sparsely plagioclase-phyric. The groundmass is a fine-grained intergrowth of plagioclase, clinopyroxene and feldspar [probably albite-oligoclase] with interstitial chlorite and a fuzzy brownish overlay of granular sphene. Epidote is generally present, but not abundant. The metamorphic grade is greenschist facies. Most rocks are somewhat foliated in an east-northeast direction with moderate to steep northward dips roughly parallel with the fold axis.

Of the 15 samples for which we have complete analyses from across Te Gobin, is that of a saturated, tholeiitic basalt with notable titania (3%) content. Its average composition is given in Table 1a, where it can be compared with averages of Povungnituk Group basalt from the Kovik section (about 70 km to the east (Baragar and Scoates, 1987)) as well as those of the Chukotat Group suite. The average form indicates that it is a fairly feldspathic basalt (~ 65% feldspar) with subordinate pyroxene (~ 36%). Note that the average normative composition shows both quartz and olivine, indicating that the composition was on either side of the silica-saturated condition. The rare-earth elements give a light-REE-enriched pattern (Baragar and Scoates, 1987) in conformity with those obtained from Povungnituk Group rocks throughout the Cape Smith Belt (Francis et al., 1983; Picard et al., 1990), to an extended element plot normalized to primitive mantle (Fig. 2b) the curve for Povungnituk Group basalt of Te Gobin shows enrichment in the more incompatible of the rare-earth elements and in Th relative to Nb. Basalt of the Povungnituk Group basalt units appear to have undergone significant fractionation prior to eruption. The setting of eruption has been interpreted as anelsio in a continental rift zone (Baragar and Scoates, 1981, 1987; Francis et al., 1983).

RELATIONSHIP TO CHUKOTAT GROUP

Because the contact between the Povungnituk and Chukotat groups transects the fairly continuous, low-Mg unit at the base of the latter group (see discussion) on the mainland, the nature of the basal sequence northeast of Te Gobin is interpreted as follows. Within the region of the fault is found at the upper margin of the folded gabbroic sill of Chukotat Group affinity and is folded with the Povungnituk Group rocks. Above the gabbro is a thin sequence of the folded shale and thin-bedded tuff units of the Povungnituk Group succeeded by a section of another projected thrust fault interpreted to follow the Illukotat River valley. At the south margin of Smith Island the faulted boundary is assumed to lie within, or at the top of, a thin sedimentary layer that overlies the sill.

CHUKOTAT GROUP

SUBDIVISION AND STRATIGRAPHY

The Chukotat Group comprises a homoclinal, steeply north dipping succession of komatiitic basalt and minor shaly interbeds with an exposed thickness of about 7000 m (Fig. 1). It is subdivided into a number of fairly distinctive, mappable units based mostly on MgO content that relates directly to the fluidity of the lavas and, in consequence, their morphology in the field. The lowest unit, the Ivakkak Member (Hynes and Francis, 1982) or Ivakkak Group (Baragar and Pichè, 1982), is a thick, basal mappable unit of high-Mg basalt (>15% MgO), medium-Mg (10–15% MgO), and high-Mg (~15% MgO) volcanic rocks. To accommodate mixed sequences, where no one type is dominant, additional categories provide for low- to medium-Mg and high- to medium-Mg lava mixes. Other units are distinguished on physical distinctions: onion-skin lavas are low-Mg flows characterized by large pillows with thick (3–10 cm), scaly rims; and thick massive units are differentiated flows as much as 250–300 m thick and kilometres in length.

Layered flows are also shown separately on the map. These are thickened lengths of massive flows or of the upper massive parts of pillow-massive flows in which extensive differentiation has resulted in a basal olivine cumulate layer surmounted by a feldspathic top (Baragar, 1984). Rarely, the upper part is marked by spinifex texture. Most are not more than 100 m thick and kilometres in length, but some are considerably thicker and extend for many kilometres. They are generally a maximum of 30 m to 100 m thick. On the map only a few are shown in their entirety, most are represented by arbitrary short lengths where they have been crossed in traverse. A majority of the layered flows are found in the high-Mg unit, some occur in the medium-Mg unit, but rarely are they present in the low-Mg unit.

The stratigraphy of the Chukotat Group is far from systematic units of differing MgO content thicken and thin as they interfinger along strike and seemingly pass arbitrarily prediction from one Mg level to another. Their lateral continuity may not persist for great distances. Note, for example, the passage of the medium-Mg unit at the core of Smith Island, eastward into a mixed unit of high- and medium-Mg lavas on the mainland. Broadly speaking, however, the incidence of high-Mg lavas tends to decrease upsection such that the northern part of the succession is dominated by medium-Mg units. This is also true of the Chukotat Group as a whole when compared to the Povungnituk Group (Smith Bell (Francis et al., 1983); Baragar and Scoates, 1987; Picard et al., 1990) where compositions evolve upward by repeating cycles of high- to medium- and/or low-Mg lavas (olivine- to clinopyroxene-phyric), each beginning with a lower Mg level than its predecessor until finally capped, in places, by a monotonous sequence of plagioclase-phyric basalt.

In this map area, similar repeating successions are interpretable from the section shown in Figure 1. Three levels of high-MgO lavas are each overlain by medium- and medium and/or low-Mg lava flows. Note also that the basal Chukotat Group units, both in the central and western parts of Cape Smith Belt (Hynes and Francis, 1982; Baragar and Scoates, 1987; Picard et al., 1990) is a low-Mg lava (clinopyroxene-phyric), suggestive of a prenatal stage of fractionation, possibly in crustal magma chambers, prior to the onset of primitive Chukotat Group magmatism from depth (Picard et al., 1990).

NORMAL HIGH- TO LOW-MG UNITS

Petrography

Metamorphism is pervasive throughout the Chukotat Group, but is of low grade (subgreenschist facies) and deformation is slight so that in most places the primary mineralogy is at least partly intact and readily interpretable. In low-Mg and rarely medium-Mg lava flows, where plagioclase is a significant component of the groundmass, pumpellyite is a common secondary constituent. Olivine is generally pseudomorphed by chlorite and/or serpentine and, less commonly, tremolite, but clinopyroxene is mostly fresh. Magnetite is extensively replaced by sphene. Primary igneous textures as described below tend to be well preserved.

The high-Mg lavas are abundantly olivine-phyric with a phenocryst content ranging from 20% to 50% and averaging about 30–35%. Although the higher concentrations are probably cumulative, many of the rocks are pillow lavas where evidence of concentration is not readily apparent. Surprisingly, the content of clinopyroxene phenocrysts tends to be quite high. Although about one quarter of the rocks examined are devoid of clinopyroxene phenocrysts, the majority contain from 5% to 15%. The olivine phenocrysts are typically euhedral and are commonly skeletal or even radiating bundles of acicular clinopyroxene and, generally, the olivine is a chromium-rich serpentinous mush. In thin section, plagioclase, interleaved with the clinopyroxene, can be recognized, but is greatly subordinate. Chromite is ubiquitous in both groundmass and olivine pseudomorphs.

Medium-Mg lavas, although highly variable, are generally much poorer in olivine phenocrysts (most 10–25%) and richer in clinopyroxene (most 5–35%) than in high-Mg lavas. In most cases, the groundmass is similarly quenched with finely intersewed clinopyroxene and plagioclase, forming fronds and radiating bundles. In the coarser grained phases, where the proportions are more evident, clinopyroxene dominates over plagioclase. Unlike the low-Mg lavas the groundmass may also contain skeletal olivine, indicative of its greater range of precipitation in this unit, and sparse quartz.

Typical low-Mg lavas are abundantly clinopyroxene-phyric (20–40%), but only scarcely olivine-phyric (0–10%). The groundmass textures vary from quench-like, interleaved plagioclase-clinopyroxene fronds overprinted by finely dispersed sphene to intergranular texture of plagioclase laths and clinopyroxene anhedra. In coarser grained rocks the plagioclase laths generally form a dominating framework enclosing clinopyroxene and rarely interstitial quartz.

Geochemistry

Compositions of low to high-Mg units can be broadly compared in Table 1a and 1b where their average compositions are presented together with those of some of the other units. Of particular interest are the progressive changes from low to high-Mg units of key oxides and elements: notably MgO, CaO, Al_2O_3, TiO_2 and, dramatically, Cr and Ni. The last three constituents have the additional significance that their range of variation is so markedly different from that of the Povungnituk Group tholeiite that they serve both as diagnostic elements of the komatiitic suite and as a mark of their consanguinity.

Compositional evolution in the stratigraphic sequence is represented in Figure 1 by the indices MgO/CaO and MgO/TiO_2 plotted above their locations within the cross-section. MgO/CaO is primarily a measure of olivine fractionation and MgO/TiO_2 of an extended range of fractionation that would include less refractory phases such as clinopyroxene and plagioclase (Baragar and Scoates, 1987). The MgO/TiO_2 is particularly applicable to rocks of the komatiitic suite that are enriched in Mg, high magnesia and low titania. Note in Figure 1 the following features.

- The Ivakkak Group komatiitic suite is distinguished from the Povungnituk Group tholeiitic rocks by their markedly higher MgO/TiO_2 even though the low-Mg komatiitic basaltic rocks are very similar to the tholeiite in many other respects.
- The komatiitic succession begins with a major low-Mg unit (as noted above), followed by an abrupt transition to high-Mg komatiitic basalt.
- Upsequence compositions (as represented by the fractionation indices) vary widely from unit to unit, but the indices do show slight overall decline, suggesting some progressive evolution of composition throughout the succession.

Multi-element plots normalized to primitive mantle are given in Figure 2a and 2b. To simplify the presentation only the averages are plotted, but they are generally representative of the multiple curves for each of the units. Note that the low-Mg unit (Fig. 2a) has a form that is very similar to that of the Povungnituk Group rocks in the same diagram, but fractionated curves. That, for instance, the curve for titanium (Ti) is by more than an order of magnitude lower than for low- to high-Mg lavas are successively offset to lower values, but for Cr to Ni (olivine compatible), to higher values. For the low- and medium-Mg curves implient fractionation of clinopyroxene or feldspar minerals might account for some of the irregularities for such elements as Rb, K, and Sr.

The normalized curves also parallel the base of the diagram, thus signifying little fractionation in this spectrum of elements from a primitive mantle composition. Compare these curves with those for N- and E-MORB in the same diagram which show depletion and enrichment, respectively, of the incompatible elements. This presents some problems in interpretation. Francis et al. (1983) demonstrated that chondrite-normalized REE diagrams for the Chukotat Group in the Cape Smith Belt are consistently light-REE-depleted and interpreted this as evidence for derivation from a source previously depleted by partial fusion. Similarly, Figure 2a shows that the least contaminated Chukotat Group komatiitic rocks show REE profiles with a variety of patterns from light-REE enriched, to parallel, to depleted (Baragar and Scoates, 1987; Picard et al., 1990; Togola, 1992). Picard et al. (1990) interpreted the light-REE-enriched profiles from low-Mg rocks at the base of the Chukotat Group as representing a transitional origin from erratic Povungnituk Group basalt to oceanic Chukotat Group komatiitic basalt, but this would be more difficult to apply to those from higher stratigraphic levels. Hence, the apparent lack of fractionation in the trace-element profiles of Figure 2a is puzzling, but not inconsistent with the varying REE profiles recorded elsewhere. Possibly, a major part of komatiitic lava flows from Smith and Ottawa islands derive from a deeper and less depleted source than those sampled in the central Cape Smith Belt.

ONION-SKIN KOMATIITIC BASALT

Onion-skin lava flows occur principally at two stratigraphic levels: in the upper part of the sequence on Smith Island where they are associated with medium- to low-Mg units, and in the lower part of the mainland sequence where they interlayer with high-Mg lava flows. They have not been reported or described from elsewhere in the Circum-Superior Belt, hence, may be a local phenomenon.

Typically, onion-skin pillow lavas are a light buff, commonly varicolric rock characterized by thick (3–10 cm), scaly rinds surrounding pillows as much as 1 m in diameter. Rarely, the pillows are small, commonly less than 0.5 m in diameter), but, in places, may be large mattresses as much as 10–15 m long. Some interspersed massive lavas contain scaly, varioltic, flow-parallel layers similar to thick pillow rims, so themselves may be parts of gigantic pillow tubes. Quartz commonly appears in interpillow interstices and in the flat drainage cavities at pillow centres, indicating the generally more siliceous nature of these lavas relative to that of the more magnesian units of the komatiitic suite.

Clinopyroxene is the most common phenocryst in these lavas (generally 1–30%), but olivine does appear as a phenocryst phase (1–9%) in ever half the specimens examined and plagioclase in one. The groundmass is typically dominated by clinopyroxene, commonly in quench-like, acicular radiating bundles or fronds interleaved with plagioclase. Generally plagioclase forms about 40–30% of the groundmass. Opaque minerals (now mainly sphene) are fine grained. Rarely, the rocks are fine grained throughout. Quartz is commonly interstitial, accompanied by pumpellyite development. Deformation is slight and textures are readily interpretable in terms of primary minerals.

Little distinction can be seen in the composition (Table 1a) of the onion-skin lavas at the two stratigraphic levels; both are similar to low-Mg komatiitic basalt. In the cross-section of Figure 1 the Smith Island onion-skin lavas are seen to be remarkably uniform in composition and distinct from the low-Mg unit (Chkl) at the base of the sequence in their consistently higher MgO/TiO_2 index. Extended element profiles normalized to primitive mantle of the two onion-skin lavas are compared in Figure 2b with those of the low- to high-Mg units. Except for minor enrichment in the most incompatible of the elements, the Smith Island profiles are generally similar to those of low- to medium-Mg lavas. Those of the onshore peninsula show extraordinary departures from the low- to high-Mg lava profiles at their incompatible end. These are not readily explained.

The compositional similarity of onion-skin and low-Mg lavas makes it difficult to account for their marked differences in physical character. The onion-skin pillows and thick, scaly rims of the onion-skin lavas must signify their much greater viscosity as melts, but their low content of phenocrysts would not indicate unusually low temperatures of emplacement, so perhaps the high viscosity is attributable to very low volatile content, possibly due to loss of pressure and devolatilization prior to eruption.

Layered flows

Layered flows range from meagre olivine concentrations near the base of massive flows, or the massive upper parts of pillow-massive flows (common among flows of the high-Mg unit (Chhkl)) to markedly uparched flow segments well differentiated into sharply defined olivine cumulate bases and feldspathic tops. Of those shown on the map several are extraordinarily well developed and are believed to have acted as major distribution conduits for lavas on the surface. They were probably responsible for some of the compositional variation that characterizes the stratigraphy. A detailed map of one such flow (west-island flow, marked WIF on map), given in Figure 3a, shows the divisions effected by differentiation and gives some insight into the mechanism of formation. Compositional variations through this, and another layered flow (mid-island flow, marked MIF on map) are shown in Figures 3b and 4, respectively. Other layered flows within this map area ("hudson's Bay layered flow" and "North Island spinifex-textured flow") have been described in detail elsewhere (Baragar, 1993, 1994; Baragar and Scoates, 1987).

West-island flow

The western end of the west-island flow shown in Figure 3a extends eastward, with about the same dimensions, for about 1400 m. The differentiated massive part of the flow rests on a pillowed base (part of the same flow) and is divided into lower dolerite (chkl zone), succeeding olivine cumulate, and upper gabbro layers with pegmatitic segregations at mid-upper levels (not shown). Clinopyroxene concentrations observed at the top of the olivine cumulate zone in one of the segregations (Fig. 3a) suggest that the segregations may be feeders to the flow tops. The pegmatitic segregations in gabbroic zones appear to be sharply divided from one another along a foliated contact. This probably indicates that segregation took place in a mobile regime. Of particular interest is the abrupt uparching at the western termination of the flow where overlying pillow lavas parallel the steeply inclined western boundary, indicating that they were in place prior to the uplift that accompanied thickening. Hence, one can postulate that after the initial emplacement of the lava flow and its subsequent burial by overlying pillows, renewal of the magma supply resulted in a surge of magma along the still-molten interior and the rapid uplift of its roof and superincumbent lavas. This was partially achieved by the spray of fluidal, flow-banded, westward from the (then) axial interior, which segmented and tilted the solidified roof and overlying lavas. Thinning westward from the magma supply seems to have been sufficiently prolonged to allow olivine accumulation to reach a third of the thickness of the flow before the overriding differentiate was intruded by the pegmatitic segregations at the lower termination of the flow.

A compositional profile through the west-island flow is shown in Figure 3b. Several features are of particular note.

- The uniformity of MgO through the bottom and top of the cumulate layer is indicative of its comparative and implies a differential flow between cumulate base and crystal-free top that draws pure fluid from the crystal mush of the former.
- Both CaO and Al_2O_3 decline steadily upward in the gabbroic upper zone, suggestive of feldspar fractionation to more alkaline compositions.
- Pegmatitic segregations at mid-upper levels result in local enrichment of FeO* (total iron as FeO), TiO_2 and Zr.

Mid-island flow

Like the west-island flow, this flow is over 80 m thick and has a length at surface of several hundred metres. Its compositional profile (Fig. 4) shows a compact olivine cumulate base similar to the west-island flow, but with better definition of its upper contact against the overlying gabbroic zone. The contact is sharp, which tends to support the view that it is formed at a zone of differential flowage. Setting in a static environment would be gradational. Other features of interest in the mid-island flow are: the enrichment of Al_2O_3, Ba, and Sr just above the cumulate base; and concentration of FeO*, TiO_2, and Zr at mid-level of the gabbroic zone. These can be attributed to some degree of plagioclase concentration at the base of the upper zone and to pegmatitic segregation near its centre.

Discussion

The importance of the layered flows is their ability to effect a separation of olivine and, to lesser degree, clinopyroxene, from erupted high- to medium-Mg lavas on surface prior to their final emplacement. The olivine-rich bases, which accumulates during transport, lags behind the more fluid, relatively crystal-free, top which arrives at its final destination variably depleted in olivine and, to some degree, clinopyroxene. These are the essential distinctions relating high-Mg to medium- and low-Mg lavas, which makes the layered flows an appropriate mechanism for achieving some of the variability in composition seen within the Chukotat Group.

THICK FLOWS

Extraordinarily thick lava flows (unit Chkl) are present in both the western part of Smith Island and near the southern margin of the mainland peninsula. The western Smith Island flow is about 270 m thick, and although complicated by faulting, is at least several kilometres in length. The peninsular unit is a composite of massive flows, some of which are unusually thick (200 m), but which may be down to as little as 20 m. Together they form a resistant, physiographic unit that is distinctive in the field and in aerophoto.

The chemistry (Table 1a, Baragar, 1983) though the thick flow on Smith Island shows significant petrochemical variations (Fig. 5) that differ from the layered flows described above. The olivine cumulate base is thin (~ 20 m) — despite the high-Mg character of the flow base and subjacent pillows — and grades (rather than passing abruptly) into the dolerite zone above. The bulk of this overlying zone is a uniform white-weathering, feldspathic, subophitic dolerite that passes up into a heterogeneous zone, about 20–30 m thick, composed of a swirly mix of mafic and felsiphatic components. Above, the rock grades into the gabbroic upper zone. Here, pegmatitic segregations are relatively rare.

The flow is covered by about 40 m of flow-top breccia containing some rounded, chilled, seemingly leucocratic lava masses.

Certain aspects of the compositional variation seen in Figure 5 are important to note in regard to the origin of the flow. The MgO/CaO index shows that the olivine cumulate base is relatively rich in olivine, as expected. The MgO/TiO_2 is an attributable better attribute as it is not dependent on variable olivine vs. Al_2O_3, CaO and Si proportions. Note that the lower part of the dolerite zone (below the zone of mixed mafic-felsiphatic components), indicative of plagioclase accumulation in this region, has the lowest index value. The MgO/TiO_2 and Zr show a distinct increase over a broad zone in the upper part of the flow, but decline toward its roof. Alkalis reach a maximum in the same zone. Hence, this probably marks a region of accumulation of late crystallizing components of magmatic mass. Finally, compositions in the rapidly cooled zones at the base and top of the flow are markedly different. The basal dolerite and related underlying pillows are medium- and/or high-Mg units (~15% MgO), whereas the upper roof and lava masses in the overlying flow-top breccia are of low-Mg (7–10% MgO) composition. These features might be best explained by a lava lake or thick ponded flow. Fractionation of olivine, followed by that of clinopyroxene and plagioclase, in the lower, cooler portions of minerals of differing densities and might, with varying and perhaps erratic intensities, lead to zones of mixed rock types, such as the mid-flow zone of mixed mafic-felsic materials. With declining convection rates, parts of the lava surface would stiffen to the point where it could bar an increasing load of debris, resulting from its own surface agitation as well as ejecta from neighbouring, still open and active convection cells. Hence, the solidifying lake surface and incumbent aggregate would be of a composition consistent with the differentiated liquid reached at that stage at the flow top. Within the magma body, closure of the surface would trap late crystallizing fluids within the upper parts of the body. Thus, the evidence clearly supports the concept of a lava lake as the progenitor of this feature.

Not enough data exists to speculate on the origin of the thick flows of the mainland peninsula. The few analyses available (Baragar and Scoates, 1987; Picard et al., 1990) are similar to those of the Smith Island flow and the Chukotat Group komatiitic lavas generally. Their thickness, however, is unusual.

CHUKOTAT GROUP SILLS

Folded gabbro sills that intrude the upper levels of Povungnituk Group sedimentary rocks on the southern margin of the mainland peninsula are considered to be comagmatic with the Chukotat Group. They are of komatiitic basalt composition, as listed in Table 1a and many of the gabbroic sills on Te Gobin are probably of this composition also (Parent). In particular, a thick, sill-like gabbroic body on the southeast coast of Te Gobin is similar in chemical values of average Te Gobin Povungnituk Group basalt: MgO (6.49%), TiO_2 (2.00%), and Cr (158 ppm). Also compare the normalized trace-element profile of average Te Gobin Povungnituk Group gabbro shown in Figure 2b with corresponding profiles of average komatiitic suite lavas and Te Gobin Povungnituk Group basalt. Despite erratic offsets of some of the REEs, the Chukotat sill profile fits more closely the komatiitic profiles than that of the Povungnituk Group lavas. Clearly, these sills are part of the Chukotat Group magmatic suite that have intruded the Povungnituk Group. Regional maps of the Cape Smith Belt (Hynes and Francis, 1982; Hynes and Rochette, 1990) show a number of gabbroic sills and the presence of similar sills at the correspondent stratigraphic level throughout much of the Cape Smith Belt (St-Onge et al., 2000). Thus, there is strong support for the view that the magmatic episodes responsible for these two major volcanic assemblages, the Povungnituk and Chukotat, are not greatly separated in time or space and are an interrelated part of an ongoing tectonic process.

SUMMARY AND CONCLUSIONS

The principal stratigraphic components of the Cape Smith Belt — the basal Povungnituk and overlying Chukotat groups — are represented in the map area by the uppermost 2000 m of Povungnituk Group, comprising tholeiitic pillow basalt and black shale, and the lowermost 7000 m of Chukotat Group of komatiitic lava flows exposed in cross-section. Their relationship is poorly defined, but is interpreted as an overthrust of Chukotat Group over Povungnituk Group. Thick gabbro sills within Povungnituk Group sedimentary rocks subjacent to the thrust contact have the chemical attributes of Chukotat Group volcanic rocks and are interpreted as intrusives of Chukotat Group magma into the earlier succession. Hence, the time and space separation of the two groups may not be of major significance.

The Chukotat Group is subdivided into a number of distinctive units based partly on magnesium content: low- (~ 10% MgO), medium- (10–15% MgO), and high-Mg (~15% MgO) lavas; and partly on distinguishing characteristics: onion-skin (thick, flaky pillow rims) pillow lavas and extraordinarily thick (~250–300 m) massive, layered flows. The onion-skin lavas are compositionally similar to the low-Mg komatiitic basalt (Ivakkak Group low-Mg unit), but were seemingly emplaced as highly viscous flows, possibly the result of pre-eruptive devolatilization. The thick flows show some of the attributes of, and are interpreted as having formed as, lava lakes. Among the high- and medium-Mg units in particular, are a number of layered flows formed of thickened, uparched segments of massive, or the massive parts of pillow-massive flows. These are typically differentiated into an olivine cumulate base and overriding gabbroic top, itself commonly differentiated to more evolved compositions. In one layered flow within the map area, spinifex texture, including parallel spinifex (primarily clinopyroxene spinifex), forms the major part of the upper zone.

Broadly based chemical profiles normalized to primitive mantle highlight the distinctions between Povungnituk Group and Chukotat Group magmas and between various units of the latter. For most of the less compatible elements Povungnituk Group lavas flows are generally tholeiitic compared to the Chukotat Group, but for the compatible elements, they are more depleted. This is indicative of the more evolved state of the former relative to the latter consistent with previously expressed views on their respective relations to an opening continental rift (e.g. Francis et al., 1983). Among Chukotat Group units, the high- to low-Mg lavas, showing roughly parallel profiles on multi-element, mantle-normalized plots, are consistent with their interpretation as parts of a single, consanguineous, komatiitic suite. Incompatible and therefore would be consistent with their derivation from a primitive mantle source; however, this would not be compatible with previous interpretations (based on REE profiles) from the Cape Smith Belt that projected an N-MORB source. Possibly the differences reflect a different source for magmas erupted along the flanking as opposed to the arch position (Cape Smith Belt) of the Circum-Superior Belt.

ACKNOWLEDGMENTS

The author is pleased to acknowledge the very able assistance rendered in the field by Mathieu Piché (senior assistant), Jacques Lefebvre, and John Leoh in 1981 and Lyn MacIntyre and Marc Baragar in 1982 and is grateful for their cheerful companionship under often trying conditions. To the people of Akulivik village are also owed thanks for the friendly reception and the assistance given the field party whenever it was required.

REFERENCES

Baragar, W.R.A.

1983: The Circum-Ungava belt of eastern Hudson Bay; geology of the Cape Smith region; *in* Current Research, Part A, Geological Survey of Canada, Paper 83-1A, p. 305–309.

1994: Pillow formation and layered flows in the Circum-Superior Belt of eastern Hudson Bay; Canadian Journal of Earth Sciences, v. 31, p. 781–792.

Baragar, W.R.A. and Pichè, M.

1982: The Circum-Superior Belt of eastern Hudson Bay: the geology of the Ottawa Islands and the Cape Smith Region; *in* Current Research, Part A, Geological Survey of Canada, Paper 82-1A, p. 11–15.

Baragar, W.R.A. and Scoates, R.F.J.

1981: The Circum-Superior Belt: Proterozoic plate margin?; *in* Precambrian Plate Tectonics, (ed.) A. Kroner; Elsevier, Amsterdam, Netherlands, p. 297–330.

1987: Volcanic geochemestry of the northern segments of the Circum-Superior Belt of the Canadian Shield; *in* Geochemistry and Mineralization of Proterozoic Volcanic Suites, (ed.) T.C. Pharoch, R.D. Beckinsala, and D. Rickard; Geological Society Special Publications, no. 33, p. 113–131.

Francis, D.M., Ludden, J.M., and Hynes, A.J.

1983: Magma evolution in a Proterozoic rifting environment; Journal of Petrology, v. 24, p. 556–582.

Hynes, A. and Francis, D.M.

1982: A transect of the early Proterozoic Cape Smith tholbelt, New Quebec; Tectonophysics, v. 88, p. 23–59.

Kerrich, R. and Wyman, D.A.

1996: The trace element systematics of igneous rocks in mineral exploration: An overview; in Trace Element Geochemistry of Volcanic Rocks: Applications for Massive Sulphide Explorations, (ed.) D.A. Wyman; Geological Association of Canada, Short Course Notes, v. 12, p. 1–50.

Picard, C., Lamothe, D., Piboule, M., and Oliver, R.

1990: Magmatic and geotectonic evolution of a Proterozoic oceanic basin system: the Cape Smith Thrust-Fold Belt (New Quebec); Precambrian Research, v. 47, p. 223–249.

St-Onge, M.R., Baragar, W.R.A., and Henderson, I.

2006: Géologie, cahirure de Cape Smith et environs, péninsule de l'Ungava, Québec–Nunavut / Geology, Cape Smith Belt and adjacent domains, Ungava Peninsula, Quebec–Nunavut; Geological Survey of Canada, Open File 4936, scale 1:300 000.

St-Onge, M.R., Lucas, S.B., and Perrish, R.R.

1992: Terrane Accretion in the Internal Zone of the Ungava Orogen, northern Quebec, Part 1: Tectonostratigraphic assemblages and their tectonic implications; Canadian Journal of Earth Sciences, v. 29, p. 746–764.

Sun, S.-s. and McDonough, W.F.

1989: Chemical and isotopic systematics of oceanic basalts: implications for mantle compositions and processes; *in* Magmatism in the Ocean Basins, (ed.) A.D. Saunders and M.J. Norrey; Geological Society Special Publications, no. 42, p. 313–345.

Togola, N.

1992: Géologie de la région de la baie Korak, Ministère de l' Énergie, des Mines, et des Ressources du Québec, ET 91-07, 39 p.



Figure 2a



Figure 2b

Figure 2. a) Profiles of average analyses for each of low-Mg, medium-Mg, and high-Mg komatiitic basalt samples normalized to primitive mantle compared to spot readings for N-MORB and E-MORB. b) Region subdivided by profiles of komatiitic lava flows shown in Figure 2a compared with corresponding profiles of Povungnituk Group tholeiitic basalt, a Chukotat Group sill from near base of the Chukotat Group, and onion-skin lava flows. Normalizing values and order of compatibility are from Sun and McDonough (1989), supplemented by data for major elements from Kerrich and Wyman (1996).



Figure 3a

Figure 3b

Figure 3. a) Detailed map of the western end of the west-island flow (WIF on map). b) Chemical variation through a sampled section of the west-island flow. Note MgO and Cr bulges marking the cumulate zone and TiO_2 and Zr enrichments signifying pegmatitic segregations. The underlying pillowed zone is interpreted as the pillowed base of the same flow.



Figure 4

Figure 4. Chemical profiles through a sampled section of the mid-island flow (MIF on map). Like the west-island flow the major features are the cumulate zone and pegmatitic segregations marked by enrichments of MgO-Cr and TiO_2-Zr, respectively. In this flow, close sampling details the sharp upper boundary of the cumulate zone, indicative of a mobile settling regime.



Figure 5

Figure 5. Chemical profiles of a sampled section through an extraordinarily thick flow (unit Chkl) of the komatiitic suite (TFS on map). Note the thin cumulate zone and its gradational top, probably indicative of static settling. This, together with lack of conformity in compositions between chilled base and top of the flow are suggestive of lava lake conditions (see text). Underlying pillows are interpreted as part of the same flow.


78°45'
40'
35'
30'
25'
20'
15'
10'
05'
78°00'
61°00'
55'
50'
45'
60°40'
BAIE D'HUDSON
HUDSON BAY
MOSQUITO BAY
Pointe Chronan
Pointe Migeon
Pointe Akulivik
Cape Smith
Hospital Bay
Knight Harbour
Babs Bay
Lac Guillaume-Delisle
Lac Minto
Lac Allaire
Lac Chevrier
Lac Gilbert
Lac Lemoine
Lac Bourgeois
Lac Bellin
Lac Bobin
Lac Lacroix
Lac Lemaire
Lac Lemoyne
Lac Lemeunier
Lac Lemay
Lac Lemieux
Lac Lemire
Lac Lemoine
Lac Lemoyne
Lac Lemeunier
Lac