



- LEGEND**
- SURFICIAL GEOLOGY**
- QUATERNARY**
- Glacier:** A mass of ice formed from compacted snow in an area where snow accumulation exceeds melting and sublimation.
- POST-FRASER GLACIATION**
- NONGLACIAL ENVIRONMENT**
- O** **ORGANIC DEPOSITS:** peat and muck; 1 to 10 m thick (typically 2 to 3 m); forming fans and bogs; organic deposits too small to be shown at this scale occur within other units; common within abandoned meltwater channels.
 - Ap** **ALLUVIAL (FLUVIAL) DEPOSITS:** gravel and sand with minor silt and clay, deposited by streams; commonly stratified; generally well sorted except in alluvial fans.
 - At** **Floodplain sediments:** sand and silt, commonly including organic materials and nodules; in many places, by gravel; 1 to 3 m thick; occurring as flat surfaces close to river level; prone to flooding.
 - Af** **Terrace sediments:** stratified sand and gravel overlain by a veneer of sand and silt; 2 to 10 m thick; forming terraces well above flood level.
 - Al** **Fan sediments:** poorly sorted sand and rubble, with clams; generally 2 to 15 m thick; forming fans at the toe of slopes.
 - Cd** **COLLUVIAL DEPOSITS:** clams and rubble deposited by various mass-wasting processes, ranging from slope wash to rock fall; composition dependent on source materials.
 - Cs** **Landslide debris:** mostly unconsolidated sediments, with texture dependent on source materials; generally 1 to 10 m thick, but may exceed 10 m near the toe of large landslides; forming hummocky accumulations on lower slopes and valley floor. Where possible, landslides were identified by type: Ch-ds, debris flow deposit; Ch-ds, debris avalanche; Ch-ds, debris slide; Ch-rs, rock slide; Ch-rs, rock avalanche; Ch-sa, snow avalanche track.
 - Cs** **Slope colluvium:** rock fragments in a matrix of boulders, gravel, sand, silt, and minor clay; 1 to 10 m thick; formed by bedrock weathering or reworking of unconsolidated deposits on steep (>30°) slopes; commonly gullied.
 - Cs** **Talus:** rubble and block accumulations at the bottom of steep (>40°) slopes; 1 to 10 m thick; forming aprons and cones.
 - Cv** **Colluvial veneer:** rock fragments in a matrix of boulders, gravel, sand, silt; usually <3 m thick; formed by bedrock weathering or reworking of unconsolidated deposits.
- FRASER GLACIATION (LATE WISCONSINAN)**
- PROGLACIAL AND GLACIAL ENVIRONMENT**
- Gmh** **Glaciomarine ice-contact deposits:** sand and gravel, well to poorly sorted, and commonly stratified; deposited by glacial meltwater; bedding disrupted locally due to melt of glacier ice.
 - Gmt** **Glaciomarine terrace sediments:** sand and gravel, stratified to massive; 1 to 10 m thick; forming flat surfaces perched well above alluvial deposits or associated with meltwater channels; may be faulted.
 - Gmd** **Proglacial deltaic sediments:** sand and gravel with minor silt and clay; on average 10 m thick, but can be >10 m; commonly overlie glaciomarine silt and clay; may form, in part, inclined surfaces. There is an inferred transition between the terraced and marine proglacial sediments.
 - Gmv** **Glaciomarine veneer:** sand and gravel, well to poorly sorted, and commonly stratified; 1 to 3 m thick; deposited by glacial meltwater; bedding disrupted locally due to melt of associated glacier ice.
 - Glf** **GLACIOFLUVIAL DEPOSITS:** sand and gravel, well to poorly sorted, and commonly stratified; deposited by glacial meltwater; bedding disrupted locally following the melting of supporting ice.
 - Gf** **Ice-contact deposits:** sand and gravel, stratified to massive and commonly faulted; generally >3 m thick; forming hummocky surfaces.
 - Gf** **Glaciofluvial terrace sediments:** sand and gravel, stratified to massive; 1 to 10 m thick; forming flat surfaces perched well above alluvial deposits or associated with meltwater channels.
 - Gv** **Glaciofluvial veneer:** sand and gravel, well to poorly sorted, and commonly stratified; deposited by glacial meltwater; bedding disrupted locally following melting of supporting ice; 1 to 3 m thick.
- GLACIAL ENVIRONMENT**
- Tu** **TILL:** Poorly sorted clastic material consisting of pebbles, cobbles, and boulders in a sandy to clayey matrix, directly deposited by glaciers; includes colluvium (reworked till) on steep slopes, and small units of glacial till, especially in valley bottoms and near the mouths and banks of meltwater channels; till surface is commonly rilled on steep slopes.
 - Tu** **Till blanket:** continuous till cover with few bedrock outcrops; 1 to 3 m thick on average; conforming to and may be locally obscuring morphology of underlying units.
 - Tv** **Till veneer:** discontinuous till cover with abundant bedrock outcrops; 1 m thick on average; reflecting topography of underlying bedrock.
- PRE-QUATERNARY**
- R** **BEDROCK:** sedimentary, low-grade metamorphic, volcanic, and intrusive rocks of Jurassic to Quaternary age; including, in places, till veneer, drift, and colluvium.
- Geological boundary (defined, inferred)
- Limit of mapping
- Escarpment
- Large meltwater channel
- Small meltwater channel
- Sand and gravel pit (large, small)
- Travel directions of landslides, mainly debris flows and snow avalanches
- Crest
- Fossil

ACKNOWLEDGEMENTS

This mapping project was funded by Climate Change Action Fund A769A. The author wishes to thank R. Schwartz and F. Kertland of CCAF for financial and managerial guidance of the project. The author is grateful to T. Millard and M. Gaudreau from BC Ministry of Forests and Range for providing digital TRIM data as well as helpful discussions. J. Chagnon, O. Hung, and F. Beaumais provided invaluable critical reviews. K. Stammers, M. Linn, C. Hickman, N. Jordan, F. Beaumais, M. Bourneau, R. Hermann, C. Jermyn, and B. Strick provided field and/or technical assistance. R. Poirier, M. Booth, F. Chagnon, and S. Herd provided GIS assistance for parts of the map area. M. Millard and N. Cole are thanked for their assistance in GIS and creating the final map product. Many thanks go to my GSC colleagues for helpful discussions and advice throughout the project.

REFERENCES

Cruen, D.M. and Veres, D.J.
1996. Landslide types and processes. In Social Report 247, Landslide investigation and mitigation. A.K. Turner and R.L. Schuster (eds). National Research Council, Transportation Research Board, Washington D.C., p. 36-76.

802.003	802.004	802.005	802.006	802.007	802.008
802.009	802.010	802.011	802.012	802.013	802.014
802.015	802.016	802.017	802.018	802.019	802.020
802.021	802.022	802.023	802.024	802.025	802.026
802.027	802.028	802.029	802.030	802.031	802.032
802.033	802.034	802.035	802.036	802.037	802.038
802.039	802.040	802.041	802.042	802.043	802.044
802.045	802.046	802.047	802.048	802.049	802.050
802.051	802.052	802.053	802.054	802.055	802.056
802.057	802.058	802.059	802.060	802.061	802.062
802.063	802.064	802.065	802.066	802.067	802.068
802.069	802.070	802.071	802.072	802.073	802.074
802.075	802.076	802.077	802.078	802.079	802.080
802.081	802.082	802.083	802.084	802.085	802.086
802.087	802.088	802.089	802.090	802.091	802.092
802.093	802.094	802.095	802.096	802.097	802.098
802.099	802.100	802.101	802.102	802.103	802.104
802.105	802.106	802.107	802.108	802.109	802.110
802.111	802.112	802.113	802.114	802.115	802.116
802.117	802.118	802.119	802.120	802.121	802.122
802.123	802.124	802.125	802.126	802.127	802.128
802.129	802.130	802.131	802.132	802.133	802.134
802.135	802.136	802.137	802.138	802.139	802.140
802.141	802.142	802.143	802.144	802.145	802.146
802.147	802.148	802.149	802.150	802.151	802.152
802.153	802.154	802.155	802.156	802.157	802.158
802.159	802.160	802.161	802.162	802.163	802.164
802.165	802.166	802.167	802.168	802.169	802.170
802.171	802.172	802.173	802.174	802.175	802.176
802.177	802.178	802.179	802.180	802.181	802.182
802.183	802.184	802.185	802.186	802.187	802.188
802.189	802.190	802.191	802.192	802.193	802.194
802.195	802.196	802.197	802.198	802.199	802.200

Author: A. Blais-Stevens
Geology by A. Blais-Stevens, 2004-2007

Compilation and interpretation was carried out using British Columbia 1994 colour aerial photography series 30P024, at 1:15,000 scale.

Digital cartography by N. Cole, Data Dissemination Division (DDD)

This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (STPS) Quality Management System, registered to the ISO 9001:2000 standard.

SURFICIAL GEOLOGY AND LANDSLIDE INVENTORY OF THE LOWER SEA TO SKY CORRIDOR
BRITISH COLUMBIA

Scale 1:50 000/Echelle 1/50 000

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2008

Projection transversale universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du chef du Canada 2008

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from Terrain Resource Information (TRIM), modified by DDD

Shaded relief image prepared by DDD, derived from the digital elevation model, based on TRIM contours elevation data
illumination: azimuth 315°, altitude 45°, vertical factor 1x

Magnetic declination 2008, 18°2' E, decreasing 12.4' annually.

Elevations in metres above mean sea level
Contour interval 20 m

OPEN FILE DOSSIER PUBLIC 5322

2008

Les données publiques sont disponibles en format PDF sur le site Web de la Commission géologique du Canada.



Recommended citation:
2008. Surficial geology and landslide inventory of the lower sea to sky corridor, British Columbia. Geological Survey of Canada, Open File 5322, scale 1:50 000.