



This legend is common to GSC Open File maps produced for NTS sheet 84 P. Not all map units in the common legend appear on this map.

**NOTE:** In areas where the surficial cover forms a complex pattern, the area is coloured according to the dominant unit and labelled in descending order of cover (e.g. O-Tr). Where buried aggregate deposits (sand and gravel - commonly associated with G1 or G2 surficial units) are known, or suspected, areas are coloured according to the overlying unit and labelled in the following manner: Lw/Gc.

**QUATERNARY SURFICIAL DEPOSITS**

**POST LAST GLACIATION**

**NONGLACIAL ENVIRONMENTS**

**ORGANIC DEPOSITS:** peat and muck: 1 to 3 m thick on average; formed by the accumulation of plant material in various stages of decomposition; generally occurs as flat, wet terrain (swamps and bogs) over poorly drained substrates.

**O<sup>1</sup>** Bog peat: sphagnum or forest peat formed in an ombrotrophic environment; wet terrain; may be treed or treeless; O<sup>1</sup>h, hummocky, mounds and plateaus; area may be underlain by ground ice or shallow permafrost conditions; O<sup>1</sup>k, thermokarst terrain related to melting ground ice.

**O<sup>2</sup>** Fen peat: peat derived from sedges and partially decayed shrubs in a eutrophic environment; forms relatively open peatlands with a mineral-rich water table that percolates seasonally near the surface; generally covered with low shrubs and sometimes a sparse layer of trees.

**O** Undifferentiated bog and fen deposits: O<sup>h</sup>, undifferentiated hummocky bog and fen deposits; area may be underlain by ground ice or shallow permafrost conditions; O<sup>u</sup>, undifferentiated bog and fen deposits with thermokarst terrain related to melting of ground ice; O<sup>c</sup>, undifferentiated bog and fen deposits cut by numerous subparallel channels on gentle slopes.

**COLLUVIAL DEPOSITS:** mass wasting debris; poorly sorted, massive to stratified debris deposited by direct, gravity-induced movement; composition dependent on source material.

**Ch** Landslide and slump debris: active and inactive landslides; hummocky topography; diamicton, generally 1 to 10 m thick, but may exceed 10 m near the toe of large landslides.

**Cv** Colluvial veneer: thin and discontinuous cover of slumped and/or soliflucted material <1 m thick; overlies bedrock or till.

**C** Undifferentiated colluvial deposits.

**ALLUVIAL DEPOSITS:** sorted gravel, sand, minor silt, and organic detritus deposited by streams; commonly stratified.

**Ap** Floodplain deposits: sorted gravel, sand, silt, and organic detritus >1 m thick; forming active floodplains close to river level with meander channels and scroll marks.

**At** Fluvial terrace deposits: inactive terraces above modern floodplains >2 m thick; represents a potential aggregate source.

**Al** Alluvial fan deposits: poorly sorted gravel, sand, and organic detritus >1 m thick.

**Av** Alluvium veneer: <1 m thick; primarily as uniform sheets of slope wash on gentle slopes.

**A** Undifferentiated fluvial deposits.

**L<sup>1</sup>** LACUSTRINE DEPOSITS: sand, silt, and minor clay deposited in a former lake; >1 m thick; generally overlain by organic deposits; exposed by recent fluctuations in lake levels.

**NONGLACIAL AND PROGLACIAL ENVIRONMENTS**

**EOLIAN DEPOSITS:** wind-deposited medium to fine sand; derived from detrital or glaciolacustrine deposits; in some areas eolian sediments are thin or absent between dunes.

**Er** Ridged eolian deposits: forming dunes; generally >2 m thick.

**Ev** Eolian veneer: discontinuous veneer of eolian sediments; <1 m thick.

**POSTGLACIAL OR LATE WISCONSINAN**

**PROGLACIAL AND GLACIAL ENVIRONMENTS**

**GLACIOLACUSTRINE DEPOSITS:** fine sand, silt, and clay, with minor debris-flow diamicton, deposited in glacial-dammed lakes in valleys and along the margin of the retreating Laurentide Ice Sheet; usually overlain by organic deposits in lowlands.

**Lb** Glaciolacustrine blanket: >1 m thick.

**Lv** Glaciolacustrine veneer: thin and discontinuous; <1 m thick.

**GLACIOFLUVIAL DEPOSITS:** well to poorly stratified sand and gravel; minor diamicton; deposited behind, at, or in front of the ice margin by glacial meltwater; represents a potential aggregate source.

**G** Proglacial outwash: cross-stratified gravel and sand deposited in front of the ice margin; G<sub>s</sub>, outwash plain deposits, generally 1 to 5 m thick, generally mantle valley floors and surfaces adjacent to glacial meltwater channel margins; G<sub>l</sub>, outwash terrace deposits, generally associated with meltwater channels and canyons; 1 to 10 m thick; G<sub>t</sub>, glaciofluvial delta deposits; 1 to >30 m thick; G<sub>v</sub>, glaciofluvial veneer thin and discontinuous; <1 m thick; G<sub>g</sub>, glaciofluvial fan deposits; >1 m thick.

**Gi** Ice-contact stratified drift: poorly-sorted sand and gravel with minor diamicton; deposited in contact with the retreating glacier; 1 to >20 m thick; G<sub>h</sub>, hummocky topography relating to melting of underlying ice; G<sub>l</sub>, surface marked by kettle holes; G<sub>r</sub>, esker ridges; G<sub>k</sub>, kame terraces; G<sub>l</sub>, ice-contact glaciofluvial delta deposits; 1 to >30 m thick, surface marked by kettles.

**TILL:** diamicton deposited directly by the Laurentide Ice Sheet; sandy to clayey matrix with scattered clasts of various lithologies, including many Canadian Shield, carbonate, and sandstone erratics; clast content is typically low (<10 %).

**Tb** Till blanket: >1 m thick, continuous till cover forming undulating topography that locally obscures underlying units.

**Ts** Streamlined and fluted till: >1 m thick, till surface marked by streamlined landforms including flutes and drumlins.

**Th** Hummocky till: >1 m thick; hummocky till surface.

**Tr** Ridged till deposits: >1 m thick, moraines or crevasse fillings forming a ridged topography.

**Tv** Till veneer: <1 m thick, discontinuous till cover; underlying bedrock topography is discernible.

**PRE-QUATERNARY BEDROCK**

**R** Sedimentary bedrock: Cretaceous Fort St. John Group shales (including the Shaleburg Formation) and Dunsmuir Formation sandstone exposed in highlands and along meltwater channel and canyon walls.

**LEGEND**

Geological boundary (defined, approximate) .....  
 Patterned ground .....  
 Obowr .....  
 Meltwater channel or undercut channel, small (paleoflow direction known, unknown) .....  
 Meltwater channel, large (paleoflow direction known, unknown) .....  
 Kettle .....  
 Esker .....  
 Major moraine .....  
 Minor moraine or crevasse filling .....  
 Ice moulded form in till (direction of flow not inferred) .....  
 Field observation site .....▲

**NOTES**

The surficial geology of the Thinahtea Lake (NTS 84 P19) map area is dominated by the effects of continental glaciation during the Late Wisconsinan (ca. 25 000-10 000 years ago). In general the ice sheet advanced from the northeast, but the ice thinned during deglaciation and divided into distinct lobes. The map area became dominated by one lobe flowing southward from the Mackenzie River valley in the north, and a second lobe in the south, occupying the Skeelike River. Glacial flutings in the northern part of the map area record the southern flow of the northern lobe. Thinahtea Lake itself occupies a large meltwater channel that was probably the mouth of a large subglacial tunnel emanating from the northern ice lobe. Numerous eskers lie along the northern edge of the lake. Initially the northern and southern lobes were in contact along an area of hummocky moraine, just south of the Pellet River, but once they separated meltwater collected between the lobes along the lowland occupied by the modern Pellet River with drainage to the west. The western part of the Pellet River valley, however, was initially occupied by the northern ice lobe and the main drainage was forced southward along the northern margin of the Esker Plateau. The Pellet River did not occupy its present course until the northern ice lobe retreated northward. Much of the map area is underlain by thick clayey till and glaciolacustrine deposits, which are poorly drained and covered by extensive muskeg. Areas of thick peat are likely underlain by permafrost and probably contain significant amounts of groundice.

Author: J.M. Bednarski

Geology by J.M. Bednarski, 2003-2005, with additional data provided by T. Ferby and A. Hicks, Resource Development and Geoscience Branch, BC Energy, Mines, and Petroleum Resources

Airphoto interpretation by J.M. Bednarski, 2006

Compilation of geology was onto 1:40 000 orthorectified airphoto mosaic by J.M. Bednarski

Digital cartography by N.Côté, Data Dissemination Division (DDD)

This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (STPSS) Quality Management System, registered to the ISO 9001:2000 standard

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map provided by the BC Watershed Atlas (1:50 000, TRIM base), modified by J.M. Bednarski

Shaded relief image prepared by DDD, derived from the digital elevation model supplied by J.M. Bednarski, based on SRTM imagery. Illumination: azimuth 310°, altitude 45°, vertical factor 4.8x

Magnetic declination 2008, 21°30' E, decreasing 23.8 annually



OPEN FILE 5479  
 SURFICIAL GEOLOGY  
**THINAHTEA LAKE**  
 BRITISH COLUMBIA  
 Scale 1:50 000/Échelle 1/50 000

Kilometres 1 0 1 2 3 4 Kilomètres

Universal Transverse Mercator Projection  
 North American Datum 1983  
 © Her Majesty the Queen in Right of Canada 2008

Projection transversale universelle de Mercator  
 Système de référence géodésique nord-américain, 1983  
 © Sa Majesté la Reine du chef du Canada 2008

84 P13	84 P14	84 P15	84 P16
OF425			OF480
84 P17	84 P18	84 P19	84 P20
OF486	OF506	OF505	OF5479
84 P21	84 P22	84 P23	84 P24
			OF481

**OPEN FILE DOSSIER PUBLIC 5479**

GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA

2008

Open file maps are products that have not gone through the GSC formal publication process.

Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la GSC.