CONTRIBUTIONS

1826

CANADIAN NATURAL HISTORY.

TO

BY W. S. M. D'URBAN AND ROBERT BELL.

(Extracted from the Report of the Geological Survey for 1858.)

This document was produced by scanning the original publication.

Ce document est le produit d'une numérisation par balayage de la publication originale.

Montreal : PRINTED BY JOHN LOVELL, ST. NICHOLAS STREET. 1860.

CONTRIBUTIONS

TO

CANADIAN NATURAL HISTORY.

Pelley of the River Day BY Print

and the set of the second states the second states

W. S. M. D'URBAN AND ROBERT BELL.

(Extracted from the Report of the Geological Survey for 1858.)

Montreal: PRINTED BY JOHN LOVELL, ST. NICHOLAS STREET. 1860.

CONTRIBUTIONS

. had evel as the set of a feet of a lot of the lot of the set of

. Land Considering Stability and West seen in the land throughout the

CANADIAN NATURAL HISTORY.

Catalogue of Animals and Plants,* collected and observed in the Valley of the River Rouge and the neighbouring Townships, in the Counties of Argenteuil and Ottawa. By Mr W. S. M. D' Urban, Assistant to Sir W. E. Logan in 1858.

VERTEBRATA.

CLASS MAMMALIA.

ORDER CHEIROPTERA.

 Vespertilio subulatus, Say.—Rouge, August 8th and 10th. There are probably several species of bats in the district, but this is the only one of which a specimen was obtained.

ORDER CARNIVORA.

 Ursus Americanus, Pallas.—Although no bears were actually seen by us, yet the evidence afforded by recent traces of them, and the information received from settlers and others, induced me to believe that they were numerous in the district.

 Mustela martes, Linn.—The pine marten does not appear to be plentiful. One specimen was seen at Hamilton's Farm on the Rouge, about fify miles from its mouth.
 " Canadensis, Schreber.—Said to be common about Hamilton's Farm; I saw a specimen which had been shot there.
 " vison, Gmel.—Abundant throughout the district. *

* The list of plants having been taken by Mr. D'Uurban to England for the purpose of reference in regard to some points, was unfortunately lost on its retumn in the Hungarian, and there has not been time to prepare another.

- 6. Mephitis chinga, Tiediman.-Common about the settlements in Grenville, &c.
- 7. Lutra Canadensis, Sabine.—Many were seen in the lakes throughout the district.
- 8. Vulpes fulvus, Desm.-Reported to be common.

ORDER RODENTIA.

- 9. Castor fiber, Linn.—Appears to be nearly extinct in the parts we explored, but seen by Sir. W. Logan between two and three miles east of Hamilton's Farm, and said to be numerous about forty miles above it.
- 10. Fiber Zibethicus, Cuvier .- Very numerous throughout the district.
- Arctomys monax, Linn.—Said to be common about clearings in Grenville. A specimen was given to me which had just been killed in that township.
- 12. Tamias Lysteri, Ray.—Township of Montcalm and about Hamilton's Farm; rare.
- 13. Sciurus Hudsonius, Pennant .- Very numerous throughout the district.
- 14. Hystrix dorsata, Linn.-This species is believed to occur in the district.
- 15. Lepus Americana, Erxlebein.-Common.

ORDER RUMINANTIA.

- 16. Cervus alces, Linn.—This animal seems to be tolerably numerous above Hamilton's Farm, but none were seen in the district we passed through.
- Virginianus, Gmel.—Tracks of this deer were frequently met with, and two were reported to have been seen near sixteen Island Lake.
- " tarandus, Linn.—One was shot on Hamilton's Farm while we were camped there. Traces of them were observed on Trembling Mountain.

Besides the animals above enumerated, I may mention the racoon, *Procyon* lotor, said by the Indians to occur in the district; a wild cat, *Lynx Canadensis*, is supposed to have been heard in the township of Montcalm; a flying squirrel, *Pteromys volucella?* is said to occur, and near the Lake of Three Mountains I had a momentary view of a small *Arvicola*.

CLASS AVES.

ORDER RAPTORES.

 Buteo ? — A buzzard was frequently seen hovering around our camps,but I was unable to obtain a specimen.
 Pandion haliætus, Linn.—On the 21st of May I shot the female of a pair of this species which had their nest on the summit of a large dead pine tree on an island in a small lake situated in the 8th and 9th ranges of Montcalm. Sir William Logan has called this sheet of water Eagle Nest Lake. An osprey was afterwards seen on several occasions when ascending the Rouge. 3. Falco sparverius, Linn.—Sixteen Island Lake ; very numerous on Hamilton's Farm in August, and last seen on the 7th of

	October.
4.	Astur palumbarius, LinnHamilton's Farm, in the end of August and beginning of September.
5.	" fuscus, GmelNear Gate Lake, May 16th; very numerous at Ha- milton's Farm in the end of August.
6	Circus cyaneus, LinnHamilton's Farm, end of August and in September.
	Syrnium nebulosum, Linn.—Observed near Trembling Lake.
0.	Otus brachyotus, Linn.—I saw a specimen of this species which had just been shot on Hamilton's Farm, and was informed that it is not uncommon there after harvest.
9.	Bubo Virginianus, GmelNumerous throughout the district.
10	Order Insessores.
	Chordeiles Virginianus, Briss.—A single bird seen at Hamilton's Farm in August.
[1.	Chætura pelasgia, Linn.—Common throughout the district. They were last seen by me at Hamilton's Farm on the 25th of August.
12.	Hirundo purpurea, Linn.—Common at Grenville Village, May 13th, but not afterwards met with.
13.	" bicolor, VieillTownships of Grenville and Montcalm, middle and latter part of May. Noticed near Hamil- ton's Farm about the middle of August.
14.	•
	Farm, August 21st.
15.	" rustica, Linn.—Common in Grenville and Harrington, May 14th and 15th; Wentworth, June 4th; Hamil- ton's Farm, July 15th to the middle of
	August.
16.	Muscicapa tyrannus, Linn.—Bevan's Lake; near the Indian Village at the Devil's rapids on the Rouge; about Hamil- ton's Farm.
17.	
	ton's Farm, August 25th.
18.	Sylvicola coronata, Lath.—Abont Sixteen Island and Eagle Nest Lakes; Hamilton's Farm; Trembling Lake. Very
	numerous from May 19th till September 9th.
19.	
20.	
	Nest Lakes, May 22nd and 24th, in company with the last two species.
21	n n
41.	and about Hamilton's Farm August 23rd and
	25th.

- 23. " maculosa? Lath .- Mouth of Devil's River, July 20th.
- 24. Certhia familiaris, Linn .--- Throughout the whole district.
- 25. Troglodytes hyemalis, Vieill.—Seen occasionally at numerous localities up to September 26th.
- 26. Parus atricapillus, Linn.—First observed, August 17th, when we were camped about a mile below Hamilton's Farm,

occasionally seen till the end of September.

- 27. Regulus satrapa, Lich.—First observed August 28th, at Hamilton's Farm 28. Sialia Wilsoni, Swains.—Grenville, October 14th.
- 29. Turdus migratorius, Linn .- Throughout the district up to October 15th.
- *mustilinus*, Gmel.—Not uncommon throughout the district up to the end of September.
- 31. Sciurus aurocapillus, Lath .- Very numerous throughout the district.
- 32. Alauda alpestris, Linn.-Hamilton's Farm, end of September.
- 33. Emberiza socialis, Wils .- About all clearings visited, up to October 18th.
- 34. Niphea hyemalis, Linn.-Throughout the district.
- 35. Fringilla melodia, Wils .- About clearings throughout the district.
- " Pennsylvanica, Lath.—Very common in the woods throughout the district.
- Erythrospiza purpurea, Gmel.—Balsam Lake, June 14th; Hamilton's Farm July 15th.
- Coccoborus ludovicianus, Linn,—Clearings about Gate Lake, May 16th and 17th.
- 39. Agleaius Phæniceus, Linn.—Grenville; Sugar-bush or RoundLake; Bevan's Lake; near Hamilton's Farm.
- 40. Icterus Baltimorus, Linn.—Said to have been heard singing at Balsam Lake, June 14th.
- 41. Quiscalus versicolor, Vieill.-Grenville, May 14th.
- 42. Corvus Americanus, Aud .- Common throughout the district.
- 43. Garrulus cristatus, Linn.—Abundant everywhere. They were seen in flocks of thirty or forty at Hamilton's Farm.
- 44. " Canadensis, Linn.-Abundant throughout the district.
- 45. Vireo olivaceus, Linn.—Common throughout the district, up to August 25th.
- 46. Bombycilla Carolinensis, Briss .- Observed only about clearings.
- 47. Sitta Canadensis, Linn.—Throughout the district, from May 26th till September 20th.
- 43. Trochilus colubris, Linn.—Occasionally seen from May 27th till August 12th.
- 49. Alcedo alcyon, Linn.-Very abundant the whole way up the Rouge till October 11th. Rarely seen on the lakes.
- 50. Picus pileatus, Linn.—One shot on Sixteen Island Lake, May 27th, and another seen on the Rouge, August 8th.
- 51. " villosus, Linn.-Grenville, Harrington and Wentworth.
- 52. " pubescens, Linn.—Throughout the district.
- varius, Linn.—Sixteen Island Lake, May 27th; Trembling Lake, September 13th.

54.	Picus articus, Swains One specimen observed in Harrington, October 15th.
55.	" auratus, Linn.—Hamilton's Farm, end of August and beginning of September.
56.	Coccyzus erythropthalamus, Wils Sugar-bush Lake, June 25th; Indian
11	Village on the Rouge, July 16th.
	ORDER RASORDS.
57.	Ectopistes migratoria, Linn.—Throughout the district, from spring till the beginning of October. Not common.
	Tetrao umbellus, LinnAbundant throughout the district.
	OEDEE GRALLATORES.
59.	Fulica Americana, Gmel.—A pair scen, September 14th, in a small lake near the Lake of Three Mountains.
60.	Totanus macularius, WilsCommon all along the Rouge and in the nume- rous lakes of the district.
61.	
	13th.
62.	" vociferus, Wils One specimen seen on Trembling Lake, Sept- ember 11th.
63.	Microptera Americana, Aud.—Said to have been heard in the swamps about Hamilton's Farm, September 2nd.
4.	Ardea nycticorax, LinnA pair seen flying over head, when we were camped near Gate Lake, May 17th.
65.	" lentiginosa? Swains Bevan's Lake during July.
	ORDER NATORES.
66.	Anas obscura, GmelSixteen Island and Bevan's Lakes; Rouge, and the small lakes on either side of it.
67.	" sponsa, Linn.—One seen on Bevin's Lake, October 16th.
68.	" discors, Linn One observed on Trembling Lake, September 11th.
69.	Fuligula marila? Linn Sixteen Island Lake, May 20th.
70.	" clangula, Linn.—Sixteen Island Lake in May; Devil's River, July 20th.
71.	Mergus servator, LinnRouge, and almost every lake we visited.
72.	" cucullatus? Linn.—Lake of Three Mountains, September 23rd and 25th.
73.	Larus argentatus, Brunnich.—A large gull, supposed to be of this species, was frequently seen at the end of May, on Sixteen Island Lake.
74.	Colymbus glacialis, LinnSeen in almost every lake visited by us.
75.	Podiceps Carolinensis? LathI observed a grebe on Balsam Lake, June 14th, which appeared to be of this species.
π	the side hunting Delicter at the I the I had a sector

The rice bunting, *Dolichonyx orizivora*, and the red-headed woodpecker, *Picus erythrocephalus*, were observed about Point Fortune on the Ottawa, but were not met within the woods.

CLASS REPTILIA.

ORDER CHELONIA.

 Chelydra serpentina, Schw.—Emysaurus serpentina, Linn.—I was given a shell of this species by G. W. Albright, Esq., P. L. S., who obtained it on the Devil's River. The carapace is one foot long and nine inches broad.

 Glyptemys insculpta, Agassiz.—Emys insculpta, Leconte.—I was shown the shell of a specimen of this species, which had been obtained on a small sandy island in the Rouge in Arundel, and I also obtained a fragment of a shell at the mouth of the Devil's River.

OBDER OPHIDIA.

 Eutainia sirtalis, Baird & Girard. Tropidonotus sirtalis, Holbrook.—Abundant in the Townships of Grenville, Harrington, De Salaberry, and at Hamilton's Farm.

No other Ophidian reptile was seen, but reports of a water snake, said to inhabit the lakes, came to my knowledge.

ORDER BATRACHIA.

- 4. Rana Catesbiana, Shaw.—R. pipiens, Holbrook.—Abounds in every lake and pond throughout the district.
- iii nigricans, Agassiz.—Abundant at Balsam, Sixteem Island and Sugarbush Lakes in May and June.
- 6. " pipiens, Gmel. R. halecina, Holbrook et aliorum.—Abundant in Sugar-bush Lake in June.
- 7. Hyla versicolor? Leconte.—Said to have been heard about Sixteen Island Lake.
- 8. Bufo Americana, Leconte .- Common throughout the district.
- 9. Plethodon erythronota, Green.—Abundant in the townships of Wentworth and Montcalm in May.
- 10. Spelerpes bilineata, Green.-Township of Montcalm.
- 11. Triton? (undetermined).—One specimen taken in Sixteen Island Lake June 2nd.
 - A "lizard" was reported as inhabiting a small stream crossing the portage between Gut and Gate Lakes.

CLASS PISCES.

OBDER ACANTHOPTERI.

 Perca flavescens, Cuvier.—Numerous in Sugar-bush, Bevan's and Bark Lakes, Montcalm; in a small lake on lot 11, range 3, of the same township, and also in a lake about three miles east of Hamilton's Farm.

ORDER MALACOPTERI.

2. Pimelodus canosus, Richardson.-Very abundant in the same lakes (with the exception of the last) as the perch.

3. Esox boreus ? Agassiz.—The specimen preserved, was caught in the small lake on lot 11, range 3, Montcalm, and agrees very well with the description of E. boreus in Agassiz's "Lake Superior," p. 317, with the exception, that the lateral line is very indistinct, instead of being "very distinct." Pike were numerous in the same lakes as the cat-fish and perch and in the Rouge as far up as we ascended.

4. Salmo fontinalis, Mitchill.—Abounds in nearly all the lakes and streams in the district. In those lakes where cat-fish, pike and perch occur, no trout were caught.

- Salmo.—A species of trout, which I have been unable to determine, was found in Sixteen Island, Trembling and Three Mountain Lakes.
- 6. Coregonus.—I saw several specimens of a Coregonus which had just been taken in Bevan's Lake, October 15th, but was unable to preserve a specimen
- 7. Catastomus.—Two species of "sucker" were said to have been taken in Sixteen Island Lake whilst I was absent, and were spoken of as the "mullet" and "black sucker."
- 8. Leuciscus.—A large fish known as the carp, usually about seventeen inches in length and about two pounds in weight, was abundant in all the lakes and in the Rouge and Devil's Rivers. On the sides, the scales have a beantiful bronze or golden lustre, and the basal half and margin of each is black. The fin-rays are as follows: Br. 3, D. 9. C. 20. V. 8. P. 16. This fish may be Cyprinus Corporalis, Mitchill, but does not agree satisfactorily, with any species I have seen described.
- *pulchellus*, Storer.—This was the most abundant fish in all the lakes and rivers throughout the district.
- 10. Leuciscus frontalis, Agassiz.—Abundant in streams flowing into the small lake on lot 11, range 3, Montcalm. The specimens collected agree exactly with the figure and description of this species in Agassiz's "Lake Superior," except that instead of fourteen, they have sixteen rays in their pectoral fins.
- "? —A small species which I cannot find described, though evidently very distinct, was common in the same stream with the last.

All the lakes swarmed with the young of various *Leucisci*, which are called dace and chub. Several species besides those above mentioned were met with in Trembling and Three Mountain Lakes, but I had no means of preserving specimens.

ARTICULATA.

CLASS INSECTA.

ORDER COLEOPTERA.

Besides the 114 species of Coleoptera enumerated in the following catalogue, many others were collected, but were unfortunately lost by the accidental fracture of the bottle which contained them.

I have added a list of 34 species, not observed in this district by myself, but brought by Mr. Robert Bell from the Augmentation of Grenville on the north, and the neighbourhood of L'Original on the south bank of the Ottawa.

1. Cicindela longilabris, Say.—Hamilton's Farm on the River Rouge, 2nd September.

- 2. " vulgaris, Say.--Very abundant on sand-banks, River Rouge, August.
 - "Baltimorensis, Herbst. (repanda, Say.)—Common on sand-banks, River Rouge, July and August.
 - 4. Lebia viridis? Say.—Huckleberry Rapids, River Rouge, DeSalaberry, 30th July.
 - 5. Patrobus longicornis, Say.—Sixteen-Island Lake, &c., Montcalm, May and June.
 - 6. Platynus sinuatus, Dej.—Under dead logs, Sixteen Island Lake, &c., township of Montcalm, May and June.
 - 7. " retractus, Lec.-With the last species.
 - 8. " obsoletus, Say .- With the last two species.
 - 9. Pacilus lucublandus, Say.--Under stones near the town of Grenville, 13th May.
- Pterostichus fastidatus, Dej.—Under bark of decaying logs, Sixteen Island Lake, Montcalm, end of May; Lake of Three Mountains, end of September.
- 11. " patruelis, Dej.-River Rouge.
- 12. " caudicalis, Say.-Under stones near Grenville, 13th May.
- 13. " orinomum, Leach (vitresis, Esch.)-Township of Montcalm, June.
- " Luczotii, Dej. (var. præc?)—Sixteen Island Lake, Montcalm, May and June.
- 15. Lophoglossus scrutator, Lec.-Under stones near Grenville, 13th May.
- 16. Rembus major, Lec.-- " " " " "
- 17. Chlænius impunctifrons, Sag " " " "
- Cychrus (Sphæroderus) Brevoorti, Lec.--Under dead logs, Bevan's Lake, Montcalm, 4th July.
- Notiophilus punctatus, Lec.—On rocks, Huckleberry Rapids, River Rouge, DeSalaberry, 27th July.
- 20 Bembidium impressum, Fabr .-- On wet sand, River Rouge, 13th August.
- 21. " punctatostriatum, Say Very abundant on wet sand, River Rouge July and August.

22. Bembidium patruelis, Dej .- Abundant on wet sand, River Rouge, 13th Aug 23. " lucidum, Lec .-- Under stones near Grenville, 13th May. 24. Agabus striatus? Say .- In Sixteen Island Lake, Montcalm, end of May. 25. Coptotomus interrogatus, Fabr .- In Sugar-bush Lake, Montcalm, 23rd June. 26. Hydroporus proximus, Aubé .- With the last species. 27. Haliplus immaculaticollis, Harris .- With the last two species. 28. " cribarius, Lec .- Very abundant in Sugar-bush Lake, Montcalm, 23rd June. 29. Gyrinus (several species not determined)-In various Lakes. 30. Dineutes (not named)---Very abundant, Sugar-bush Lake, Montcalm, 23rd June. 31. Philhydrus cinctus, Say .- In a small stream crossing the portage between Gate and Gut Lakes, Wentworth, and in Sugar-bush Lake, Montcalm. 32. Necrophorus lunatus, Lec .-- Huckleberry Rapids, River Rouge, De Salaberry, 27th July. 33. pygmæus, Kirby .- Township of Montcalm, 20th June. 34. Silpha marginata, Fabr.-Abundant under putrid fish, Sixteen Island Lake, Montcalm, 1st June. 35. Homalota (not determined)-Township of Montcalm, June. u' 36. Tachyporus (not determined) 66 37. Tachinus fumipennis, Say .-- In bear's dung, Chain Lake, Montcalm, 17th June. conformis, Dej .- Township of Montcalm, June. 38. 39. Philonthus cyanipennis, Fabr .-- In a fungus on a rotten tree, River Rouge, 13th August. (not determined)-Under stones near Grenville, 13th May. 46 40. 41. Stenus (not determined)-Numerous on wet sand, River Rouge, Arundel, July. (not determined)-Numerous on wet sand, River Rouge, near 42. Hamilton's Farm, 13th August. 43. Oxytelus Pennsylvanicus, Er.-Common in our tents throughout the district. 44. Anthobium dimidiatum, Mels.-Township of Montcalm, June. 45. Platysoma parallelum, Say.-" " " 66 " 46. Carpophilus niger, Er.-" ٤٢ 47. Epurcea, (not determined) " 48. Cucujus clavipes, Oliv .- One specimen taken as it pitched on the mane of a horse, Township of Harrington, 15th May. 49. Pediacus planus, Lec .- Very abundant in the tents, Huckleberry Rapids, end of July. 50. Dermestes lardarius, Linn .- Observed about the provisions, Sixteen Island Lake, Montcalm. 51. Anthrenus castaneæ, Mels.-Township of Montcalm, June. 52. Platycerus depressus, Lec .- Near Huckleberry Rapids, River Rouge, De-Salaberry, July. 53. Onthophagus Hecate, Pz .- Near Huckleberry Rapids, River Rouge, De-Salaberry, 2nd August. 54. Geotrupes Egeriei, Germ. (microphagus, Say.) Woods near Hamilton's Farm, 31st August.

55. Aphodius fimetarius, Fabr.-Abundant in cow-dung, Hamilton's Farm, August. 56. Dichelonycha subvittata, Lec .-- Abundant throughout the district, June to August. 57. Osmoderma scabra, Beauv.-River Rouge, July and August. 58. Nichius piger, Fabr .-- On blossoms of Viburnum opulus, Sugar-bush Lake, and on white clover blossoms, and bleeding stumps of yellow birch, Bevan's Lake, Montcalm, end of June and beginning of July. 59. Ancylocheira maculiventris, Say .- Near Silver Mountain, River Rouge, 12th August. 60. Cryptohypnus silaceipes, Germ .-- Under stones near Grenville, 13th May. 61. Dolopius fucosus, Lec .-- Township of Montcalm, June. " stabilis, Lec .---٤ ٢ " 65 62. 63. Corymbites triundulatus, Randall .- Township of Montcalm, end of May. 64. Pyractomena angulata, Say .-- Common, Sugar-bush Lake, Montcalm, 23d to 26th June. 65. Ellychnia corrusca, Linn .--- Under stones near Grenville, 13th May. 66. " lacustris, Lec .- Abundant in the woods of Harrington, middle of May; Hamilton's Farm, and Lake of Three Mountains, August and September. 67. Digrapha terminatis, Say .- Bevan's Lake, 29th June, and 5th July, and Hamilton's Farm, 31st August. 68. Eros coccinatus, Say .-- Sixteen-Island Lake, &c., Montcalm, end of May. molis, Lec .-- Huckleberry Rapids, River Rouge, DeSalaberry, 69. 11 2nd August. 70. Podabrus modestus, Say .-- About clearings, Bevan's Lake, Montcalm, 2nd July. " 28 71. Telephorus rotundicollis, Say .- Abundant " 72. 66 carolinus, Fabr .---٤ĥ " " 66 66 fraxini, Say .- Township of Montcalm, June. 73. 74. Anobium foveatum, Kirby.-Abundant in a rotten tree, Bevan's Lake, 4th July. 75. Cis (not determined)-Township of Montcalm, June. 66 76. Pedilus collaris, Say .--44 66 77. Mordella nigricans, Mels.- " 66 z 78. Meloe rugipennis, Lec .- Hamilton's Farm, 31st August, and Grenville, 14th October. 79. Cistela (not determined)-Very abundant on leaves of bass-wood, Sugarbush Lake, Montcalm, 26th June. 80. ٤٤ (not determined)-River Rouge. 81. Nyctobates (not determined)-Under logs on grass-land, Hamilton's Farm August. 82. Upis reticulatus, Say .-- (ceramboides, Linn.)-With the last species. 83. Bolitophagus cornutus, Pz.-Larvæ and Pupa in a boletus, Huckleberry Rapids, DeSalaberry, 3rd August. 84. Apion (not determined)-Township of Montcalm. 85. Sitona lepidus, Sch.-Near Hamilton's Farm.

86. Hylobius (near pineti)-Sixteen Island Lake, 1st June.
87. " pales, HerbstTownship of Montcalm, June.
88. Tomicus (not named) """""
89. Saperda tridentata, OlivBase of Silver Mountain, Rouge, 10th Aug.
91. " scutellatus, SayNumerous, Bevan's Lake, 7th July; and
abundant the whole way up the Rouge, to
the end of August.
92. Encyclops cœruleus, Say One specimen taken on blossoms of Viburnum
opulus, Sugar-bush Lake, Montcalm, 26th
June.
93. Acmaops proteus, Kirby Township of Montcalm, June.
94. Evodinus monticola, RandallSixteen-Island Lake, 30th May; and abun-
dant on blossoms of Viburnum opulus, Sugar-
bush Lake, end of June.
95. Leptura Canadensis, OlivAbundant on blossoms of Spirae salicifolia,
River Rouge, July and August.
 96 "vittata, Oliv.—Near Huckle-berry Rapids, DeSalaberry, 15th July. 97 "pubera, Say.—Abundant on blossoms of Viburnum opulus, Sugar-
97 " pubera, Say.—Abundant on blossoms of Viburnum opulus, Sugar- bush Lake, Montcalm, 25th June.
and the second s
 98. "proxima, Say.—Near Huckleberry Rapids, DeSalaberry, 26th uy. 99. "mutabilis, Lec.—On blossoms of Viburnum opulus, Sugar-bush Lake,
end of June.
100. Donacia palmata, Oliv In blossoms of Nuphar advena, (Yellow Water-
lily), Sugar-bush Lake, end of June.
101. " subtilis, Kunze In a small Lake near Lake of Three Moun-
tains, 14th September.
102. " pusilla, Say Sugar-bush Lake, Montcalm, end of June.
103. " Aavipes, Kirby " " "
104. Syneta tripla, Say Township of Montcalm.
105. Chrysomela scalaris, LecAbundant on alders throughout the district,
from the end of June to the end of September
106. " spira, SayVery abundant, Sugar-bush Lake, 25th June.
107. " interrupta, FabrAbundant on alders, Sixteen-Island and
Sugar-bush Lakes, Montcalm, May and June.
Larva abundant on alder leaves, June 25.
108. Chrysomela Vitellinæ, LinnAbundant on oak and poplar leaves, Sixteen
Island and Sugar-bush Lakes, May and June. 109. Systema pontalis, Fabr.—Township of Montcalm, June.
110. Phyllobrotica decorata, Say. (Oliviéri, Kirby.)-Very abundant on Scu-
tellaria galericulata and laterifolia, River
Rouge, July and August.
111. Adoxus vitis, Fabr.—Amongst dead leaves, Gate Lake, Wentworth, 16th
May.
112. Chrysochus auratus, FabrAbundant on Apocynum androsæmifolium and
cannabinum, Bevan's Lake, Huckleberry Ra-
pids, &c., July.
113. Galleruca sagittariæ, KirbyTownship of Montcalm, June.
114. Coccinella picta, Randall.— "" "

The following are the thirty-four species of Coleoptera from L'Orignal and the Augmentation of Grenville, collected by Mr. R. Bell.

Cymindis reflexa, Lec. Calathus gregarius, Say. Platynus capripennis, Say. Pterostichus erythropus, Dej. 66 adjunctus, Lec. Amara angustata, Say. 11 impuncticollis, Say. Anisodactylus Baltimorensis, Say. " Harrisii, Lec. ٤6 rusticus, Say. Harpalus Pennsylvanicus, Geer. " herbivagus, Say. Chlanius sericeus, Forst. 66 tricolor, Dej. Acilius fraternus, Harris. Silpha Surinamensis, Latr. Pæderus littorarius, Grav.

Hister perplexus ? Lec. Ips quadrisignatus, Say. Cytilus varius, Fabr. Lachnosterna fusca, Frolich. Osmoderma eremicola, Knoch. Photuris Pennsylvanica, Geer. Trichodes Nuttalii, Kirby. Thanasimus dubius, Fabr. Tenebris molitor, Linn. Ipthinus Pennsylvanicus, Geer. Orthosoma unicolor, Drury. Saperda vestita, Say. Chelymorpha cribaria, Fabr. Haltica collaris, Fabr. Chrysomela trimaculata, Fabr. Helodes trivittata, Say. Hippodamia 13-punctata, Linn.

ORDER LEPIDOPTERA.

With the exception of the *Rhopalocera*, the greater portion of the *Lepidop*tera collected are still undetermined. Some of the *Heterocera* enumerated below were named for me at the British Museum by Mr. Francis Walker, towhom I am much indebted.

Rhopalocera.

115.	Papilio turnus, Linn	Abundant throughout the whole district, from
		May 30th till the end of July.
116.	" asterias, Fa	bA large black butterfly, seen on the 17th June at
	,	Balsam Lake I supposed to be of this species.
117.	Colias philodice, God	t Grenville Village, June 5th; along the Rouge
		from the 30th June till the middle of Septem-
		ber; again seen at Grenville, October 14th
		and 18th.
118.	Pieris oleracea, Harris	sAbundant throughout the whole district, from
		the middle of May till the end of June. A
		few seen at Hamilton's Farm, end of August.
119.	Danais Archippus, Fa	bA single specimen seen flying across the Rouge
		a little above Silver Mountain on the 12th of
		July.
120.	Debis Portlandia, Bo	isd First seen on the 2nd July, at Bevan's Lake.
	,	Abundant in the woods along the Rouge as
		far as Silver Mountain till the first week in
		August. As this is generally supposed to
		be a southern species, it is not a little re-
		markable that it should be so abundant to
		the north of the Ottawa.

121. Hipparch	ia nephele ? Kirby Abundant amongst grass on Hamilton's
	Farm, from the 22nd August till the begin-
DANS 3 - BRALLAND	ning of September.
i singul the	Arthemis, DruryVery abundant throughout the district, from the 26th June till the end of July, a few
一种 一方的 医骨盆膜的	lingering till the middle of August.
123. Cynthia	cardui, LinnOne specimen met with at Hamilton's Farm on the 21st August.
124. Vanessa	Atalanta, Linn I observed a butterfly which appeared to be of
一日 うちちち うちち	this species, at Sugar-bush Lake on the 24th of June.
125. "	Antiopa, LinnGrenville Village, May 13th; a few seen in the
when the share	township of Montcalm in June and near
les tenet test	Silver Mountain on the 12th of August.
126. "	Milberti, Godt., furcillata, Say Grenville Village, May 14th ;
Simult be bland	Rouge, July 10th, and occasionally seen at
	Hamilton's Farm, up to the 31st August.
	J. album, Boisd Common throughout the district, from May
	19th till the end of September. One observed
	near Grenville on the 18th of October.
	Fogne, FabAbundant everywhere, from the 14th May till the
A data taken ta	middle of September.
129. "	C. album, Godt I took several specimens of a Grapta along
	the Rouge which I believe to be of this
100 0	species.
130. Argynnis	Dalphnie (?), CramerFirst seen, July 2nd, and last, Sep- tember 12th. Abundant. I am of opinion
	Aphrodite, Fab. and A. Cybele, Fab., as the
	same species. There are at least three closely
	allied species of Argynnis inhabiting Canada,
	but nothing short of breeding each from the
	larva will satisfactorily separate them. My
	specimens are all too small for \mathcal{A} . Cybele,
101 Augumente	Fab.
	Myrina, CramerFrom June 5th till August 31st. Common.
132. "	Bellona, Fab.—The only specimen met with, was captured in Arundel on the 30th June.
133. Melitæa	Thurse, Oramer Sugar-bush Lake, June 29th; Bevan's Lake,
	July 2nd ; Devil's River, July 14th.
134. Thecla (?)-I observed a large Thecla at Huckleberry Rapids, July 30th,
	but did not succeed in capturing it.
135. Lycæna A	Imericana, Harris Numerous on grass land at Hamilton's Farm,
	from the 21st to the 31st August.
136. Polyomm	atus pseudargiolus, BoisdNumerous in Grenville and about
	Sixteen Island Lake in May. Worn speci-
	mens were seen about Bevan's Lake as late as
	the 2nd of July.

137. Pamphila.—One specimen of a species resembling P. paniscus of Europe was captured near Bevan's Lake, July 2nd. Specimens of a dingy grey species and of two or three other Pamphilas were taken in various localities in June, July and August. I cannot find descriptions of any of these and some of them are probably new.

Heterocera.

138. Sphinz.—Two species of Sphinz were captured in July, in Arundel and DeSalaberry, allied to S. Kalmiz, A. & S. and S. gordius, Cramer, but not agreeing satisfactorily with Dr. Harris's descriptions of these species given in the Amer. Jour. Sci. Vol. 28.

139. Smerinthus.—Two larvæ belonging to this genus were obtained at Hamilton's Farm on the 3rd and 4th September, of which the following are descriptions.—No. 1. Pale green, whitish on the back, with oblique stripes of white and dark green on the sides.—No. 2. Green, with oblique tuber-culated stripes on the sides and two tubercles on each of the second and third segments.

140. Trochilium.—On the 25th June, at Sugar-bush Lake, I captured a beautiful and apparently undescribed species of Trochilium, sitting on the blossoms of Viburnum opulus (high-bush cranberry). The anal tuft is deep orange; antennæ black; expansion of the wings 11 lines; length of the body 5 lines.

141. Ctenucha Latreillana, Kirby .- One specimen taken in Arundel, July 16th.

- 142. Crocota brevicornis, Walker.—Township of DeSalaberry; Hamilton's Farm, July and August.
- 143. Medaria mendica, Walker.-Near Bevan's Lake, July.
- 144. Arctia Parthenos, Harris.—I took a fine Arctia on the Devil's River, July 19th, agreeing in every respect with the description and figure of this species in Agassiz's "Lake Superior," with the exception that it has five, instead of three cream-coloured spots on the costal edge of the anterior wings.
- 145. Hypercompa Lecontei, Boisd.—Montcalm, Arundel and DeSalaberry, during the month of July.
- 146. Halesidota aunulifascia, Walker.—Coccoons, apparently of this species, were found near Sixteen Island Lake, May 22nd.
- 147. Orgyia leucostigma, A. & S.-Hamilton's Farm, end of August and beginning of September.

148. Telea Polyphemus, Hübner .- Throughout the whole district.

149. Thyatira scripta, Gosse.-Montcalm and Arundel, end of June and beginning of July.

150. " cymotaphoroides, Guén.-Montcalm and Arundel, June and July; Trembling Lake, September 7th.

151. Graphiphora C. nigrum, Linn.—One specimen taken in DeSalaberry, July 24th, and another at Hamilton's Farm, August

28th.

"Dahlii, Höbner.—One specimen taken in Wentworth, May 17th.
 Euplexia lucipara, Linn.—Common in Montcalm in June.

154. Plusia mortuorum .- Hamilton's Farm, end of August.

155. Angerona crocataria, Fab .-- Common in Arundel and Montcalm in July.

156. Sicya solfaiarina, Guén .- Not uncommon in DeSalaberry, end of July.

157. Ellopia æqualiaria.-Montcalm, June.

158. Nematocampa filamentaria, Guén.-DeSalaberry, July 22nd.

159. Endropia tigrinaria, Guén.-Very abundant in Montcalm at the end of June.

160. Melanippe Goihicata, Guén.—Extremely numerous in Montcalm during the month of June.

161. Scotosia undulata, Linn.—Common in Montcalm, end of June and beginning of July.

162. Pyralis n. sp?-DeSalaberry, June 27th. Mr. Walker supposes this to

be a new species, and the following is a description of it.—Anterior wings, dull pink, crossed by two black tranverse lines, the first of which, situated near the base, is straight and has a yellow spot on its inner side, occupying the angle which it forms with the costa; the second, situated beyond the middle, is bent, forming an obtuse angle before it reaches the costal margin, where it has a yellow crescent-shaped spot on the outer side. Posterior wings, dusky-white at the base, with a broad, pale black, sub-marginal band and crossed by two black tranverse lines. Expansion of the wings 12¹/₂ lines; length of body 4 lines.

163. Bleptina surrectalis, Guén .- DeSalaberry, August 4th.

164. Anania octomaculata, Linn .-- One specimen taken in Montcalm, July 2nd.

165. Hydrocampa.—A species of Hydrocampa was abundant near Hamilton's Farm, August 15th.

166. Batys verticalis, Linn.-DeSalaberry, not uncommon about the first of August.

167. Eubulea.—A small species apparently closely allied to the European E. sambercalis, Schiff., was very numerous on the blossoms of the rasberry (Rubus strigosus), near Bevan's Lake, at the begining of July.

B

168. Tortrux.-Several pupe of a Tortrix, which I collected on the Devil's River, produced the perfect insect, but I have been unable to determine either this or any other of my Micro-Lepidoptera.

MOLLUSCA.

CLASS GASTEROPODA.

ORDER PULMONIFERA.

1.	Tebenn	ophorus Carolinensis, BoscThroughout the district.
2.	Succin	ea obliqua, Say.—Occurred plentifully at Hamilton's Farm, and sparingly in wild parts of the district.
3.	Helix o	Bolabris, Say Wentworth ; Montcalm ; Harrington.
4.	66	exoleta, BinneyWentworth ; DeSalaberry ; Harrington.
5.	66	monodon, RackettArundel; Hamilton's Farm; near the Lake of
		Three Mountains.
6.	"	concava, Say Wentworth ; Montcalm ; Arundel. Abundant.
7.	66	pulchella, Müller Under stones at Carillon, but not elsewhere
		met with.
8.	13	Sayii, Binney.—Near Doran's Lake, Grenville.
9.	66	labyrinthica, Say Wentworth ; Montcalm ; Arundel. Common.
10.	22	alternata, Say Abundant throughout the district.
11.	66	striatella, AnthonyVery abundant throughout the district.
12.	23	arborea, SayPlentiful throughout the district.
13.	22	chersina, Say. " " " "
14.	66	lineata, Say.—Abundant throughout the district.
15.	Bulim	us marginatus, SaySugar-bush Lake and near Gate Lake.
16.	Achati	na lubrica, Müll.—Bevan's and Gate Lakes. Common.
17.	Vertig	o Gouldii, BinneySixteen Island Lake.
18.	Pupa	(undetermined)With the last species.
19.	Carycl	ium exiguum, Say One specimen found near Sixteen Island Lake.
		(Fresh Water.)

- 20. Physa heterostropha, Say .--- Sugar-bush Lake, and near Grenville Village.
- 21. " aurea, Lea.-Small Lake near Hamilton's Farm.
- 22. Physa elliptica, Les.-In a small lake one mile west of the Indian Village in Arundel.
- 23. " elongata, Say .- Near Grenville Village.
- 24. Limnæa exigua, Lea. (young) .- In a small lake near Hamilton's Farm.
- 25. " galbanus, Say.—Abundant in shell marl from the bottom of Eagle Nest Lake.
- 26. " " plicata, Lea.---Sugar-bush Lake. Abundant.
- 27. " reflexa, Say .- Near Grenville Village.
- 28. " umbilicata, Say .--- With the last species.

29. Planorbis truvolvis, Say .- In the small lake one mile west of the Indian Village in Arundel.

30. " bicarinatus, Say.—Eagle Nest Lake and a small lake neaar Hmilton's Farm.

31. " campanulatus, Say.—Near Grenville Village and in numerous lakes throughout the district.

32. " parvus, Say.—In shell marl in Eagle Nest Lake, and living in the lake one mile west of the IndianVillage, Devil's Rapids, and in the lakes near Hamilton's Farm.

33. " deflectus, Say .-- Sixteen Island and Sugar-bush Lakes.

ORDER PROSOBRANCHIATA.

34. Paludina decisa, Say.—Very abundant the whole way up the Rouge and its tributary the Devil's River. Those collected are of a reddish-brown color, very unlike the light green of specimens from L'Orignal, opposite the mouth of the Rouge, and from the St. Lawrence near Montrsal.
35. Valvata tricarinata, Say.—A few specimens found in shell marl from the

bottom of Eagle Nest Lake.

CLASS LAMELLIBRANCHIATA.

1. Unio complanatus, Lea. This was the only species of Unio met with. It

inhabits nearly every lake in the district and was abundant in the Rouge as far as we ascended it. It was extraordinarily abundant in the shallow stream by which the waters of Bevan's and Bark Lakes are discharged into the Rouge; in fact they were crowded together as closely as they could lie, in the same manner as a bed of

Margaritana rugosa, Barnes.
 Margaritana rugosa, Barnes.
 Margaritana rugosa, Barnes.
 Margaritana rugosa, Barnes.
 Month small lake west of Balsam Lake, lot 11, range 3. Montcalm.

3. Anodonta cygnea (?), Linn.—This species was found in almost every lake we visited.

4. Anodonta edentula, Say.—One specimen obtained from the lake on los 11, range 3, Montcalm.

5. " fragilis, Linn.-Sixteen Island, Eagle Nest, and Bevan's Lakes.

6. " Footiana, Lea.-With the last species.

 Cyclas similis, Say.—Sixteen Island and Sugar-bush Lakes; lake ene mile west of the Indian Village; in shell, marl in Eagle Nest Lake.

- 8. " partumeia (?), Say:—Ponds near Eagle Nest Lake; Sugar-bush Lake; small lake near Hamilton's Farm.
- 9. " dubia (?), Say.—In shell marl, Eagle Nest Lake; living in the small lake near Hamilton's Farm.

Catalogue of Animals and Plants collected and observed, on the south-east side of the St. Lawrence from Quebec to Gaspé, and in the Counties of Rimouski, Gaspé and Bonaventure. By Mr. Robert Bell Jr., Assistant to Mr. James Richardson, Geological Explorer under Sir W. E. Logan, in 1858.

VERTEBRATA.

CLASS MAMMALIA.

ORDER CHEIROPTERA.

1. Vespertilio subulatus, Say .- Restigouche, Matapedia and Patapedia Rivers.

ORDER INSECTIVORA.

2. Sorex Forsteri, Richardson .- Counties of Rimouski and Bonaventure.

ORDER CAENIVORA.

	3.	Ursus	Americanus,	Pallas	Through	hout t	he d	listrict.
--	----	-------	-------------	--------	---------	--------	------	-----------

4.	Mustela martes, Linn	23	55
5.	" vison, Gmel	66	66
6.	" vulgaris, Linn	88	66
7.	" Canadensis, Schreber	. 66	66
8.	Mephitis Chinga, Tiedimann	66	55
9.	Lutra Canadensis, Sabine	66	6.6
10.	Canis fulvus, Desm	56	66

11. " lupus, Linn .- Said to come no farther north than the St. John River.

12. Lynx Canadensis, Linn.-Gaspé, and probably throughout the whole district.

ORDER RODENTIA.

44

٤٤

13. Castor fiber, Linn.-Throughout the district.

14. Fiber zibethicus, Cuv.,— "

15. Mus musculus, Linn .- In settled parts throughout the district.

16. Pteromys volucella, Desm.-Gasp6.

17. Tamias Lysteri, Ray.-Rimouski and Gaspé.

18. Sciurus Hudsonius, Penn.-Throughout the district

19. Hystrix dorsata, Linn.— "

20. Lepus Americanus, Erxl.- "

ORDER RUMINANTIA.

21. Cervus alces, Linn .-- Rimouski, Bonaventure and western part of Gaspé.

22. " tarandus, Linn.-Among the Shickshock Mountains.

CLASS AVES.

Determined by Mr. D'Urban.

ORDER RAPTORES.

 Haliætus leucocephalus, Linn.—Along the St. Lawrence from Green Island to Martin River, in June and July; seen on the Restigouche in August.

2. Astur fuscus, Gmel.-Capucin, August 8th.

3. Surnia funerea, Gmel.-Green Island, middle of October.

4. Syrnium nebulosum, Gmel.-Marsouin River, end of July.

ORDER INSESSORES.

 Chordeiles Virginianus, Briss.—Chat River, June 18th; [Ste. Anne, June 28th to July 17th; at the mouth of the Matapedia, August 28th.

6. Hirundo bicolor, Vieill.-Chat River, June 18th; Ste. Anne, June 30th: Martin River, July 20th.

7. " fulva, Vieill.-Metis, beginning of June.

8. " rustica, Linn.-Trois Pistoles, May 30th ; Metis, June 10th ; Long Point, June 15th.

9. " riparia, Linn.-Ste. Anne, June 28th.

10. Sylvicola coronata, Lath .- Green Island Village, May 25th.

11. Troglodytes hyemalis, Vieil1 .- Patapedia River, September 5th.

- 12. Parus atricapillus, Linn.—First seen on the Patapedia River, September 5th, and afterwards in various localities.
- 13. Regulus satrapa, Lich .--- Rivière du Loup, May 18th.
- 14. Turdus migratorius, Linn .- In settled parts, throughout the district.
- 15. Anthus Ludovisianus, Lich.-Rivèire du Loup to Rimouski, from May 10th to June 5th.

16. Alauda alpestris, Linn .-- Bimouski Village, beginning of October.

- 17. Plectrophanes nivalis, Linn .- Kamouraska, beginning of November.
- 18. Emberiza socialie, Will.---Various localities from Rivière du Loup to Cape Chat.
- 19. Niph as hyemalis, Linn .- Throughout the district.

 Carduelis tristia, Linn.---Along the coast from St. Fabien to Martin River, from May 31st to July 19th; on the Restigouche, September 2nd.

21. Fringilla Pennsylvanica, Lath.-About clearings along the whole coast.

22. Erythrospiza purpurea, Gmel.-St. Fabien, May 30th; Ste. Anne, July 18th.

- 23. Agelaius Phaniceus, Linn,-Ste. Anne, July 17th.
- 24. Quiscalus ferrugineus, Lath.—Metis River, and between Metis and Rimouski, September and October.

25. Corress Americanus, Aud.-Along the whole coast, and on the Restigouche, but not seen in inland parts.

26. Garrulus cristatus, Linn .- Lake Matapedia, August 19th.

27. " Canadensis, Linn.-Throughout the district.

 Bombycilla Carolinensis, Briss.—Metis, June 8th; Ste. Anne, in July; Marsouin River, August 2nd. 29. Trochilus colubris, Linn .- Metis, middle of August.

30. Sitta Canadensis, Linn .- Matapedia Lakes, August 19th.

31. Alcedo alcyon, Linn.—Throughout the district; observed from May 19th to the end of September.

32. Picus pileatus, Linn .- Green Island Seigniory.

33. " villosus, Linn.-Bic, Ste. Anne, Marsouin and Martin Rivers.

ORDER RASORES.

- 34. Ectopistes migratoria, Linn.—From Metis to Ste. Anne; about Lake Matapedia, and along the Restigouche, from Jane 18th to August 31st.
- 35. Tetrao umbellus, Linn.—Near Rimouski. This species was not met with in Gaspé, and is believed by the Indians not to extend so far to the north-east.

to extend so tar to the hor

36. " Canadensis, Linn.-Throughout the district.

ORDER GRALLATORES.

- 37. Strepsilas interpres, Linn.-Green Island Village, October 26th.
- 38. Tringa pusilla, Wils.—Rivière du Loup and Green Island in May; Chat and Martin Rivers in July.
- 39. Tringa (undetermined) .-- Mouth of Marsouin, August 4th.
- 40. Totanus solitarius, Wils .- Matapedia and Restigouche Rivers in August.
- 41. " vociferus, Wils .- Rivière du Loup, May 20th.
- 42. Scolopax Noveboracensis, Gmel.-Green Island, May 25th.
- 43. Ardea nycticorax, Linn.—Dalhousie, N. B., August 25th; Patapedia River, September 9th; Metis Lake, October 1st.

ORDER NATORES.

- 44. Anser Canadensis, Linn.—Rimouski, beginning of June; Cape Chat, June 17th, and near Green Island and Cacouna in the end of October.
- 45. " leucopsis, Bechst .-- Rimouski and Green Island in October.
- 46. Fuligula fusca, Linn .- Coast of Rimouski and Gaspé in June and July.
- 47. " perspicillata, Linn.-Green Island and various localities further down.
- " clangula, Linn.—Bic and Green Island in October, and Metis Lakes, September 18th.
- 49 " histrionica, Linn.—Ste. Anne River in July; Restigouche in August, and Patapedia in the beginning of September.
- 50. Mergus servator, Linn.-Along the whole coast and on every river visited; first seen at Ste. Anne, June 30th.
- 51. Phalacrocorax carbo, Linn.-Between Bic and Green Island, middle of October.
- 52. Larus atricilla, Linn.-Whole coast.
- 53. Uria Grylle, Linn.-Hare Island; Green Island; Ste. Anne and near Martin River.

54. Colymbus glacialis, Linn .--- Metis Lakes ; Marsouin River and Rimouski. septentrionalis, Linn .- Skins of this bird were procured by Mr. 55. 22 Richardson in Anticosti.

CLASS REPTILIA.

66

Labor at the

die Reinerschließen und Annen. Annen

66

66

and the state - Black Soulites occup in

- 1. Tropidonotus sirtalis, Linn.-Throughout the district.
- 2. Rana pipiens, Gmel.-
- 3. Salamandra erythronota, Green .all. Coursess ---- have been allowed the lite
- 4. Bufo Americana, Leconte .--

CLASS PISCES.

ORDER ACANTHOPTERI.

- 1. Gasterosteus (not determined).-Metis River, above the high falls.
- 23 12 pungitius, Linn .- In numerous localities along the coast, from 2. an malaers Rivière du Loup downwards.
- biaculeatus, Mitch .- With the preceding species, but more 3. 11 abundant. Found also in Lake Matapedia.

4. Cottus Virginianus, Willughby .- Coast of Gaspé and Rimouski.

- gracilis ? Heck .- Restigouche River and Metis Lakes. 24 5.
- 6. Scomber vernalis, DeKay .- Ascends the St. Lawrence to Rimouski.

ORDER MALACOPTHEI.

- 7. Salmo salar, Linn .- Ascends all the rivers in the peninsula which are free from mill-dams.
- fontinalis, Mitch .--- In every stream and lake throughout the district. 8.
- trutta, Linn.-Abundant for a short distance up the clear streams of 9.

Gaspé.

10. Osmerus viridescens, Lesueur .- Whole coast below Green Island.

- 11. Alosa præstabilis, DeKay .-- Coast of Rimouski, middle of May.
- 12. " tyrannus, DeKay.-Rimouski Village.
- 13. Clupea virescens (?), DeKay .-- Whole coast as far up as the salt water extends.
- 14. " elongata, Lesueur .-- Whole coast also.
- 15. Mallotus villosus, Cuvier .-- With the last two species ; extremely abundant.

ORDER ANACANTHINI.

16. Ammodytes Americanus, DeKay .--- Coast of Gaspé,

17. Morrhua Americana, Storer .-- Ascends the river as far as Trois Pislotes.

- æglefinus, Cuvier .-- Taken with cod on the Gaspé coast. 18. "
- 19. pruinosa, DeKay.-Caught at the mouths of various rivers from the Chat upwards.

20. Motella cimbria (?), Parnell.-Ste. Anne.

21. Zoarcus viviparus, Cuvier .- Off the mouth of Marsouin River.

22. Hippoglossus vulgaris, Cuvier.-Ascends the river to Green Island.

- 23. Platessa vulgaris, Flem .- Several of the fishing stations on the Gaspé coast.
- 24. Cyclopterus lumpus,-Linn.-Ste. Anne; Green Island.

ORDER PLAGIOSTOMI.

25. Spinax acanthias, Cuvier.-Les Islets.

26. Raia radiata, Don .- Ste. Anne.

MISCELLANEOUS.

- 27. Coregonus.—Herring Trout, probably C. clupeiformis, are abundant in the Metis Lakes and River.
- 28. Cyprinus .-- Lake Matapedia and the Restigouche River.
- 29. Catasiomus.-Black Suckers occur in the Restigouche and the larger lakes of the district.
- 30. Anguilla.—Probably A. acutirositris, about the mouths of the rivers all along the coast.

31. Salmo.—An important species of Salmo, known as "Toag," abounds in the lakes of Rimouski County, but as no specimens were preserved nothing certain can be said about it.

ARTICULATA.

CLASS INSECTA.

ORDER COLEOPTERA.

Determined by Dr. J. L. Leconte of Philadelphia.

1.	Cicindel	a longilabris, Say Green Island Seigniory ; between Metis and
		Lake Matapedia; Ste. Anne.
2.	66	vulgaris, Say Ste. Anne ; Ruisseau de la Grande Vallée ;
		between Metis and the mouth of the Mata-
		· pedia.
3.	66	duodecimguttata, Dej Metis River; between Metis and the Mata-
		pedia ; Ste. Anne.
4.	66	Baltimorensis, Herbst. (repanda, Say.)-Rimouski; Metis River;
		Capucin.
5.	Brachin	48, (not determined)Abundant on Metis River.
		s reflexa, Lec. (marginata, Kirby) Rivière du Loup ; Rimouski ;
	÷	Metis ; Matanne.
7.	Calathus	gregarius, Say St. Simon ; from the mouth of the Marsouin to
		the Shickshock Mountains, fourteen miles up
		that river; Mount Commis on the Metis River.
8.	Platynus	sinuatus, Dej Point Levi ; St. Simon ; Marsouin River.
9.	66	extensicollis, Say Metis River.
10.	\$ 6	melanarius, Dej Point Levi, opposite Quebec.
11.	66	tenuis, Say Berthier and Ste. Anne.
12.	66	cupripenne, Say Point Levi, St. Simon and Ste. Anne.
13.	66	retractus, Lec Berthier, Rivière du Loup, and Ste. Anne.
14.	2.8	picipennis, Kirby, (lenum, Lec.)-Berthier, Marsouin River, and
		hetween Metis and the Matanedia.

		the second s
15.	Platynu	s lutulentus, Lec.—Point Levi.
16.		placidus, Say Berthier, Matanne, and Ruisseau de la Grande
17.	Pæcilus	lucublandus, Bay Very abundant at Point Levi, Berthier, Bivière
		da Loup, Green Island Village, St. Simon
		and Metis.
18	Dierosti	chus erythropus, Dej.—Point Levi.
19.	1 001 00000	
20.		mandibularis, KirbyGetween the mouth of the Marsouin and
AV.		the Shickshock Mountains.
0.1		A REAL PLACE AND A REAL AND A
21.	66	caudicalis, Say.—Bertleier and Green Island Seigniory. corvinus. Lec.—Point Levi.
22.		The second se
23.		or monomy router (our out room) - another the second
dine.		Loup to Ste. Anne; Mount Commis near the Metis.
24.		Luczotii, Dej. (var. præc.?)-Metis and Ste. Anne.
		adjunctus, LecRivière du Loup to Ste. Anne.
		ibera, LecRivière du Loup.
27.	66	pallipes, Kirby, (depressa, Loc.)—Rimouski.
28.	55	impuncticollis, Say Berthier and Ste. Anne.
29.	33	fallax, LecGreen Island Seigniory and Matanne.
30.	66	interstitialis, DejRimouski and Matanue.
		tylus Harrisii, Lec. (agricels, vide Harris.)-Point Levi and Berthier.
32.]	Harpalus	s viridiæneus, Beauv Very abundant at Green Island Seigniory,
		between Metis Lake and the Matapedia,
		Matanne, and Ste. Anne.
33.	66	pleuriticus, Kirby.—Abundant from Berthier to Rimouski.
34.	66	megacephalus, LecRivière du Loup.
35.	66	rufimanus, Lec -Ste. Anne.
36.	Chlæniu	s sericeus, Say Point Levi, Berthier, and St. Simon.
37.	4.6	chlorophanus, DejMetis River.
38.	66	tricolor, DejBerthier.
39.	Cychrus	(Sphæroderus) Brevoortii, LecRivière du Loup, St. Simon, Mount
	•	Commis twenty miles up Metis River, Ste.
		Anne and Marsouin River.
40.	Carabus	serratus, Say Rivière du Loup to Matanne, and between Metis and
		the Matapedia River.
41.	22	Lapilayi, LecRivière du Loup and Green Island Seigniory.
	Calosom	a calidum, FabrL'Islet, Rimonski, Metis, Matanne, and Ste. Anne.
		s Californicus, Mann. var. punctatissimus, Lec.—St. Simon.
		s longicornis, Say.—Berthler, Metis, and mouth of the Matapedia.
45.		angicollis, Randall.—Metis River.
	Rembidi	um dilatatum, Lec.—Metis River.
47.	Demonari ((
		lucidum, Lec.—Point Levi.
40.	Anabus	confluens, Say. (O. Oligbukii, Kirby.)-Mouth of Metis River.
49.	ng an as	striatus (?), Say Rivière du Loup, Green Island Seignlory, and
80	37.0000	Ste. Anne.
50.	тестори	orus velutinus, Fabr.—Metis River.

- 51. Silpha lapponica, Herbst.-Very abundant at Ste. Anne.
- 52. Staphylinus villosus, Grav.---Bimouski, Metis, Matanne, and Ste. Anne.
- 53. Omosita colon, Fab.-In vast numbers in fields manured with capeling.
- 54. Pediacus planus, Lec.-Between Metis and Matapedia.
- 55. Byrrhus picipes, Kirby .- Ste. Anne.
- 56. Platycerus depressus, Lec.-Ste. Anne.
- 57. Aphodius fussor, ("absolutely the same as the European," Leconte, in lit.)-Rivière du Loup and Ste. Anne.
- 58. " fimetarius, Fabr.-Abundant from Metis to the Matapedia.
- 59. " n. sp. (?)-Metis.
- 60. Lachnosterna fusca, Frohlich .-- Point Levi and Rivière du Loup.
- 61. Dichelonycha subvittata, Lec.-Ste. Anne.
- 62. Ancylochira maculiventris, Say .--- Metis River, and between Metis and the Matapedia.
- 63. Ellychnia corrusca, Dej.—Capucin, Ste. Anne, and Ruisseau de la Grande Vallée.
- 64. Meloe rugipennis, Lec .- Between Metis and the mouth of the Matapedia.
- 65. Serropalpus substriatus, Hd.-Metis River.
- 66. Upis reticulata, Say .- Metis.
- 67. Tomicus (not named) .--- Between Metis and the Matapedia.
- 68. Physocnemum ligneum, Fabr.-Green Island Seigniory.
- 69. Monohammus confusor, Kirby .- Metis.
- 70. " scutellatus, Say .--- Metis and Ste. Anne.
- 71. Chrysomela scalaris, Lec.-Metis.
- 72. Galleruca (not named) .- Between Metis and the Matapedia.
- 73. Coccinella novemnotata, Fabr .-- Rimouski and Metis.

ORDER LEPIDOPTERA.

Determined by Mr. D'Urban.

(a) Rhopalocera.

- 74. Papilio turnus, Linn.—From Cape Chat to Martin River, from June 18th till the end of July. Extremely abundant.
- 75. Colias philodice, Godart.—Cape Chat and Ste. Anne, from the middle of June till the middle of July; between Metis and Lake Matapedia, August 17th; along the Restigouche during the latter half of

August ; last seen September 1st.

- 76. Pieris oleracea, Harris.—St. Simon, May 28th; Ste. Anne, from June 20th to the middle of July. Common.
- 77. Limenitis Arthemis, Drury.-Ste. Anne, July 16th ; Marsouin River, July 26th.
- Cynthia cardui, Linn.—Seigniory of Grand Metis, August 16th; Dalhousie N. B., August 25th.
- Vanessa J. album, Boisd.—Junction of the Patapedia and Awaganasees, September 12th.
- 80. " Antiopa, Linn .- Metis and near Rimouski, September 29th.

- 81. Grapta Progne, Fab.—From Rivière du Loup to Ste. Anne, from May 18th till July 19th ; Lake Matapedia, August 17th; along the Restigouche and Patapedia Rivers till September 12th.
- 82. Grapta C. aureum, Cramer (?)-Mouth of Awaganasees Brook, September 12th.

83. Argynnis Aphrodite, Fab.—First observed at Ste. Anne on the 20th of June and very abundant there for the next month; Marsouin River, July 26th; between Metis and Lake Matapedia and along the Rest gouche in August, and last seen at the mouth of the Awaganasees, September 12th.

- myrina, Cramer.—Ste. Anne, end of June and beginning of July; between Metis and Lake Matapedia, August 16th.
- Bellona, Godart.—Mouth of Matapedia River, Angust 27th.
 Melitaa Tharos, Oramer.—Ste. Anne, beginning of July.
- 87. Polyommatus pseudargiolus, Boisd.-Rivière du Loup, May 19th, and thence as far down as Chat River, till June 18th.
- 88. Hesperia ------ (?) -- Metis, August 13th ; Lake Matapedia, August 17th.

(b) : Heterocera.

- 89. Orgyia ----- (?) -- Matapedia River, August 20th.
- 90. Ctenucha Latreillana, Kirby .- Ste. Anne, June 28th. Abundant.
- 91. Phragmatobia fuliginosa, Linn.-Matanne, June 12th.
- 92. Mamestra _____ (?) -Ste. Anne.
- 93. Plusia ----- (?) --Common in Gaspé and on the Restigouche.
- 94. Pyralis ----- (?) Mouth of the Matapedia River.
- 95. Crambus ----- (?) -Very abundant in meadows at Ste. Anne, and at the

mouth of the Matapedia.

Five undetermined species of Geometric Moths.

CLASS CRUSTACEA.

OBDER DECAPODA.

1. Cancer irroratus, Say .--- Whole coast below Green Island.

- 2. Hyas fissirostra, Say sp.-With the preceding species.
- 3. Pagurus Bernhardus, Fabr.-Coast of Gaspé and Rimouski.
- 4 Homerus Americanus, Milne-Edw.-Rars on the coast of Rimouski and on

the north coast of Gaspé, but abundant in Gaspé Bay, on Anticosti and in the Bay of Chaleur.

- 5. Astacus Bartonii, Bosc.-Metis, Matapedia and Restigouche Rivers.
- 6. Crangon vulgaris, Fabr .- Coast of Gaspé and Rimouski.
- 7. " sculptus (?); Bell.-Off Cape Chat.
- 8. Hippolyte (not determined)-Near Metis.
- 9. Orchestia (not determined)-Whole coast.

CLASS ANNULATA.

ORDER TUBICOLE.

Determined by Dr. J. W. Dawson.

1.	Spirorbis	porrectu,N	orth	coast of	Gaspé.
2.		sinisi''0811,	66		"
3.	66	eurinaie,	6.6		65
4.	44	viirea,-	66		-4.6
5.	4.6	cancellata,	44		44

spiril'un....On littoral Algae, whole coast below Rivière du Loup.
 Serpula (vermilia) serrula, Stimpson...-North coast of Gaspé.

MOLLUSCA.

CLASS GASTEROPODA.

ORDER PULMONIFERA.

(Terrestrial.)

l.	Helix	alternata,	Say Common	from	Qu	ebec	along	; the	whole	coast	into
			G	aspé;	it	appe	ars t	o be	diffused	lover	the
whole peninsula.											

 albolabris, Say.—From Quebec to Metis; Lake Matapedia; along ille Restigoucle River from Dalhousie to the mouth of the Patapedia. I never met with this species in the County of Gaspé.

- "monodon, Rackett.—Point Levi; along the banks of the Restigouche from Dalhousie to the mouth of the Patapedia.
- 4. " Sayii, Binney.--Restigonche River, about five miles above the mouth of the Matapedia.
- 5. " concava, Say .- Point Levi ; abundant.
- 6. " hortensis, Müll.—From all that I could ascertain regarding this species, it appears to have diffused itself over a strip of country several miles in width, bordering on the St. Lawrence and extending from Metis to Gaspé Bay.
- 7. " arborea, Say.—Throughout the whole district; very abundant. Occurs on the Island of Anticosti.
- 8. " striatella, Anthony.-With the last species and equally abundant.
- 9. " *lineata*, Say.—Numerous localities on the coast, from Berthier to Marsouin River.
- 10. " labyrinthica, Say .- Rivière du Loup and Green Island.

11. Helix pulchella, Müll.—Berthier, mouth of Magdalen River and Dalhousie, N. B.

- 12. " asteriscus, Morse.-Valley of the Marsouin River.
- "chersina, Say.—Trois Pistoles; Capucin; Ste. Anne; along the vallies of the Marsouin, Magdalen and Mata-

pedia Rivers, and at the mouth of the Patapedia.

14. Helix (undetermined) .- A young shell of one of the larger species, but differing from any of the preceding ; Rivière du Loup.

15. Succinea awara, Say .- Matanne; mouth of Magdalen River; several localities on the Restigouche.

ovalis, Gould .- Metis, Matanne and Ste. Anne. 16. 66

obliqua, Say .- Throughout the district. " 17.

18. Achalina lubrica, Mull.-Rivière du Loup ; Trois Pistoles ; Metis Lakes and along the Restigouche.

19. Bulimus harpa, Say sp .- Metis; mouth of Magdalen River, and very abundant in the Marsouin Valley.

20. Vitrina pellucida, Drap .-- Rivière du Loup; Trois Pistoles; Ste. Anne; Restigouche River ten miles above its janction with the Matapedia.

21. Pupa (Vertigo) simplex, Gould .- Valley of the Marsouin ; along the Restigouche and at Metis.

(Fresh Water.)

22.	Physa heterostropha, SayThroughout the district; very abundant.						
23.	" aurea, LesSeveral localities in the County of Rimouski.						
24.							
25.							
26.	66	marginata, SayNear Rimouski Village.					
27.	7. Limnaa stagnalis, Lam Extremely abundant in the Metis Lakes, and in						
		the lakes on the Rimouski River.					
28.	66	caperata, Say Lakes Metis and Matapedia, and the Metis and					
		Restigouche Rivers. Abundant.					
29.	13	umbrosa, Say Ste. Anne; a creek about two miles below Chat					
		River ; Metis and Restigouche Rivers.					
30.	66	catiscopium, Say Rimouski, Restigouche, and Dartmouth rivers.					
31.	. " apacina, LeaLiving in the St. Lawrence at Point Levi; in the						
		Metis, Rimouski and White Rivers.					
32.	66	acuta, Lea.—Upper Lake Metis ; abundant in Marl Lake, Anticosti.					
83.	66	umbilicata, Say Metis and Ste. Anne.					
34.	66	reflexa, Say Upper Metis Lake.					
35.	61	pallida, AdamsLarge Lake Matapedia; near Cape Chat.					
36.	66	modicella, SayGreen Island Village ; Rimouski ; Ste. Anne.					
37.	66	parva, LeaRivière du Loup.					
38.	66	decollata, SayLarge Lake Matapedia; Rimouski Village.					
39.		alternata, or new.—Point Levi.					
40	. Plan	orbis trivolvis, Say.—Rimouski, Metis and Restigouche Rivers.					
41	. "	campanulatus, Say Lakes Metis and Matapedia.					
42		oreal summer's Dal HERMROACHE MILLET.					
43	- "	parvus, Say Throughout the district.					
44	68	deflecture Say - Larva Laka Matanadia					

ORDER PROSOBRANCHIATA.

(Fresh Water.)

45. Amnicola porata, Say .- Little Lake Matapedia.

46. Valvata tricarinata, Say .- Matapedia Lakes.

47. " humeralis, Say (or a new species).—Matanne; small lake at the head

of Awaganasees Brook ; Little Lake Matape-

dia.

48. " sincera, Say .-- Marl Lake, Anticosti. Abundant.

Note.—Many of the above species of land and fresh water Gasteropoda were kindly determined for me by W. G. Binney Esq., of Burlington N. J. and Dr. Isaac Lea, of Philadelphia.

(Marine.)

49. Fusus scalariformis, Gould .- Peter River; Ste. Anne ; Marsouin.

50. " gracilis Alder .- Trent; Ste. Anne ; Marsouin.

51. " tornatus, Gould .- Rimonski Village ; near Ste. Anne.

52. " decemcostatus, Say. --- Near Cape Gasp6 (collected by Sir W. E. Logan in 1844.)

"

53. " rufus, Gould .- Ruisseau de la Grande Vallée.

- 54. " Bamffius, Flem.— " "
- 55. Bela cancellata, M. & A.- "

56. Pleurotoma bicarinata (?), Couth.-

57. Buccinum undatum, Linn .- Whole coast below Rivière du Loup.

58. " Donovani, Gray .- Several localities below St. Flavie.

59. Nassa trivittata, Say .--- Gaspé Bay and Bay of Chaleur.

60. " obsoleta, Say .- Vicinity of Cape Gaspé.

61. Purpura lapillus.-Lam.-Whole coast below Metis.

62. Trichotropis borealis, Sowerby .- Ste. Anne and near Cape Chat.

63. Velutina halioloides, Müll .- Ste. Anne and Marsouin.

64. Lamellaria perspicua, Lovèn.-Ruisseau Vallée.

- Natica heros, Say, ampullaria, Lam.—In sandy bays on the Gaspé coast; at Dalhousie, Bay of Chaleur.
- 66. " clausa, Brod. & Sow .- Several localities between Bic and Marsouin
- 67. " triseriata, Say .--- Magdalen Bay.
- 68. " flava ? Gould .- Rimouski; Les Islets ; Claude.
- 69. " helicoides, Johnston.-Marsouin.

70. Chemnitzia .- One or more species of Chemnitzia dredged off Marsouin.

71. Aphorhais occidentalis, Gould .- Bic ; Ste. Anne ; Claude ; Marsouin.

72. Rissoa minuta, St .- Green Island and Long Point.

- 73. Lacuna vincta, Turt .-- Whole coast below Rimouski.
- 74. Littorina littoralis, F. & H., palliata, Gould.-Whole coast below Rivière Ouelle.

75. Littorina rudis, Gould, (including tenebrosa).-With the preceding species.

 Margarita cinerea, Gould.—Ste. Anne; Ruisseau Vallée; Peter River and Marsouin.

- 77. " undulata, Sow.-Ste. Anne; Ruisseau Vallée.
- 78. " helicina, Müll.-Trent; Les Islets; Ste. Anne.

- 79. Skenea costulata, F. & H.-Marsouin.
- 80. Diadora Noachina, Gray .-- Capucin ; Ste. Anne ; Marsouin.
- 81. Crepidula fornicata, Lam .- Dalhousie, Bay of Chaleur.
- 82. Acmæa testudinalis, Hanley.-Whole coast below Rivière du Loup, also in Bay of Chaleur.

83. " cæca.-Marsouin.

84. Chilon-marmoreus, Fabr.-Bic, and whole coast of Gaspé

CLASS L'AMELLIBRANCHIATA.

Transienty bad toul Lagrad on to (Marine.)

- 1. Pholas crispata, Linn.-Bic; Rimouski; near the Trent.
- 2. Saxicava rugosa, Lam.—Les Islets; Ste. Anne; Cape Chat; Marsouin Claude.
- 3. Mya arenaria, Linn.-Whole coast below Rivière Ouelle, and in Bay of Chaleur.
- 4. " truncata, Linn.-Numerous localities on the coast of Rimouski and Gaspé.
- 5. Glycymeris siliqua, Lam .- Cape Chat; Ruisseau Vallée; and Marsouin.
- 6. Osteodesma hyalina, Couth.-Ste. Anne.
- 7. Machæra costata, Gould .-- Rimouski,
- 8. Solen ensis, Linn.—Bic ; Rimouski, and numerous localities on the coast of Gaspé.
- 9. Tellina proxima, Brown.-Ste. Anne ; Ruisseau Vallée ; Marsouin.
- "Grænlandica, Beck.—Whole coast below Bay St. Paul (fifty-five miles below Quebec), and in the Bay of Chaleur.
- 11. Mactra ovalis, Gould.-Bic; Rimouski; Metis, and in sandy bays everywhere on the Gaspé coast.
- 12. Mesadesma gratatum, Gould.-Whole coast below Green Island. Extremely abundant.
- 13. Venus gemma, Tott.-Green Island.
- 14. Aphrodite Grænlandica, St.-Bio; Rimouski; Metis; Ste. Anne; Ruisseau Vallée.
- 15. Cardium Islandicum, Linn,-Bic; Rimouski; Metis; Ste. Anne.
- 16. Cardita borealis, Con.—Marsouin ; Capucin ; Ste. Anne ; Ruisseau Vallée.
- 17. Astarte sulcata, Costa.-Bic, and various localities on the Gaspé coast.
- 18. " elliptica, Brown.-Marsouin.
- 19. " compressa, Mont.-Marsouin.
- 20. Lucina flexuosa, Gould .- Ste. Anne; Ruisseau Vallée and Marsouin.
- 21. Lima subauriculata, Mont .- Ste. Anne.
- 22. Mytilus edulis, Linn .- Whole coast below Kamouraska.
- 23. Modiola discors, Linn .- Ste. Anne; Marsouin.
- 24. " plicatula, Lam.-Vicinity of Gaspé Bay.
- 25. " glandula, Tott.-Ste. Anne; Ruisseau Vallée; Marsouin.
- 26. " pectinula, Gould.-Ruisseau Vallée ; Marsouin.
- 27. " nexa, Gould.-Ruisseau Vallée.

28. Nucula myalis, Couth .- Numerous localities on the Gaspé coast.

29. " icauis, Turt .- Capucia ; Ste. Anne ; Ruisseau Vallée.

30. Pecien Magailanicus, Lam.-Ste. Anne; Claude and Gaspé Bay.

31. " Islandicus, Müll .-- Whole coast below Metis.

32. Anomia ephippium, Linn.-Ste. Anne ; Marsouin.

(Fresh Water.)

- 33. Unio complanatus, Lea.—Living in the St. Lawrence as far down as Berthier. Valves both of this species and of Uradiatus were frequently found on the beach the whole way to Gaspé. They had probably drifted from the fresh water of the St. Lawtence, as no species of Unio was found in any of the rivers or lakes of our present district.
- 14. Margaritana arcuata, Barnes sp.—Green and Limouski Rivers, and both the Matapedia Lakes.
- 35. Anodonta subcylindracea, Lea.—Gand Lac (ten miles south of Rimouski Village); Lake Matapedia; small lake six miles S. W. of Metis.

36. " new species .- Berthier.

- 37. " edentula, Say.-Lake Matapedia.
- 38. " fragilis, Lam.-Metis Lakes.
- 39. " implicata, Say.-Berthier.

40. Cyclas similis, Say .- Meiis Lakes and a small lake six miles S. W. of Metis.

- 41. " dubia (?), Say .- Throughout the district.
- 42. " (undetermined).-Ste. Anne.

CLASS BRACHIOPODA.

1. Hypothyris psittacea, King.—Ste. Anne; Ruisseau Vallée; Marsouin. Abundant.

CLASS POLYZOA.

ORDER CHEILOSTOMATA.

The Polyzoa dredged at Marsouin on the north coast of Gaspé, were kindly determined by Dr. J. W. Dawson, Principal of McGill College. The following is his communication in full.

The Polyzoa in Mr. Bell's Collections are numerous and fine, but much time and care would be required for their accurate determination. The appearances presented in various stages of growth and preservation, are so perplexing, and the characters given for the species of authors, of so little value, that little can be done with a collection of dead cells, except to indicate the described species with which they seem to be identical. The following species were all attached to dead shells and stones, from a depth of about thirty fathoms.

1. Hippothoa catenularia, Jameson.

2. " divaricata, Elliot.

3. Hippothoa expansa. New species. Description. Cells oval, depressed, and

expanded at the sides, not contracted at the base, branching dichotomously. When magnified the surface presents indistinct transverse wrinkles and delicate longitudinal lines. Aperture, small, round, with a slight sinus. Texture hyaline, but less delicate than H. divaricata.

All the three species above mentioned are found associated on small pebbles and shells.

4. Lepralia pertusa, Thompson.-Very abundant.

44 8. Peachii, Johnston .- Very abundant also.

- 33 trispinosa, Johnston.-Abundant. 6.
- hyalina? *, Johnston.-Rare. 7. 66 Redenations gainer (of the WO of a

11 punctata, Hassal.---Rare. 8.

San File A. Market Y to

puncturata, Busk .- A little group of three cells on a shell of Mactra ovalis have the precise characters of this species, obtained by Busk from the English Crag. It appears still to live, though as a rare species, in the Gulf of St. Lawrence.

10.

9.

16

"

New species. Description. In large patches. Young Belli.

cells granular, semi-hyaline, confluent; mouth immersed, sinuated, with a vibraculum or avicularium inside the middle of the lower lip; ovi-cells rounded, granulous like the cells. Old cells white, opaque, flat above, and separated by a deep sinuous furrow. Cells having a strong tendency to form rows radiating from the centre of the patch. I can find no described species possessing the above characters. It is allied to L. concinna, Busk, but differs in essential points from his description and figure.

11.

"

plana. New species. Description. Cells flat, confluent, shallow; walls deeply and irregularly furrowed; mouth rounded above, straight below, often with a narrow sinus in the middle. Young cells hyaline; old cells, opaque and deeply furrowed in a stellate manner. Forms very thin and flat expanding crusts. L. adpressa, Busk, from Chilce, resembles it more nearly than any other species known to me.

12. Membranipora Lacroixi, Busk, or a nearly allied species.

lineata *, Busk, Flustra lineata, Fabricius, "Fauna Groen-" 13. landica."

* The species marked thus were found by Fabricius in Greenland.

- 14. Cellepora pumicosa *, Ellis .--- On Sertularia.
- 15. " cervicornis, Borlase.
- 16. " ramulosa, Linn., -- or allied species.
- 17. Carbasea papyrea, Gray.—The frond is narrower than in British examples, but the cells are of the same form.
- 18. Diastopora obelia, Fleming, or closely allied species.
- 19. Tubulipora flabellaris.* Fabricius.
- 20. " hispida.* Johnson.-It is the Madrepora verrucaria of Fabricius.
- 21. " phalangea ? * Couch-Of the form of T. flabellaris, but dotted with

pores and having larger tubes, which are grouped in bundles. Perhaps it is *T. densa*, Stimpson. Its colour is often light blue. Fabricius seems to have seen it and placed it with *T. fabellaris*.

22. Cellularia (species undetermined.)

Many more species were dredged but have not yet been determined.

RADIATA.

CLASS ECHINODERMATA.

ORDER ASTEROIDEA.

- 1. Ophiocoma bellos, Link .--- Ste. Anne and Marsouin ; abundant.
- 2. " Gordsiri? Forbes.-Marsouin.

 Astrophyton scutatum, Link.—Green Island; Gaspé Bay; St. Nicolas (north shore). Said to be common on the coast of Rimouski.

- 4. Cribella oculata (?) Pennant.-Near Ste. Anne.
- 5. Solaster papposa, Linn.-Marsouin.
- Asteracanthion polaris, Müll.—Very abundant along the whole coast below Rimouski.
- 7. " rubens, Linn.—Les Islets.

ORDER ECHINOIDEA.

- Echinarcahnius Atlanticus.—On muddy and sandy bottoms, along the whole coast below Rimouski.
- 9. Echinus granularis, Lam .- Whole coast below Rivière du Loup.

ORDER HOLOTHURIDEA.

- 10. Cucumaria communis, Forbes .- Between Cape Chat and Ste. Anne; abdt.
- 11. Psolus phantapus, Linn .--- Various localities between Metis and Ste. Anne.

CLASS ACALEPHÆ.

ORDER HYDROLDEA.

"

...

- 1. Sertularia polyzonia *, Johnston .- Dredged off Marsouin.
- 2. " argentea*, Ellis.— "
- 3. " filicula, Ellis -----
- 4. " latiuscula?, Stimpson, or a closely allied species.

None of the above have ovicapsules.

Six or more different Sponges, some of them beautiful forms, were collected.

PLANTS.

1 am indebted to Mr. D'Urban, late of the Geological Survey, for preparing the following catalogue of Plants collected by me in the eastern peninsula of Lower Canada. Numerous species, about which Mr. D'Urban was in doubt, were kindly determined by George Barnston Esq., of the Hudson's Bay Company.

Ranunculaceæ.

- 1. Anemone Pennsylvanica, Linn.-F. F.*, August 12th, Metis.
- 2. Thalictrum cornuti, Linn .-- F. F., July 16th, Ste. Anne.
- 3. Ranunculus repens, Linn. " "
- 4. " acris, Linn. " "
- 5. " (undetermined) .- No flower, September 1st, River Restigouche.
- 6. Caltha palustris, Linn .- F. F., June 5th Rimouski.
- 7. Aquilegia Canadensis, Linn .- F. F., May 16th, L'Islet.

 Nuphar advena, Ait., (a very small form.)—F. F., August, west end of Lake Matapedia.

Sarraceniacea.

- Sarracenia purpurea, Linn.—F. F., June, Ste. Anne. Fumariacea.
- Corydalis aurea, Pursh.—F. F., August 30th, Restigouche River. Crucifera.
- Sinapis arvensis, Linn.—F. F., July 11th, Ste. Anne. Violacea.
- Viola cucullata, Ait.—F. F., May 30th, St. Simon. Cistaceæ.
- Hudsonia tomentosa, Nutt.—F. F., August 31st, River Restigouche. Parnassiaceæ.
- 14. Parnassia Carolinianum, Michx.-F. F., August 30th. Caryophyllaesee.
- 15. Silene inflata, Smith .- F. F., July 6th, Ste. Anne.
- Mœhringia lateriflora, Linn.—F. F., July 23rd, Portage between Martin and Marsouin rivers.
- Spergula arvenşis (?), Linn:-No flower, August 12th, Metis. Oxalidaceæ.
- 18. Oxalis acetosella, Linn.-Very abundant up the River Marsouin.
- "stricta, Linn.—Going to seed, August 30th, River Restigouche. Anacardiaceæ.
- Rhus Toxicodendron, Linn.—Fruit ripe, August 31st, River Restigouche. Sapindaceæ.
- Acer spicatum, Linn.—Abundant everywhere on low land; just out of flower, July 5th, Ste. Anne. In seed, Sept. 11th, mouth of the Awaganasees Brook.
- 22. " saccharinum, Wang. (Hard Maple) .- On rich soil only.

Nymphaacea.

- Leguminosæ. 23. Trifolium repens, Linn .- Abundant round clearings, &c., throughout the district. 24. Desmodium Canadense, D. C .- F. F., August 12th and 31st, River Restigouche. 25. Vicia Cracca, Linn .- F. F., July 11th, Ste. Anne. 26. Lathyrus palustris, Linn .- F. F., August 4th, mouth of the Marsouin. 27. Oxytropus Lamberti, (?) Pursh .- F. F., August 31st, River Restigouche. Rosaceæ. 28. Prunus pumila, Linn.-Fruit nearly ripe, August 31st, River Restigouche. 66 Pennsylvanica, Linn .- Abundant throughout the Counties of Ri-29. mouski and Bonaventure. 30. Virginiana, Linn .- Fruit ripe, Sept. 1st, River Restigouche. 31. Agrimonia Eupatoria, Linn .-- In seed, August 21st, fifteen miles up the River Matapedia. 32. Potentilla anserina, Linn .- F. F., August 4th, mouth of the River Marsouin. 33. Fragaria Virginiana, Ehrhart.-Grass land throughout the district. Fruit ripe beginniug of July, Ste. Anne. 34. Rubus triflorus, Rich .- Fruit ripe, July 12th, Ste. Anne; mouth of the Awaganasees. strigosus, Miche .- Extremely abundant on burnt land and about 35. fences throughout the district. 36. Rosa blanda, Ait .- In blossom, July 5th and 20th at Ste. Anne, and August 12th at Metis. 37. Cratægus tomentosa, Linn.-River Restigouche. 38. Pyrus Americana, D. C.-Moderately abundant throughout the district. Onagraceæ. 39. Epilobium augustifolium, Linn.-F. F., July 16th, Ste. Anne. 66 coloratum, Muhl.-In seed, July, three miles up the River Marsouin. 40. 41. Enothera biennis, Linn .-- F. F., July 11th, Ste. Anne; August 30th, mouth of the River Matapedia. 42. Circæa Alpina, Linn.-In flower, July 31st, mouth of the River Marsouin. Saxifragacea. 43. Mitella nuda, Linn.-Seed ripe, July, 3 miles up the River Marsouin. Umbelliferæ. 44. Heracleum lanatum, Michx .--- F. F., July 16th, Ste. Anne. 45. Sium lineare, Mich.-F.F., August 12th, Metis. Cornacea. 46. Cornus Canadensis, Linn.-F. F., July 5th, Ste. Anne. 47.
 - "stolonifera, Michx.—F. F., June, Ste. Anne. Caprifoliaceæ.
 - 48. Linnea borealis, Gronov.-F. F., June, Ste. Anne, and abundant everywhere.
 - 49. Lonicera ciliata, Muhl.-In fruit, July 30th, Marsouin river.
 - 50. Diervilla trifida, Mænch.-F. F., August 30th, River Restigouche.

51.	Sambucus Canadensis, Linn Abundant on low land.
	Viburnum opulus, Linn F. F., July 16th, St. Anne.
	Composite
53.	Eupatorium purpureum, Linn F. F., Sept. 3rd, mouth of the River Pata-
	. Pediasted Caraterial, Long - F
54.	" ageratoides, Linn F. F., July 31st, mouth of the River Mar-
	souin, and August 30th, River Restigouche.
55.	Aster miser, Linn, Ait F. F., August 12th, Metis.
56.	" simplex, (?) Willd "
57.	" longifolius, (?) Lam"
	Diplopappus umbellatus, Torr. and GrF. F., June 30th, mouth of the River Matapedia.
59.	Solidago bicolor, Linn Going out of flower, August 30th, River Resti-
	gouche.
60.	" Canadensis, LinnF. F., August 12th, Metis.
61.	Achilles millefolium, Linn - F. F., July 11th, Ste. Anne, and mouth of the Awaganasees, September.
62.	Leucanthemum vulgare, Lam F. F., July 4th, Ste. Anne, and August
	30th, River Restigouche.
63.	Cirsium Muticum, MichxF. F., August 30th, mouth of the River Matape- dia.
64.	" pumilum (?), Spreng Out of flower, August 30th, River Resti-
	gouche.
65.	Hieracium Canadense, Michx F. F., August 30th, River Restigouche.
66.	Nabalus racemosus, Hook. ("variety with truncate and obcordate leaves."
	G. B.)-August 30th, River Restigouche.
07	Lobeliaceæ. Lobelia Kalmii, Linn.—F. F., August 30th, River Restigouche.
01.	
~~	Campanulaceæ.
68.	Campanula rotundifolia, LinnF. F., August 4th, mouth of the River
	Marsouin, and August 30th, River Resti- gouche.
	Ericaceæ.
69.	Vaccinium Pennsylvanicum, (?) Lam In great profusion on hills which
	had been burnt over.
70.	Chiogenes hispidula, Torr. and GrIn great abundance throughout the district.
71.	Andromeda polifolia, LinnF. F., July 16th, Ste. Anne.
	Pyrola rotundifolia, Linn " " "
	Plantaginaceæ.
73.	Plantago maritima, LinnF. F., August 4th, mouth of the River Marsouin.
17.4	Primulaceæ.
44.	Primula farinosa, Linn.—Abundant all along the southern shore of the Gulf. F.F., end of May and June.
	Lentibulaceæ.
75	Utricularia zulgaria (2) Linn Matia

75. Utricularia vulgaris (?) Linn .--- Metis.

Scrophulariacea.

- 76. Chelone glabra, Linn.-F. F., August 12th, Metis.
- 77. Veronica Americana, Schweinitz.—Nearly out of flower, July 12th, Ste. Anne.
- Pedicularis Canadensis, Linn.—F. F., August 10th, Matan. Labiatæ.
- 79. Lycopus Virginicus, Linn., (a very coarse form).—In flower, August 30th, River Restigouche.
- 80. Brunella vulgaris. Linn .- In flower, July 11th. Ste. Anne.
- 81. Scutellaria nervosa, Pursh.—In flower, August 12th, Metis.

Borraginaceæ.

- Mertensia maritima (?), Don.—In flower, beginning of July, Ste. Anne. Apocynacea.
- Apocynum androsæmifolium, Linn.—F. F., August, between Metis and Lake Matapedia.

Asclepiadaceæ.

84. Asclepias cornuti, Decaisne .- Abundant all along the Restigouche.

Oleaceæ.

85. Fraxinus sambucifolia, Lam., (Black Ash).—In valleys, and along the shores of the Lakes.

Polygonaceæ.

- Rumex acetosella, Linn.—Coming into flower, July 16th, Ste. Anne. Urticacez.
- 87. Ulmus Americana, Linn., (Swamp Elm).--Very abundant, and of large size, along the River Restigouche.

Cupuliferæ.

88. Corylus rostrata, Ait., (Hazel-nut).-Marsonin River.

Betulaceæ.

- Betula papyracea, Ait., (White Birch).—The most abundant deciduous tree throughout the eastern peninsula, and reaching a large size.
- 90. " excelsa, Ait., (Yellow Birch).--Most abundant round Lake Matapedia, and in the valleys of the Rivers Marsouin and Restigouche; generally associated with Hard Maple on rich soil.
- 91. Alnus incana, Willd., (Alder.)-Everywhere bordering the streams and rivers, forming dense thickets.

Salicaceæ.

92. Populus tremuloides, Michx., (Common Poplar).-Abundant on high lands.

93. " balsamifera, Linn., (Balsam Poplar, Balm of Gilead).—Abundant on the borders of rivers and lakes.

Conifera.

- 94. Pinus resinosa, Ait., (Red Pine).—Abundant, but of small size, along the upper part of the River Patapedia.
- 95. " strobus, Linn., (White Pine).-Abundant everywhere.

96. Abies balsamea, Marshall, (Balsam Fir) .--- Very abundant.

- 97. " nigra, Poir., (Black Spruce).—The principal, and in many places the sole tree covering the hilly country of the eastern peninsula.
- "alba, Michx., (White or "Sea Spruce" of the Indians).-The commonest tree along the coast and rivers.
- Larix Americana, Michx., (Tamarack).---Bather scarce, but occurring in every variety of situation throughout the district.
- 100. Thuja occidentalis, Linn., (White Gedar).—Very abundant in the vallies of all the rivers, reaching a very large diameter, but no great height.
- 101. Taxus baccata, Linn., var. Canadensis, (Ground Hemlock).-Abundant amongst trees on low ground.

Alismaceæ.

 Sagittaria variabilis, Engelm.—F. F., August 15th, Metis. Orchidacea.

- 103. Platanthera flava, Gray .-- F. F., September 1st, River Restigouche.
- 104. "psycodes, Gray.—F. F., August 17th, West end of Lake Matapedia.
- 105. Spiranthes decipiens, (?) Hooker.—Coming into flower, July 30th, Marsouin River.
- 106. Corallorhiza Macræi, Gray.—Going to seed, July 31st, three miles up the River Marsouin.

Iridaceæ.

108. Sisyrinchium Bermudianum, Linn., (variety mucronatum, Gray).—In flower, July 16th, Little Ste. Anne.

Smilaceæ.

109. Trillium erectum, Linn., (very large).—Fruit ripe, July 31st, three miles up the Marsouin River.

Liliaceæ.

- 110. Smilacina stellata, Desf .--- F. F., June, Ste. Anne.
- 111. " bifolia, Ker.-In seed, but not ripe, July 20th, Marsouin River.
- Clintonia borealis, Raf.—Throughout the district. Melanthaceæ.
- 113. Streptopus roseus, Michx.-F. F., June, Ste. Anne.
- 114. Tofielda glutinosa, Willd.—Seed ripe, August 30th, River Restigouche. Cyperacew.
- 115. Eriophorum vaginatum, Linn.-Ste. Anne. Gramineæ.
- 116. Phleum pratense, Linn., (Timothy).—Table-topped Mountain, 3800 ft. above the sea; upper part of Magdalen River grows luxuriantly along roadsides, in openings in the woods &c.
- 117. Calamagrostis Canadensis, Beauv.-Shickshock Mountains.
- 118. Elymus Canadensis, Linn.-River Restigouche.

^{107.} Iris versicolor, Linn .- F. F., July 4th, Ste. Anne.

		triata, Michx('		. ,			ock						
		ææ.			and the second								
	D. Equisetum pratense, Ehrh.—Metis.												
-12	Filices.												
121.	121. Asplenium feliz-formina, R. BrMouth of the Awaganasees Brook.												
122.	22. Aspidium spinulosum, Swartz "												
	123. Osmunda regalis, Linn Round Metis Lake, &c.												
124. Botrychium Virginicum, SwartzFertile fronds ripe, July 28th, River Marsouin.													
	Lycopodia	iceæ.	God II.	91									
125.		ium lucidulum, Mi			River Res	tigouche.	12						
126.	64	dendroideum.	Michx "	4	ALC: NO.	44	505						
127.	66	clavatum, Li				66							
128.	66	complanatum	, Linn.,-"	13 13		. 46							
	Musci.				Sahang libi								
129.	Polytric	hum commune. Lir	nCollec	ted on the Riv	er Marson	ain.							
	-	splendens, Hedw.					897						
131.	66	Schreberi, Willd.			25								
132.	66	Crista-Castrensis,	L	66	48								
133.	66	reptile, Michx	44	25	4								
	Lichenes.												
134.	Peltigera	aphthosa (?) Hoff	en, infert.	River Marson	in.		241						

135. Sticta pulmonaria, Ach.-

