

LEGEND

This legend is common to GSC OF4754, OF5070, OF5237, OF5305, OF5306, OF5307, OF5309, OF5527, and OF5540. Coloured legend blocks indicate units that appear on this map.

NOTE: In areas where the surficial cover forms a complex pattern, the area is coloured according to the dominant unit and labelled in descending order of cover (e.g. O-T). Where buried aggregate deposits (sand and gravel - commonly associated with O¹ or O² surficial units) are known, or suspected, areas are coloured according to the overlying unit and labelled in the following manner: LV/Gd.

QUATERNARY SURFICIAL DEPOSITS

- POST-LAST GLACIATION**
- NONGLACIAL ENVIRONMENTS**
- AN** ANTHROPOGENIC DEPOSITS: culturally-made or modified geological materials such that their original physical properties (e.g. structure, cohesion, compaction) have been drastically altered; >2 m thick.
 - O¹** **Bog peat:** sphagnum or forest peat formed in an ombrotrophic environment; wet terrain; may be tread or treeless; O¹h, hummocky, mounds and plateaus; area may be underlain by ground ice or shallow permafrost conditions; O¹h, thermokarst terrain related to melting ground ice.
 - O²** **Fen peat:** peat derived from sedges and partially decayed shrubs in an eutrophic environment; forms relatively open peatlands with a mineral-rich water table that persists seasonally near the surface; generally covered with low shrubs and an occasional sparse layer of trees.
 - O** **Undifferentiated bog and fen deposits:** O¹, undifferentiated hummocky bog and fen deposits; area may be underlain by ground ice or shallow permafrost conditions; O¹, undifferentiated bog and fen deposits with thermokarst terrain related to melting of ground ice; O¹, undifferentiated bog and fen deposits with numerous subparallel channels on gentle slopes.
 - COLLUVIAL DEPOSITS:** mass wasting debris; poorly sorted, massive to stratified debris deposited by direct, gravity-induced movement; composition dependent on source material.
 - Ch** **Landslide and slump debris:** active and inactive landslides; hummocky topography; diamictic, generally 1 to 10 m thick, but may exceed 10 m near the toe of large landslides.
 - Cv** **Colluvial veneer:** thin and discontinuous cover of slumped and/or soliflucted material <1 m thick; overlies bedrock or till.
 - C** **Undifferentiated colluvial deposits.**
 - ALLUVIAL DEPOSITS:** sorted gravel, sand, minor silt, and organic debris deposited by streams; commonly stratified.
 - Ap** **Floodplain deposits:** sorted gravel, sand, silt, and organic debris >1 m thick; forming active floodplains close to river level with meander channels and scroll marks.
 - At** **Fluvial terrace deposits:** inactive terraces above modern floodplain; >2 m thick; represents a potential aggregate source.
 - Ad** **Deltaic sediments:** stratified sand and gravel underlain by silt and clay; generally 2 to 15 m thick; occurring at the mouths of streams entering lakes.
 - Af** **Alluvial fan deposits:** poorly sorted gravel, sand, and organic debris >1 m thick.
 - Av** **Alluvium veneer:** <1 m thick; primarily as uniform sheets of slope wash on gentle slopes.
 - A** **Undifferentiated fluvial deposits.**
 - L¹** **LACUSTRINE DEPOSITS:** sand, silt, and minor clay deposited in a former lake; >1 m thick; generally overlain by organic deposits; exposed by recent fluctuations in lake levels.


- NONGLACIAL AND PROGLACIAL ENVIRONMENTS**
- Eolian DEPOSITS:** wind-deposited medium to fine sand; derived from detritic or glaciolacustrine deposits; in some areas eolian sediments are thin or absent between dunes.
 - Er** **Ridged eolian deposits:** forming dunes; generally >2m thick.
 - Ev** **Eolian veneer:** discontinuous veneer of eolian sediments; <1 m thick.

- POSTGLACIAL OR LATE WISCONSINAN PROGLACIAL AND GLACIAL ENVIRONMENTS**
- GLACIOLACUSTRINE DEPOSITS:** fine sand, silt, and clay, with minor debris-flow diamictic, deposited in glacier-dammed lakes in valleys and along the margin of the retreating Laurentide Ice Sheet; usually overlain by organic deposits in lowlands.
 - Lb** **Glaciolacustrine blanket:** >1 m thick.
 - Lv** **Glaciolacustrine veneer:** thin and discontinuous; <1 m thick.

- GLACIOFLUVIAL DEPOSITS:** well to poorly stratified sand and gravel; minor diamictic; deposited behind, at or in front of the ice margin by glacial meltwater; represents a potential aggregate source.
- G** **Proglacial outwash:** cross-stratified gravel and sand deposited in front of the ice margin; G₁, outwash plain deposits, generally 1 to 5 m thick; generally marlly valley floors as well as surface adjacent to glacial meltwater channel margins; G₂, outwash terrace deposits, generally associated with meltwater channels and canyons; 1 to 10 m thick; G₃, glaciolacustrine delta deposits; 1 to >30 m thick; G₄, glaciolacustrine fan deposits; 1 to 10 m thick.
- Gi** **Ice-contact stratified drift:** poorly-sorted sand and gravel with minor diamictic; deposited in contact with the retreating glacier; 1 to >20 m thick; Gi₁, hummocky topography relating to melting of underlying ice; Gi₂, surface marked by kettle holes; Gi₃, esker ridges; Gi₄, kame terraces; Gi₅, ice-contact glaciolacustrine delta deposits; 1 to >30 m thick, surface marked by kettles.
- TILL:** diamictic deposited directly by the Laurentide Ice Sheet; sandy to clayey matrix with stratified clasts of various lithologies, including many Canadian Shield, carbonate and sandstone erratics; clast content is typically low (<10 %).

- Tb** **Till blanket:** >1 m thick, continuous till cover forming undulating topography that locally obscures underlying units.
- Ts** **Streamlined and fluted till:** >1 m thick, till surface marked by streamlined landforms including flutes and drumlins.
- Th** **Hummocky till:** >1 m thick; hummocky till surface.
- Tr** **Ridged till deposits:** >1 m thick, moraines or crevasse fillings forming a ridged topography.
- Tv** **Till veneer:** <1 m thick, discontinuous till cover, underlying bedrock topography is discernible.

- PRE-QUATERNARY BEDROCK**
- R** **Sedimentary bedrock:** Cretaceous Fort St. John Group shales (including the Shalebury Formation) and Devonian Formation sandstones exposed in highlands and along meltwater channel and canyon walls.



OPEN FILE DOSSIER PUBLIC 5307

GEOLOGICAL SURVEY OF CANADA / COMMISSION GÉOLOGIQUE DU CANADA

2009


Open file products that have not gone through the GSC formal publication process.

Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la GSC.



Author: I.R. Smith
 Geology by I.R. Smith, 2003-2005
 Geological completion by I.R. Smith, 2003-2006
 Digital completion by L. Robertson, GSC Northern Canada Division, 2006
 Digital cartography E. Everett, Data Dissemination Division (DDD)

GSC OPEN FILE 5307
 BCMEMPR MAP 2006-1
 SURFICIAL GEOLOGY
KYKLO CREEK
 BRITISH COLUMBIA



This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (DDD) Quality Management System, registered to the ISO 9001:2000 standard.

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada.

Digital base map provided by I.R. Smith, modified by DDD

Shaded relief image prepared by DDD, derived from the digital elevation model supplied by I.R. Smith, based on 3" arc second SRTM imagery. Illumination: azimuth 310°, altitude 25°, vertical factor 4.8x

Mean magnetic declination 2009, 21°01'E, decreasing 22.1' annually