

LEGEND
This legend is common to the GSC maps that are shown in Figure 1 below.
Coloured legend blocks indicate map units that appear on this map only.

QUATERNARY SURFICIAL DEPOSITS

NONGLACIAL ENVIRONMENT

- ORGANIC DEPOSITS:** peat, muck; 1 to 3 m thick; commonly underlain by fine-grained glacial lake deposits; occurs in confined, low-lying, poorly drained parts of the shield.
- 7b** Fan peat: wet ridge and moss peat; includes string fan, floodplain, and riverine terraces; occurs as flat grassy surfaces with few trees, and commonly visible surface waters; permafrost present in isolated plateaus or small peat plateaus which occur within the fan.
- 7a** Bog peat: moss and woody peat; occurs as raised irregular surfaces with an open to closed tree cover; derived from spruce forest vegetation; thermokarst depressions and ponds, wooded plateaus and forested peat plateaus are common; contains some areas of collapse scar fans.

- 6** ALLUVIAL DEPOSITS: silt, sand, and minor gravel, 1 to 30 m thick; floodplain and channel-fill sediments deposited in modern drainage ways.

PROGLACIAL ENVIRONMENT

- 5c** GLACIAL LAKE DEPOSITS: massive to stratified clay, silt, sand, and gravel; thickness ranges from a thin veneer to tens of metres; glacial sediments reworked by wave action in glacial Lake Agassiz, or carried to the basin in large part by glacial meltwater and deposited in deep water of Lake Agassiz.
- 5b** Offshore sediment blanket: clay, silt and silty sand, minor sand, gravel, and diamicton; fine-grained deep water sediments are non to weakly calcareous and commonly massive near surface; 2 to 45 m thick; form flat plains in low relief areas, mantled with peat; underlain by till or bedrock.
- 5a** Offshore sediment veneer: clay, silt, and silty sand; less than 2 m thick; forms a discontinuous glaciolacustrine sediments deposited in deep water beyond or near the ice margin; surfaces locally inscribed by ledging scour.

GLACIAL ENVIRONMENT

- GLACIOFLUVIAL DEPOSITS:** stratified sand and gravel, minor diamicton; sorted coarse-grained sediment deposited by flowing glacial meltwater in contact with or near the glacier.
- 4** Subaqueous outwash sediments: well sorted fine sand, commonly rippled and/or crossbedded; interbedded with clay, gravel, and diamicton units of variable thicknesses; 1 to 20 m thick; deformation and faulting common; deposits occur as outwash fans or down-ice of bedrock highs; sediments deposited in glacial Lake Agassiz at or near the retreating ice front by meltwater turbidity currents.
- 3** Proximal glaciofluvial sediments: moderately to well sorted and well rounded interstratified sand and gravel, minor diamicton; 3 to 25 m thick; forming eskers and crevasse fillings; deposited by sub- or epiglacial meltwater streams.
- 2** Till blanket: forms a continuous cover, 2 to several metres thick, locally up to 20 m thick on unchanneled landforms, masking underlying bedrock topography; deposits form drumlinized till plain and minor De Geer moraines.
- 1** Till veneer: forms a moderately discontinuous cover, 1 to 2 m thick, reflecting underlying bedrock structure; commonly thinsens on the down-ice side of Precambrian bedrock outcrops; surfaces may be covered by a thin veneer of Lake Agassiz offshore sediments or littoral sand and gravel.

PRE-QUATERNARY BEDROCK

- R** Precambrian rocks: metacarbonate and metasedimentary rocks, associated intrusive bodies; glacially scoured outcrops forming abundant rocks moutonnées and striated or grooved surfaces; gently rolling topography with thin patchy drift cover.

- Geological boundary (defined)
- Thermokarst depression, large
- Thermokarst depression, small
- Meltwater channel, large
- Meltwater channel, small (direction unknown, known)
- Drumlin, drumlinoid ridge, fluting, undifferentiated
- Crag-and-tail landform
- Palaeo
- Kettle
- Striae (ice flow direction known, unknown)
- Striae (poorly defined; ice flow direction known, unknown)
- Crossed striae (numbers indicate relative age, 1 being the oldest)
- Small bedrock outcrop
- Gravel pit
- Till sample

REFERENCES

Henderson, P.J., McMartin, L., Hall, G.E.M., Perivel, J.B., and Walker, D.A. 1990: The chemical and physical characteristics of heavy metals in humus and fill in the vicinity of the base metal smelter at Fim Flon, Manitoba, Canada. *Environmental Geology*, 34, p. 35-68.

McMartin, L., Henderson, P.J., and Nielsen, E. 1999: Impact of a base metal smelter on the geochemistry of soils of the Fim Flon region, Manitoba and Saskatchewan. *Canadian Journal of Earth Sciences*, 36, p. 141-160.

McMartin, L., Henderson, P.J., Nielsen, E., and Campbell, J.E. 1998: Surficial geology, till and humus composition across the Shield Margin, northwestern Manitoba and Saskatchewan: peatland analysis of a glaciated environment. *Geological Survey of Canada, Open File 3277*.

McMartin, L., Henderson, P.J., Pleuflin, A., and Knight, R.D. 2002: Comparison of Cu, Ni, Pb, Zn concentrations in soils adjacent to anthropogenic point sources: examples from four Canadian sites. *Geochimica et Cosmochimica Acta*, 66, p. 37-74.

Nielsen, E. 1982: Surficial geology and till geochemical sampling in the Neosap Lake area (NTS 68K14); in Manitoba Energy and Mines, Report of Activities 1980, Manitoba Energy and Mines, p. 47-49.




Figure 1. Index map showing the location of the Naosap Lake map and surrounding surficial geology maps published within the NATMAP Shield Margin Project area (grey outline). Sources used in the compilation are as follows: 1757A, Kazuyuki, C.A. and Way, New, V.J., 1:250 000; 1919A, Campbell, J.E. and Henderson, P.J., 1:50 000; 1954A, McMartin, L., 1:100 000; 2010A, Henderson, P.J., 1:50 000; OF285, McMartin, L., 1:100 000; OF300, McMartin, L. and Boucher, R., 1:100 000; OF310, McMartin, L., Campbell, J.E., and Boucher, R., 1:100 000; OF324, McMartin, L., 1:100 000; OF345, McMartin, L., 1:100 000; OF346, McMartin, L., 1:100 000; OF348, Campbell, J.E., McMartin, L., and Millard, M., 1:100 000; OF356, McMartin, L., 1:100 000; OF366, Campbell, J.E., Millard, M., and McMartin, L., 1:100 000; OF368, Henderson, P.J. and McMartin, L., 1:50 000; OF369, Henderson, P.J. and McMartin, L., 1:50 000.

Authors: P.J. Henderson and I. McMartin

Geology by C.A. Kazuyuki (formerly at Geological Survey of Canada) as part of the Canada-Manitoba Mineral Development Agreement, 1984-1989, by E. Nielsen (formerly at Manitoba Energy and Mines) and I. McMartin as part of the Shield Margin NATMAP Project, 1991-1994, and by P.J. Henderson and I. McMartin as part of the TGI-3 Fim Flon Project, 2008

Geological compilation by P.J. Henderson and I. McMartin, 2008

Digital compilation by Gléboch and L. Robertson, GSC Northern Canada Division, 2008

Digital cartography by J. Dohar, Data Dissemination Division (DDD)

Co-ordinated through the auspices of the TGI-3 Fim Flon Project under the Targeted Geoscience Initiative-3 of Natural Resources Canada

This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (DDG) Quality Management System, registered to the ISO 9001:2000 standard

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by DDD

Shaded relief image prepared by DDD, derived from the digital elevation model supplied by GSC Northern Canada Division
Illumination: azimuth 315°, altitude 45°, vertical factor 5x

Mean magnetic declination 2009, 7°21' E, decreasing 12.3" annually

Elevations in feet above mean sea level



OPEN FILE 5940
SURFICIAL GEOLOGY
NAOSAP LAKE
MANITOBA
Scale 1:50 000 / Échelle 1/50 000

Kilometres 0 1 2 3 4 Kilomètres

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2009

Projection transversale universelle de Mercator
Système de référence géodésique nord-américain, 1983
© Sa Majesté la Reine du Chef du Canada 2009

63 N4	63 N0	63 N2
63 K13	63 K14	63 K15
63 K12	63 K11	63 K10

OPEN FILE
DOSSIER PUBLIC
5940
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
2009

Open file are products that have not gone through the GSC formal publication process.

Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la GSC.