



Authors: P.J. Henderson and I. McMartin

Geology by E. Nielsen (formerly Manitoba Energy and Mines), C.A. Kaszycki (formerly Geological Survey of Canada) and I. McMartin, 1986-1990-1992; and by P.J. Henderson and I. McMartin, 2007-2009

Geological compilation by P.J. Henderson and I. McMartin, 2008-2009

Digital compilation by L. Robertson, GSC Northern Canada Division, 2008-2010

Co-ordinated through the auspices of the TGI-3 Flin Flon Project under the Targeted Geoscience Initiative-3 of Natural Resources Canada

Digital cartography by E. Everett, Data Dissemination Division (DDD)

OPEN FILE 6556
SURFICIAL GEOLOGY
FILE LAKE
MANITOBA

Scale 1:50 000/Échelle 1:50 000

Universal Transverse Mercator Projection
North American Datum 1983
© Her Majesty the Queen in Right of Canada 2010

Projection transversale universelle de Mercator
Système de référence géodésique du Nord-Américain 1983
© Sa Majesté la Reine du chef du Canada 2010

This map was produced from processes that conform to the Scientific and Technical Publishing Services Subdivision (DDD) Quality Management System, registered to the ISO 9001:2000 standard.

Any revisions or additional geological information known to the user would be welcomed by the Geological Survey of Canada

Digital base map from data compiled by Geomatics Canada, modified by DDD

Shaded relief image prepared by DDD, derived from the digital elevation model supplied by L. Robertson, based on the TRIM topographic data
Illumination: azimuth 315°, altitude 45°, vertical factor 2x

Magnetic declination 2010, 6°12'E, decreasing 11.0' annually

Elevations in feet above mean sea level

63 N2	63 N1	63-04
63 K15	63 K16 OF6556	63 J13 OF5889
63 K10	63 K9	63 J12

NATIONAL TOPOGRAPHIC SYSTEM REFERENCE AND NOTES
TO ACCOMPANY GEOLOGICAL SURVEY OF CANADA MAPS

LEGEND

This legend is common to the GSC maps that are shown in Figure 1 below.
Coloured legend blocks indicate map units that appear on this map.

SURFICIAL DEPOSITS QUATERNARY

NONGLACIAL ENVIRONMENT

ORGANIC DEPOSITS: peat, muck; 1 to 3 m thick; commonly underlain by fine-grained glacial lake deposits; occurs in confined, low-lying, poorly drained parts of the Shield.

7b Fan peat: wet sedge and moss peat; includes string fen, floodplain, and riverine marshes; occurs as flat grassy surfaces with low trees, and commonly visible surface waters; permafrost present in isolated patches or small peat plateaus which occur within the fen.

7a Bog peat: moss and woody peat; occurs as raised irregular surfaces with an open to closed tree cover; derived from spruce forest vegetation; thermokarst depressions and ponds, wooded peats and forested peat plateaus are common; contains some areas of collapse scar fens.

6 ALLUVIAL DEPOSITS: silt, sand, and minor gravel, 1 to 30 m thick; floodplain and channel-fill sediments deposited in modern drainage ways.

PROGLACIAL ENVIRONMENT

GLACIAL LAKE DEPOSITS: massive to stratified clay, silt, sand, and gravel; thickness ranges from a thin veneer to tens of metres; glacial sediments reworked by wave action in glacial Lake Agassiz, or carried to the basin in large part by glacial meltwater and deposited in deep water of Lake Agassiz.

5c Nearshore and littoral sediments: sand, gravel, or rock shingle, moderately well sorted and commonly horizontally bedded; occurs as isolated or series of ridges, 1 to 3 m in height, including beaches, bars, and spits; blankets of sand, commonly less than 2 m thick, grade basinward into finer sediments.

5b Offshore sediment blanket: clay, silt and silty sand, minor sand, gravel, and diamicton; fine-grained deep water sediments are non to weakly calcareous and commonly massive near surface; 2 to 45 m thick; form flat plains in low relief areas, marked with peat, surfaces locally inscribed by iceberg scours.

5a Offshore sediment veneer: clay, silt, and silty sand, less than 2 m thick; forms a discontinuous blanket mimicking underlying glacial and bedrock topography; includes undifferentiated glaciolacustrine sediments deposited in deep water beyond or near the ice margin; surfaces locally inscribed by iceberg scours.

GLACIAL ENVIRONMENT

GLACIOFLUVIAL DEPOSITS: stratified sand and gravel, minor diamicton; sorted coarse-grained sediment deposited by flowing glacial meltwater in contact with or near the glacier.

4 Subaqueous outwash sediments: well sorted fine sand; commonly rippled and/or cross-bedded; interbedded with clay, gravel, and diamicton; units of variable thicknesses; 1 to 20 m thick; deformation and faulting common; deposits occur as outwash fans or down-ice of bedrock highs; sediments deposited in glacial Lake Agassiz at or near the retreating ice front by meltwater turbidity currents.

3 Proximal glaciofluvial sediments: moderately to well sorted and well rounded interstratified sand and gravel, minor diamicton; 1 to 25 m thick; forming eskers and crevasse fillings; deposited by sub- or englacial meltwater streams.

GLACIAL DEPOSITS: unsorted to poorly sorted diamictons deposited at the ice margin or beneath the glacier. The area has been glaciated by ice originating from two dispersal centres, the Keweenaw Sector to the north, and the Labrador Sector to the east. Deposits have been subdivided into two units based on provenance: a) till of northern provenance overlying Precambrian rocks is generally sandy, permeable, non to slightly calcareous, and locally derived; b) till of eastern provenance is silty sandy, weakly permeable, moderately to strongly calcareous, and contains fewer Shield clasts.

2a Till blanket: forms a continuous cover, 2 to several metres thick, locally up to 20 m thick in streamlined landforms, masking underlying bedrock topography; deposits form drumlinized till plain and minor de Geer moraines; 2a - till of northern provenance underlain by Precambrian rocks; 2b - till of eastern provenance.

2b

1a Till veneer: forms a moderately discontinuous cover, 1 to 2 m thick, reflecting underlying bedrock structure; commonly occurs on the down-ice side of Precambrian bedrock outcrops; surface may be covered by a thin veneer of Lake Agassiz offshore sediments or littoral sand and gravel; 1a - till of northern provenance underlain by Precambrian rocks; 1b - till of eastern provenance.

1b

PRE-QUATERNARY BEDROCK

R Precambrian rocks: metavolcanic and metasedimentary rocks, associated intrusive bodies; glacially scoured outcrops forming abundant roches moutonnées and stratified or grooved surfaces; gently rolling topography with thin patchy drift cover.

Geological boundary (defined)
Thermokarst depression (large)
Thermokarst depression (small)
Drumlin, drumlinoid ridge, fluting; undifferentiated
Striae (ice flow direction known, unknown, unknown poorly defined)
Rock escarpment
Mine
Quarry
Small bedrock outcrop
Till sample

REFERENCES

Gober, G. and Nielsen, E., 1991. Till geochemistry of the Snow Lake-File Lake area (NTS 63016, J13); in Manitoba Energy and Mines, Report of Activities 1991, p. 47-48.

Kaszycki, C.A., Nielsen, E., and Gober, G., 1996. Surficial geochemistry and response to volcanic-hosted massive sulphide mineralization in the Snow Lake region; in G.F. Borhan-Carter, A.G. Galley, and G.E.M. Hall (eds.), EXTECH 1: A Multidisciplinary Approach to Massive Sulphide Research in the Rusty Lake-Snow Lake Greenstone Belts, Manitoba, Geological Survey of Canada, Bulletin 426, p. 139-154.

McMartin, I., 1994. Ice flow events in the Comond Lake-Veluxo Lake area, Northern Manitoba; in Current Research, Part C, Geological Survey of Canada, Paper 94-1C, p. 175-182.

McMartin, I., Henderson, P.J., Nielsen, E., and Campbell, J.E., 1996. Surficial geology, till and humus composition across the Shield Margin, north-central Manitoba and Saskatchewan: geospatial analysis of a glaciated environment, Geological Survey of Canada, Open File 3277.

Mhyzhuk, M., 1988. Surficial geology and aggregate resources in the Snow Lake area; in Report of Activities, Manitoba Energy and Mines, Map 1988 SL-2 and SL-3, 1:50 000 scale.

Figure 1. Index map showing the location of the File Lake map and surrounding surficial geology maps published within the NatMap Shield Margin Project area (grey outline). Sources used in the compilation are as follows: 1757A, Kaszycki, C.A. and Way Nee, V.J., 1:250 000; 1919A, Campbell, J.E. and Henderson, P.J., 1:50 000; 1954A, McMartin, I., 1:100 000; 2010A, Henderson, P.J., 1:50 000; OF2855, McMartin, I., 1:100 000; OF2860, McMartin, I. and Boucher, R., 1:100 000; OF2100, McMartin, I., Campbell, J.E., and Boucher, R., 1:100 000; OF3324, McMartin, I., 1:100 000; OF3342, McMartin, I., 1:100 000; OF3406, McMartin, I., 1:100 000; OF3485, Campbell, J.E., McMartin, I., and Milard, M., 1:100 000; OF3535, McMartin, I., 1:100 000; OF3565, Campbell, J.E., Milard, M., and McMartin, I., 1:100 000; OF5889, Henderson, P.J. and McMartin, I., 1:50 000; OF5828, Henderson, P.J. and McMartin, I., 1:50 000; OF5940, Henderson, P.J. and McMartin, I., 1:50 000.

OPEN FILE DOSSIER PUBLIC 6556

Geological Survey of Canada
COMMISSION GÉOLOGIQUE DU CANADA
2010

Open files are products that have not gone through the GSC formal publication process.

Les dossiers publics sont des produits qui n'ont pas été soumis au processus officiel de publication de la GSC.

Recommended citation:
Henderson, P.J. and McMartin, I., 2010. Surficial geology, File Lake, Manitoba; Geological Survey of Canada, Open File 6556, scale 1:50 000.