

Figure 1. Two ancient coastal cirques with floors below sea level, Leopold Island. These cirques probably formed during periods of low eustatic sea levels (glaciations) when continental ice sheets did not reach this area. 2011-057

Figure 2. Plateaux and fiords near Cape Mercy. 2011-059

Abstract

In 2009, as part of the GEM Program of the Geological Survey of Canada, Cumberland Peninsula east and southeast of the National Park was mapped. Mapping included several aspects of the regional Precambrian geology and the Quaternary geology. Regional till was sampled for sedimentological and geochemical purposes, and moraines and other deposits were sampled for cosmogenic exposure dating.

The eastern part of the peninsula is today an area of intensive alpine glaciation and evidently has experienced this style of glaciation throughout the Quaternary. During the last glacial maximum (28–12ka), alpine glaciers thickened to form regional ice divides over the mountains. Ice flow from these divides covered most, possibly all, of the region and supplied ice streams along several fiords. This local ice coalesced with the Laurentide Ice Sheet in Cumberland Sound. Deposits and landforms associated with this glaciation also show little or no sign of glacial erosion and retain Tertiary surfaces mantled with block fields and tors.

Résumé

En 2009, dans le cadre du programme GEM, la Commission géologique du Canada a cartographié les régions de la péninsule de Cumberland à l'est et au sud-est du parc national. La cartographie incluait plusieurs aspects de la géologie régionale précambrienne de même que la géologie du Quaternaire. Des échantillons de till régional ont été effectués à des fins d'analyses sédimentologiques et géochimiques ainsi que des moraines et d'autres dépôts à des fins de détermination des âges d'exposition aux rayonnements cosmogéniques.

La partie est de la péninsule est aujourd'hui une région de glaciation alpine intense et de toute évidence, ce type de glaciation existait durant le Quaternaire. Durant le dernier maximum glaciaire (28–12ka), les glaciers alpins se sont épaissis pour former des lignes de partage glaciaire au-dessus des montagnes. L'écoulement glaciaire provenant de ces lignes de partage glaciaire couvrait une grande partie sinon toute la région et formait des courants glaciaires en bordure de plusieurs fjords. Cette glace locale entrait en coalescence avec l'Inlandsis laurentidien dans le bas Cumberland. Les dépôts et les formes de relief associés à cette glaciation ne montraient que très peu ou même aucun signe d'érosion glaciaire et retenaient les surfaces du Tertiaire recouvertes de champs de blocs et de tors.



National Topographic System reference and index to adjoining published Geological Survey of Canada maps

Cover illustration
Ulvaalik Glacier, Cumberland Penninsula.
Photograph by Art Dyke.
2003-255

© Her Majesty the Queen in Right of Canada 2011

Printed map:
Catalogue No. M183-1/15-2010E
ISBN 978-0-660-20018-7

Digital map:
Catalogue No. M183-1/15-2010E-PDF
ISBN 978-1-100-17456-3

doi:10.4095/287205

Natural Resources Canada | Ressources naturelles Canada

CANADIAN GEOSCIENCE MAP 15
(preliminary version)
SURFICIAL GEOLOGY
HOARE BAY–ABRAHAM BAY NORTH
Baffin Island, Nunavut
1:100 000

Author: A.S. Dyke

Geology by A.S. Dyke, 2009

Geological compilation by A.S. Dyke, 2009

Cartography by L. Robertson

Initiative of the Geological survey of Canada, conducted under the auspices of the Eastern Arctic Cumberland Peninsula Project (Geo-mapping for Natural Resources and Development) and the Energy and Minerals (GEM) program

Logistical support provided by the Polar Continental Shelf Project as part of its mandate to promote scientific research in the Canadian North. PCSP 002-09

CANADIAN GEOSCIENCE MAP 15
SURFICIAL GEOLOGY
HOARE BAY–ABRAHAM BAY NORTH
Baffin Island, Nunavut
1:100 000

2 0 2 4 6 8 km

Map projection Universal Transverse Mercator, zone 20. North American Datum 1980

Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations in feet above mean sea level

Mean magnetic declination 2011, 32°45'W, decreasing 27.6' annually. Readings vary from 32°15'W in the SW corner to 33°52'W in the NE corner of the map.

The Geological Survey of Canada welcomes corrections and additional information from users. You may write to us by mail or e-mail. 2009 standard used by the Standards Council of Canada's Publishing Services Quality Management System.

This publication, including digital data, can be downloaded free of charge from GeoPub (http://geopub.nrcan.gc.ca/). It is also available from the Geological Survey of Canada Bookstore (http://gsc.nrcan.gc.ca/bookstore).

SURFICIAL DEPOSITS

QUATERNARY

HOLOCENE

NEOGLACIAL

Ice: glacier ice, 1–600 m thick forming cold-based and polythermal plateau ice caps and cirque and valley glaciers; extent as of AD 1958 with AD 2008 extent superimposed.

Till: nonsorted glacial debris commonly very bouldery with a silty sand matrix

Latero-frontal moraines: 5–100 m high moraine ridges with over-steepened, failing slopes on shallowly buried glacier ice cores and associated ground ice-cored terrain; mainly fresh looking deposits, distinguished from older moraines by lack of mature hi/hm and plant covers.

POSTGLACIAL (including Neoglacial)

FLUVIAL SEDIMENTS: alluvium; gravel and sand deposited beyond primary influence of Holocene glaciers

Alluvial fans: gravel and sand commonly bouldery, with detrital organic layers and buried soils, 1–20 m thick, formed by steep-gradient streams and debris flows forming channels and levees.

Alluvial plains: gravel and sand, 1–10 m thick.

Alluvial terraces: gravel and sand, 1–10 m thick, above limit of modern flooding.

COLLUVIUM: block and rubble accumulations, 1–50 m thick

Colluvial fan deposits: blocky to gravelly debris-flow accumulations mixed with scree, 1–50 m thick, traversed by narrow channels and debris levees; typically interrupt scree slopes (Cs) at the exits of prominent debris hopper in cliff surface slopes less than angle of repose but steeper than those of alluvial fans.

Talus: generally active accumulations of blocks and rubble, as much as 50 m thick forming talus (scree) aprons at angle of repose below cliffs derived from rock falls.

Rock glacier debris: talus, generally 10–50 m thick, deformed by interstitial flow of buried ice to form talus glaciers; irregular terraces on valley floors and lower valley sides; mainly fresh looking, some relict forms, some risers stable and well vegetated; most risers instable, unvegetated, and at angle of repose.

MARINE SEDIMENTS: gravel, sand, silt, and minor clay, 1–20 m thick, deposited on beach, bar, and offshore environments during regression of postglacial sea

Beach sediments: gravel and sand, commonly bouldery, 1–5 m thick, forming raised beach ridges and swales and the modern, transgressive beach, a barrier beach in places.

Deltaic sediments: sand and gravel, typically overlying fine sand and silt bottomset beds, 5–20 m thick, forming raised terraces; terraces at marine limit formed at or near the ice margin.

EARLY HOLOCENE AND WISCONSINAN

GLACIOFLUVIAL SEDIMENTS: gravel and sand, 1–10m thick, deposited behind, at, and in front of the ice margin

Proglacial outwash: gravel and sand, 1–10 m thick, forming fan-shaped deposits.

Proglacial outwash: gravel and sand, 1–10 m thick, forming inactive braided plains.

Proglacial outwash: gravel and sand, 1–10 m thick, forming terraced deposits.

TILL: nonsorted bouldery diamictons, 1–40 m thick, deposited in subglacial and ice marginal environments, lithic composition generally reflecting underlying bedrock

Latero-frontal moraines: 5–40 m high ridges and hummocks comprised mainly of till probably overlying debris-rich glacier ice cores, forming lateral and end moraine ridges and less organized, hummocky accumulations formed during glacier recession; commonly over-steepened, failing slopes on ice-cored terrain; degradation of ice cores; matrix somewhat more sandy and less silty than till forming ground moraine; locally contains ice-contact stratified drift and outwash.

Rock-glacierized moraines: ice-cored end or lateral moraines; 5–40 m high; commonly with lobate form and hummocky surface; mainly fresh, low-if ice-debris mixture; till mixed in places with scree; commonly ice-cored as local base level for scree accumulation, hence difficult to distinguish from Cr in places; mainly stable inserts.

Till veneer: variously bouldery (10–60% cover; typically 20–40%) diamicton with silty sand matrix, 0.5–2 m thick and discontinuous; insufficiently thick to obscure relief of underlying bedrock.

Till blanket: variously bouldery (10–60% cover; typically 20–40%) diamicton with silty sand matrix, 2–10 m thick, sufficiently thick to obscure relief of underlying bedrock.

PRE-WISCONSINAN

Weathered bedrock: Residuum: bitaxmeuver, rock rubble, and gruss, 1–2+ m thick, mantling bedrock; formed by disintegration of Precambrian bedrock prior to Last Glacial Maximum but including sparse erratics and possibly morphologically degraded glacial deposits; commonly comprised of rock rubble in sandy silt and sand matrix, with typically 60–80% block cover; finer rubble and gruss on coarser, monzogranite of Cumberland Batholith, interrupted by tors 1–6 m high; bedrock surface shows tills or no sign of glacial scouring; hence retains its pre-Quaternary form; mainly on flat or gently graded, cliff-bounded, upland plateaux, but on moderately steep slopes in places; most surfaces probably covered by cold-based ice during one or more glaciations.

BEDROCK

PRE-QUATERNARY

ROCK: rock of various compositions and Precambrian ages; Cumberland Batholith comprised largely of monzogranite dominant west of Kingnait Fiord, lonalite and metasediments farther east

Cliffs: major escarpments, typically hundreds of metres high, forming serrated faces with multiple debris hoppers; glacially scoured surfaces removed by postglacial scarp retreat producing basal (or submarine) talus accumulations.

Scoured rock: hilly and hummocky surfaces with lake basins and ice moulded eminences resulting from glacial scouring and with patchy veneers of till, commonly depleted of matrix material, probably covered by warm-based ice during stadial intervals of Wisconsin Glaciation, including Last Glacial Maximum.

- Glacier ice margin for 2009 from SPOT5 satellite imagery
- Geological boundary (defined)
- Geological boundary (gradational)
- Fluted bedrock (direction known)
- Latero-frontal moraine
- Lateral meltwater channel, barb on upslope side
- Proglacial meltwater channel
- Cirque
- Arête
- Cliff
- Glacial lake limit
- Dyke
- Field observation (point colour relative to geological units)
- Striae (ice flow direction known)
- Striae from literature (Kaplan et al., 2001)
- Station, marine shoreline elevation in metres
- Station, marine shoreline elevation in metres from literature (Kaplan, 2001)

Recommended citation:
Dyke, A.S., 2011. Surficial geology, Hoare Bay–Abraham Bay north, Baffin Island, Nunavut; Geological Survey of Canada, Canadian Geoscience Map 15 (preliminary version), scale 1:100 000. doi:10.4095/287205

ess.nrcan.gc.ca

Canadian **Geoscience** Maps
Cartes **géoscientifiques** du Canada

Canada

Four trim marks around perimeter of map sheet. Trim map sheet first, then fold at folding marks.

Cover and additional panels are 175m wide when folded.

CANADIAN GEOSCIENCE MAP 15
(preliminary version)
SURFICIAL GEOLOGY
HOARE BAY–ABRAHAM BAY NORTH
Baffin Island, Nunavut