



**GEOLOGICAL SURVEY OF CANADA
OPEN FILE 5279**

Metamorphic map of the western Churchill Province, Canada

R. G. Berman

2010



Natural Resources
Canada

Ressources naturelles
Canada

Canada



**GEOLOGICAL SURVEY OF CANADA
OPEN FILE 5279**

**Metamorphic map of the western Churchill Province,
Canada**

This report contains marginal notes to accompany 3 map sheets.

R. G. Berman

2010

This publication is available from the Geological Survey of Canada Bookstore (http://gsc.nrcan.gc.ca/bookstore_e.php).
It can also be downloaded free of charge from GeoPub (<http://geopub.nrcan.gc.ca/>).

Recommended citation:

Berman, R.G., 2010. Metamorphic map of the western Churchill Province, Canada; Geological Survey of Canada, Open File 5279, 3 sheets, 1:2 500 000 scale, + 49 p. report.

Open files are products that have not gone through the GSC formal publication process.

TABLE OF CONTENTS

INTRODUCTION	1
Overview.....	1
Acknowledgements.....	2
METAMORPHIC MAP LEGEND	4
GEOLOGICAL BASE.....	6
DESCRIPTION OF METAMORPHIC EVENTS.....	8
Pre-2.56 Ga Archean metamorphic events	8
MacQuoid Orogeny (ca. 2.56 - 2.50 Ga)	9
Arrowsmith Orogeny (ca. 2.5 - 2.3 Ga)	10
Thelon-Taltson Orogeny (ca. 2.0 – 1.92 Ga)	14
Snowbird phase of Hudsonian Orogeny (ca. 1.91 – 1.87 Ga)	17
Main phase of Hudsonian Orogeny (ca. 1.86 – 1.75 Ga)	23
<u>Saskatchewan – Manitoba segment of the Trans-Hudson orogen</u>	23
<u>North Quebec – Baffin Island segment of the Trans-Hudson orogen</u>	26
<u>Central Rae segment of the Trans-Hudson orogen</u>	29
REFERENCES	34

LIST OF FIGURES

Figure 1: Regional geology of the western Churchill Province and environs	3
---	---

LIST OF TABLES

Table 1: References to geochronological data displayed on sheets 1-3.....	7
---	---

INTRODUCTION

1. Overview:

The western Churchill Province represents a large region of dominantly Meso- to Neo-Archean continental crust comprised of at least four major crustal blocks: Rae craton, Hearne craton, Chesterfield block, and Meta Incognita microcontinent (Fig. 1). Recent tectonic interpretation (Berman et al., 2007) suggests that the Rae and Chesterfield blocks were amalgamated at ca. 2.65 Ga (prior to the emplacement of ca. 2.6 Ga stitching plutons), whereas the Hearne craton accreted to the composite Rae-Chesterfield block at ca. 1.9 Ga, prior to the ca. 1.88 – 1.865 Ga collision of Meta Incognita microcontinent with the Rae craton (St-Onge et al., 2002).

The western Churchill Province has a complex metamorphic history with some regions affected by more than eight thermotectonic events between ca. 2.8 – 1.7 Ga. Not surprisingly, the most recent of these events is the most extensively preserved and the most reliably documented. A key tool in deciphering earlier metamorphic events has been *in situ* geochronology of metasedimentary rocks (e.g. Williams et al., 1999; Stern and Berman, 2000; Berman et al., 2005). Application of this technology both as an integral part of regional mapping studies and via reconnaissance studies has been successful in helping to define distinct regional events at ca. 2.56 – 2.50 Ga (MacQuoid orogeny; Stern and Berman, 2000), 2.5 – 2.3 Ga (Arrowsmith orogeny; Berman et al., 2005; Schultz et al., 2007; Hartlaub et al., 2007), 2.0 – 1.92 Ga (Thelon - Taltson orogeny; Hoffman, 1988; Bostock and van Breemen, 1992; Thériault, 1992; Henderson et al., 1999; De et al., 2000; McNicoll et al., 2000), 1.91 – 1.87 Ga (Snowbird phase of Hudsonian orogeny; Berman et al., 2007), and 1.86 – 1.75 Ga (main phase of Hudsonian orogeny; e.g. Hoffman, 1988; Lewry and Collerson, 1990; Ansdell, 1995; St-Onge et al., 2002;

Berman et al., 2005). In many cases, these geochronological data have been linked to metamorphic assemblages and/or calculated pressure - temperature conditions, making it possible to infer overall metamorphic conditions at specific times. However in very few places has pre-2.5 Ga metamorphism been dated convincingly in spite of the widespread presence of ca. 2.6 Ga and older basement. In the few places where ca. 2.6 Ga and older metamorphism has been dated (e.g. Skulski et al., 2003; Davis et al., 2004; Berman et al., 2005, 2008, 2010a; Mahan et al., 2006; Harper et al., 2007; Martel et al., 2007), the metamorphic grade is largely obscured by younger events. For these reasons no attempt has been made to construct a metamorphic map of pre-2.5 Ga metamorphism.

After a brief description of the map legend, the five main orogenic events depicted on Sheets 1-3 are briefly described.

2. Acknowledgements:

The author is greatly appreciative of the excellent cartographic support he has received especially from M. Methot, but also from P. Arscott, C. Hemingway, T. Houlahan, and D. Paul during the early stages of the work that led to this publication. I am also very grateful for the contributions made by a number of geologists at various stages in the compilation of the metamorphic zones portrayed in these regions: Saskatchewan (K. Ashton, C. Card, R. Maxheiner), Manitoba (D. Schledewicz, H. Zwanzig, T. Corkery, P. Gilbert), southern and central Baffin Island (M. St-Onge), and Thelon tectonic zone (P. Thompson). Lastly, I thank those who provided unpublished geochronology results (see Table 1), D. Corrigan for his constructive review, and Sylvia Frohberg-Junginger for her very careful cartographic check.




Figure 1. Regional geology of the western Churchill Province and environs

METAMORPHIC MAP LEGEND

The metamorphic map legend used on Sheets 1-3 follows the format described by Berman et al. (2000a) and applied elsewhere (Easton et al., 2005; Ashton et al., 2009). Increasing metamorphic grade is shown with generally warmer colours, except for granulite facies (purple). Where pressure information is available, the metamorphic pressure regime is shown by the orientation of superimposed coloured lines. Pressure is classified as high (vertical lines) or low-intermediate (horizontal lines), primarily on the basis of whether kyanite (Barrovian facies-series) or andalusite-sillimanite (Abukuma facies-series) is present in pelitic rocks. For granulite-facies rocks, a medium-pressure (diagonal lines) subdivision is also used, based on thermobarometric results.

Instead of showing complex overprinting relationships on a single map (e.g. Berman et al., 2000a; Easton et al., 2005), separate metamorphic time slices are presented on Sheets 1-3, as done on the metamorphic map of Saskatchewan (Ashton et al., 2009). On Sheets 1 and 2, two different ages of metamorphism are shown by means of a bright and a pastel colour series of metamorphic zones. In two small regions where separate metamorphic events overlap (Sheet 2), overprint “circles” are used to show the metamorphism of the younger event.

In order to convey the degree (or lack) of control on the age of assigned metamorphic zones, selected geochronological data (together with reference numbers given in parentheses) are shown on Sheets 1-3. Table 1 lists the reference that corresponds to each of these reference numbers. Because of spatial limitations on the maps, the displayed data do not represent a comprehensive compilation, but show instead the most constraining data available in each region. The majority of displayed data are U-Pb monazite and zircon ages that are interpreted as metamorphic, typically crystallizing near the metamorphic peak. Most ^{40}Ar - ^{39}Ar ages are not shown because they generally record younger cooling ages through the closure temperature of

hornblende (500 °C; Harrison, 1981) and biotite (300 - 350°C; Harrison et al., 1981). Selected ^{40}Ar - ^{39}Ar and K-Ar ages are used to show regions where metamorphic grade contrasts can be identified by differences in hornblende or biotite ages. In most of these regions, the preservation of old biotite or hornblende ages is used to indicate that these regions have experienced metamorphic grade lower than greenschist or lower amphibolite, respectively. Where other data are lacking, some K-Ar biotite or hornblende ages are also used to indicate regions that experienced at least greenschist- or amphibolite-facies metamorphism, respectively.

For those data that have not been previously published, the ages displayed on the maps (Sheets 1-3) are rounded to the nearest 5 Myr and associated uncertainties are not shown. The references for these geochronological data are presented in Table 1. An important point to note is that the distribution of the geochronological data provides a visual representation of the regions across which metamorphic zones have been interpolated or extrapolated. Areas of each map that are uncoloured represent either regions that were unmetamorphosed at the time of metamorphism shown on each map, or regions in which there are no data to evaluate whether they were metamorphosed.

Thermobarometric data are not shown on the metamorphic maps for several reasons. First, their display obscures metamorphic relationships shown on the maps, particularly where the data are most dense in parts of Saskatchewan and Manitoba. Second, unlike geochronological data which have very accurately known decay constants, thermobarometric data are based on empirical observations, sometimes conflicting sets of experimental data, and thermodynamic data that are not all accurately known (e.g. Berman, 1990; Spear, 1993). Third, numerical results are highly sensitive to interpretation of mineral zoning patterns (which are not always documented), and to whether results correspond to prograde, near-peak, or retrograde conditions.

GEOLOGICAL BASE

The metamorphic maps shown on Sheets 1-3 were compiled on a composite geologic base, simplified primarily from the following sources:

- Saskatchewan: Macdonald, 1999 (1:1,000,000 scale)
- Manitoba: Manitoba Geological Survey, 1979 (1:1,000,000 scale)
- Hearne/Rae craton (60-64 °N, 88-100 °W): Tella et al., 2007 (1:550,000 scale)
- Hearne/Rae craton (60-64 °N; 100-112 °W): Pehrsson et al., 2011 (1:550,000 scale)
- Rae craton (64-68 °N; 81-102 °W): Patterson and Lecheminant, 1985; Paul et al., 2002 (1:1,000,000), Wheeler et al., 1996 (1:5,000,000)
- Rae craton (68-70 °N; 81-86 °W): Schau, 1993 (1:500,000 scale)
- Rae craton (north of 68 °N) and Baffin Island: Wheeler et al., 1996 (1:5,000,000 scale), Fraser et al., 1978 (1:3,500,000 scale), Jackson and Berman (2000)

Table 1. References for geochronological data plotted on Sheets 1-3.

#	Reference	#	Reference
1	Annesley et. al. (1992a)	72	Heaman et. al. (1999)
2	Annesley et. al. (1992b)	73	Heaman et. al. (1993)
3	Annesley et. al. (1993)	74	Heaman et. al. (1991)
4	Annesley et. al. (1999)	75	Heaman et. al. (1994)
5	Annesley et. al. (1997)	76	Henderson et. al. (1992)
6	Ansdell and Stern (1997)	77	Hunt and Roddick (1987)
7	Ashton et. al. (1999a)	78	Hunt et. al. (1988)
8	Ashton et. al. (1999b)	79	Hunt et. al. (1993)
9	Ashton et. al. (1992)	80	Jackson (1978)
10	Aspler et. al. (2002)	81	Jackson et. al. (1990)
11	Baadsgaard and Godfrey (1967)	81a	Joyce, N. and Ryan, J. (2007, unpub. data)
12	Baldwin et. al. (2006)	82	Koster and Baadsgaard (1970)
13	Baldwin et. al. (2003)	82a	Leech et al. 1963)
14	Baldwin et. al. (2004)	83	Loveridge et. al. (1988)
15	Banks (1980)	84	Lowdon et. al. (1963)
16	Beaumont-Smith and Bohm (2002)	85	Machado et. al. (1990)
17	Bell (1985)	86	MacLachlan, K. and Davis, W. J. (2005, unpub. data)
18	Berman, R.G. (2005, unpub. data)	87	MacLachlan et. al. (2005a)
19	Berman, R.G., Ashton, K., Davis, W.J. (2005, unpub. data)	88	MacLachlan et. al. (2005b)
20	Berman, R.G., Card, C., Ashton, K., Davis, W.J. (2005, unpub. data)	89	MacLachlan et. al. (2004)
21	Berman et. al. (2002b)	90	Mahan et. al. (2006a)
22	Berman, R.G., Davis, W.J., LeCheminant, T. (2005, unpub. data)	91	Mahan et. al. (2006b)
23	Berman, R.G., Davis, W.J., Pehrsson, S. (2005, unpub. data)	92	Martel et. al. (2007)
24	Berman et. al. (2007)	93	McDonough et. al. (2000)
25	Berman et. al. (2002a)	94	McNicol et. al. (2000)
26	Berman, R.G., Davis, W.J., Sanborn-Barrie, M. (2005, unpub. data)	95	Mills et. al. (2007)
27	Berman, R.G., Davis, W.J., Sandeman, H.S. (2004, unpub. data)	96	Parent et. al. (1999)
28	Berman, R.G., Davis, W.J., Tella, S. (2004, unpub. data)	97	Parrish (1989)
29	Berman, R.G., Maxeiner, R., Davis, W.J. (2005, unpub. data)	98	Rayner et. al. (2005b)
30	Berman, R.G., Qui, H., Davis, W.J., Pehrsson, S. (2006, unpub. data)	99	Reilly (1993)
31	Berman, R.G., Rayner, N. and Sanborn-Barrie, M. (2006, unpub. data)	100	Roddick et. al. (1992)
32	Berman, R.G., Rayner, N., Sanborn-Barrie, M., Chakungal, J. (2009, unpub. data)	101	Roddick and van Breemen (1994)
33	Berman et. al. (2011)	102	Ryan et. al. (2000)
34	Berman, R.G., Ryan, J.J., Davis, W.J., Nadeau, L. (2008, unpub. data)	103	Sanborn-Barrie et. al. (2001)
35	Berman et. al. (2008)	104	Sandeman, H.A. and Berman, R.G. (2001, unpub. data)
36	Berman, R.G., Sanborn-Barrie, M., Rayner, N. (2003, unpub. data)	105	Sandeman (2001)
37	Berman et. al. (2010a)	106	Schultz et. al. (2007)
38	Berman et. al. (2005)	107	Scott, D. (2005, unpub. data)
39	Berman, R.G., Stern, R.A., Tella, S. (1997, unpub. data)	108	Scott and Gauthier (1996)
40	Bethune and Scammell (1997)	108a	Scott and St-Onge (1995)
41	Bethune and Scammell (2003)	109	Scott and Wodicka (1998)
42	Bickford et. al. (1994)	110	Scott (1997)
43	Bohm, C. (2006, unpub. data)	111	Scott (1999)
44	Bostock and van Breemen (1992)	112	Scott et. al. (2002)
45	Bostock et. al. (1991)	113	Shaerer and Deutsch (1990)
46	Bostock and Loveridge (1988)	114	Stern et. al. (2000)
47	Brouand et. al. (2002)	115	Stern et. al. (2003)
48	Carpenter et. al. (2005)	116	Stevens et. al. (1982a)
49	Carson et. al. (2004)	116a	Stevens et. al. (1982b)
50	Cavell et. al. (1992)	117	St-Onge et. al. (2002a)
51	Chiarenzelli et. al. (1998)	118	St-Onge et. al. (2002b)
52	Clark et. al. (1988)	119	St-Onge et. al. (2003a)
53	Clarke et. al. (2005)	120	St-Onge et. al. (2003b)
54	Corrigan et. al. (2001)	121	St-Onge et. al. (2003c)
55	Cranstone et. al. (1976)	122	St-Onge et. al. (2005a)
56	David et. al. (1996)	123	St-Onge et. al. (2005b)
57	Davis, W.J. (2005, unpub. data)	124	St-Onge et. al. (2005c)
58	Davis et. al. (2004)	125	St-Onge et. al. (2005d)
59	Davis et. al. (2006)	126	St-Onge et. al. (2007)
60	Fedorowich et. al. (1995)	127	ter Meer (2001)
61	Flowers et. al. (2006b)	128	Thomas (1994)
62	Flowers et. al. (2006a)	129	Tran (2001)
63	Flowers et. al. (2008)	130	van Breemen and Bostock (1994)
64	Frisch and Hunt (1988)	131	van Breemen et. al. (2007)
65	Frisch and Hunt (1993)	131a	Wanless et al. (1966)
66	Gordon et. al. (1990)	131b	Wanless et al. (1970)
67	Harper et. al. (2006)	131c	Wanless et al. (1972)
68	Hartlaub (2004)	131d	Wanless et al. (1974)
69	Hartlaub, R.P. (2006, unpub. data)	132	Wanless et. al. (1979)
70	Hartlaub et. al. (2005)	133	Williams and Jercinovic (2002)
71	Heaman, L.M. (2004, unpub. data).	134	Wodicka and Scott (1997)

refers to reference number given in parentheses with ages plotted on Sheets 1-3

DESCRIPTION OF METAMORPHIC EVENTS

The five main orogenic events depicted on Sheets 1-3 (ca. 2.56 – 2.50 Ga, ca. 2.50 – 2.30 Ga, ca. 2.0 – 1.92 Ga, ca. 1.91 – 1.87 Ga, ca. 1.86 – 1.75 Ga) are briefly described here with the aim of providing an overview of the main geochronological constraints, styles of metamorphism, and interpreted tectonic settings. For regions within Saskatchewan, most of the descriptions as well as the map linework and geochronological constraints are taken from Ashton et al. (2009). Also included below is brief mention of earlier metamorphic events that are not displayed on Sheets 1-3.

Pre-2.56 Ga Archean metamorphic events

Although there is no shortage of evidence of metamorphic events between 2.8 – 2.6 Ga, associated metamorphic assemblages are generally not known and the data are too dispersed geographically to be displayed meaningfully on a map. Regions with the most unambiguous evidence for specific, pre-2.56 Ga metamorphic events include: the high-grade, ca. 2.85 Ga metamorphism in the Chipman domain of the central Snowbird tectonic zone in northeastern Saskatchewan (Mahan et al., 2006) – southeastern Northwest Territories (Martel et al., 2007), ca. 2.66 Ga, largely greenschist-facies metamorphism of the Kaminak belt (Davis et al., 2004), a ca. 2.60 Ga granulite-facies belt on southern Boothia Peninsula (Hinchey et al., 2006; Berman et al., 2008), the cryptic, ca. 2.58 Ga, greenschist-facies regional contact metamorphism of the Committee Bay belt (Skulski et al., 2003; Berman et al., 2010a), and the ca. 2.7 Ga greenschist-to amphibolite-facies metamorphism of the Ege Bay region on Baffin Island (Bethune and Scammell, 2003).

MacQuoid Orogeny (ca. 2.56 - 2.50 Ga; pastel colour series on Sheet 1)

Evidence for this event was first discovered in the MacQuoid supracrustal belt within the Chesterfield block (Berman et al., 2000b; Stern and Berman, 2000). A strong, northwest-dipping, northeasterly-striking tectonic fabric is associated with low- to intermediate-P, middle amphibolite-facies metamorphism dated between 2.56 and 2.50 Ga (*in situ* monazite and zircon) in the MacQuoid Lake supracrustal belt (Berman et al., 2000b; Stern and Berman, 2000). Similar ages have been obtained in detailed studies of low-P, upper amphibolite to granulite facies rocks in domain 2 at Angikuni Lake (*in situ* U-Pb monazite; Berman et al., 2002b) and in the Yathkyed Lake region (K-Ar hornblende, Hunt and Roddick, 1987; ^{40}Ar - ^{39}Ar hornblende, Sandeman, 2001; U-Pb monazite, MacLachlan et al., 2005a; U-Pb titanite, MacLachlan et al., 2005a; W. J. Davis, unpub. data, 2005). This event is also documented in structurally deeper, granulite-facies rocks of the Snowbird tectonic zone, both to the southwest in in northwestern Saskatchewan (Baldwin et al., 2003; Mahan et al., 2006a, 2006b; Flowers et al., 2008) and to the northeast in the Big Lake shear zone (Ryan et al., 2000), Kramanituar complex (Sanborn-Barrie et al., 2001) and Uvauk complex (Mills et al., 2007). Scant geochronological data (see Sheet 1) suggest that this event extends into felsic gneisses further to the northeast of Chesterfield Inlet, but the grade of metamorphism there is unknown (assumed to be amphibolite facies), except for an upper amphibolite- to granulite-facies region mapped to the east of the Uvauk complex (Tella et al., 1993) and assumed here to be ca. 2.5 Ga in age.

High-grade rocks with ca. 2.55 – 2.53 Ga metamorphic ages have also been identified in northern Manitoba (C. Bohm, unpub. data, 2006) and southern Nunavut (Loveridge et al., 1988) as well as in the region of Peter Lake domain, Saskatchewan (Annesley et al., 1992a, b; R. G. Berman, R. Maxeiner, R., W. J. Davis, unpub. data, 2005). These metamorphic rocks are

considered to be unrelated to those discussed above because the Hearne craton is thought to have had a separate and independent history from the Rae craton until their collision at ca. 1.9 Ga (Berman et al., 2007).

A possible tectonic setting for the MacQuoid orogeny is that it initiated as a continental arc (northwest-dipping subduction under the southeast margin of the Rae - Chesterfield block) that generated a large volume of 2.61 – 2.58 Ga plutons in the Rae - Chesterfield block (e.g. van Schmus et al., 1986; LeCheminant and Roddick, 1991; Skulski et al., 2003; Davis et al., 2006; Hinchey et al. 2006), and caused low- to intermediate-*P* metamorphism at ca. 2.56 Ga in the MacQuoid supracrustal belt (Stern and Berman, 2000). A subsequent collisional event may have produced the higher pressure, more calcic outer core compositions and microporphyroblasts observed in garnet from the MacQuoid supracrustal belt and dated at ca. 2.50 Ga (Berman et al., 2000; Stern and Berman, 2000). If this model is correct, it implies that the colliding block rifted away between 2.2 – 2.1 Ga, prior to the ca. 1.9 Ga Snowbird phase of the Hudsonian orogeny (see below, and Sheet 2). The general lack of recognition of a ca. 2.5 Ga tectonic fabric in parts of the Rae craton outside of the Snowbird tectonic zone may reflect strain being taken up in the Rae-Chesterfield boundary zone, as documented in the granulite-facies Big Lake shear zone (Ryan et al., 2000).

Arrowsmith Orogeny (ca. 2.5 - 2.3 Ga; bright colour series on Sheet 1)

The Arrowsmith Orogeny was defined on the basis of ca. 2.35 Ga monazite interpreted to date a distinct deformation and metamorphic event in the western and northwestern parts of the Committee Bay belt (Berman et al., 2005). The clockwise P-T paths and absence of pre- or syn-metamorphic magmatism (Sanborn-Barrie et al., 2011) indicate that low-P (~ 4 kbar), lower amphibolite-facies metamorphism in this region was a response to crustal thickening (Berman et

al., 2005). Subsequent work to the west has revealed ca. 2.5 – 2.43 Ga felsic plutonism, sedimentation (Sherman gneisses), and ca. 2.38 Ga metamorphism (monazite growth) in low-*P* (~ 5 kbar, R. G. Berman, unpub. data), granulite-facies rocks of the eastern part of the Queen Maud block, south of Queen Maud Gulf (Schultz et al., 2007). On the southern part of Boothia Peninsula a cryptic, ca. 2.43 – 2.35 Ga high grade event has been documented in metasedimentary rocks that experienced granulite facies at ca. 2.6 Ga (Berman et al., 2008). North of this ca. 2.6 Ga granulite-facies belt, metasedimentary rocks record episodic, low-*P* (~ 5 kbar), at least upper amphibolite-facies monazite growth at ca. 2.5 Ga, ca. 2.43 Ga, and ca. 2.38 Ga, with the ca. 2.5 Ga monazite dating a regional tectonic fabric (R. G. Berman, W. J. Davis, J. J. Ryan, and L. Nadeau, unpub. data, 2008).

For large areas of the Rae craton, data are not available to evaluate the timing of metamorphism. Nevertheless much of the western and northern Rae craton is considered to have been affected by this orogenic event on the basis of widely distributed, but sparse geochronological data. In the northern Rae craton, this event is suggested by ca. 2.48, 2.4, and 2.32 Ga zircon ages of granulite-facies gneiss on northern Boothia peninsula (Frisch and Hunt, 1993), a ca. 2.52 Ga zircon age of granulite-facies gneiss on Devon Island (Frisch and Hunt, 1988), a ca. 2.53 Ga zircon age of enderbite on Bylot Island (northeast Baffin Island; D. Scott, unpub.data, 2005), and ca. 2.31 and 2.39 Ga Rb-Sr whole rock isochron ages and a ca. 2.5 Ga zircon age for upper amphibolite-facies gneiss on northwest Baffin Island (Jackson, 1978; Jackson et al., 1990). The above three age determinations on northern Baffin Island suggest that this region lies within the Arrowsmith orogen, and the 6 – 8 kbar pressures determined from these granulite-facies areas yield a consistent indication of low- to medium-*P* granulite facies metamorphism at this time. There is insufficient geochronological control to allow the southern

extent of Arrowsmith orogen on northern Baffin Island to be delineated from Hudsonian orogenic reworking. With the exception of the high-*P* northeast Baffin thrust zone (see Sheet 3), Hudsonian metamorphism is thought to decrease in grade northward from its culmination in the high-*P* Dexterity granulite belt on the north side of the Isortoq fault zone (Jackson and Berman, 2000; Sheet 3), an interpretation supported by the preservation of pre-Hudsonian Rb-Sr isochrons (ca. 1.97 Ga and 2.39 Ga, Jackson, 1978; ca. 2.31 Ga, Jackson et al., 1990) on northwestern Baffin Island. Thus upper amphibolite-facies regions of northwestern Baffin Island (Fraser et al., 1978) more likely represent an earlier metamorphic event, which is assumed here to be the Arrowsmith orogeny. This interpretation gains some support from a 2.52 Ga upper intercept zircon age from migmatitic gneiss near the tip of Cambridge Fjord (Jackson et al., 1990), which is used here as the southern limit of the Arrowsmith orogen, but it may well extend further south to the Isortoq fault zone, coincident with the southern limit of Jackson and Berman's (2000) Committee orogen. It should be noted that data are not available to evaluate the extent that northern Melville Peninsula may have been affected by the Arrowsmith orogeny. Granulite-facies plutonism and/or metamorphism on northwestern Melville Peninsula (Schau et al., 1993) is assumed to be ca. 2.6 Ga in age, based on its proximity to the 2.6 Ga granulite belt on southern Boothia Peninsula (Berman et al., 2008).

In the southwestern Rae craton, the Arrowsmith orogen is marked by low-*P* (~ 4 – 7 kbar), granulite-facies gneisses southwest of the Thelon basin (Sheet 3), which record ca. 2.5 – 2.3 Ga monazite ages (Berman, R.G., Qui, H., Davis, W.J., Pehrsson, S., unpub. data, 2006), and by upper amphibolite- to granulite-facies gneiss with 2.38 – 2.32 Ga monazite and zircon ages in the Taltson basement (Bostock and Loveridge, 1988; McNicoll et al., 2000). Granitic plutons of 2.33-2.29 Ga age in the Uranium City region of northwestern Saskatchewan are interpreted as a product of syn- to post-orogenic magmatism associated with the Arrowsmith orogeny (Hartlaub

et al., 2007). Rare ca. 2.35 Ga chemical monazite ages in variably overprinted pelitic gneisses probably record the Arrowsmith metamorphic event in this region (Hartlaub, 2004). Granitoid rocks north of Tazin Lake, Saskatchewan yield ca. 2.5 – 2.3 Ga K/Ar ages in hornblende (Koster and Baadsgaard, 1970; Banks, 1980), suggesting that peak temperatures there exceeded the $500 \pm 40^\circ\text{C}$ estimated closure temperature of hornblende (Harrison, 1981), thus reaching at least lower amphibolite facies. Beyond these inferences, the grade of metamorphism in northwestern Saskatchewan is largely unconstrained as is the extent of this event, which is currently based on known occurrences of the 2.33-2.29 Ga plutons (Hartlaub et al., 2007; Ashton et al., 2009).

“Arrowsmith” ages have also been found on the east side of the Rae craton in parts of the Snowbird tectonic zone, occurring in high-grade gneiss and greenschist-grade mylonite at Angikuni Lake (MacLachlan and Davis, unpub. data, 2005) and in upper amphibolite- to granulite-facies gabbro on the south side of the Kramanituar complex (Sanborn-Barrie et al., 2001). The ca. 2.45 Ga ages observed in the Sask craton exposed in structural windows to the south are considered to be unrelated to the Arrowsmith orogeny (Ashton et al., 2009).

The Arrowsmith orogeny is considered to represent the transition from a ca. 2.5 – 2.4 Ga continental arc (east-dipping subduction beneath the western flank of the Rae craton; e.g. Hoffman, 1988) to a ca. 2.4 – 2.3 Ga collisional orogen (Berman et al., 2005), and the evidence for ca. 2.5 Ga regional foliation formation in the Boothia Peninsula region (see above) support this model. However, Schultz et al. (2007) interpret their Queen Maud data to indicate an extensional environment, possibly a continental rift, with sedimentary basin formation (Sherman gneisses) penecontemporaneous with ca. 2.5 – 2.43 Ga plutonism. This region may thus represent a back-arc setting or relatively short-lived rift within an overall convergent tectonic setting.

Thelon-Taltson Orogeny (ca. 2.0 – 1.92 Ga; pastel colour series on Sheet 2)

While metamorphic zones and the Barrovian style of metamorphism in Thelon orogen north of the MacDonald fault have been quite well defined (Henderson et al., 1999), the timing of metamorphism in this region is only very loosely constrained between ca. 2.0 and 1.9 Ga (Henderson and van Breemen, 1992; Roddick and van Breemen, 1994).

Limited geochronological data indicate that this event extends to the northeast where it has been dated in high-grade gneiss at 1.91 Ga on northern Boothia Peninsula (Frisch and Hunt, 1993), 1.93 Ga within the Haughton impact structure on Devon Island (Shaerer and Deutsch, 1990), and 1.94 Ga on Ellesmere Island (Frisch and Hunt, 1988). The eastward extent of reworking during the Thelon orogeny is very poorly constrained. In the eastern Queen Maud block (south of Queen Maud Gulf), only one grain of ca. 1.9 Ga zircon was found among abundant ca. 2.6 – 2.43 Ga detrital zircon in a sample of the Sherman gneiss (Sheet 2; Schultz et al., 2007). It is not clear what grade of metamorphism produced this ca. 1.9 Ga monazite, nor how much of the area experienced, but may not record this event in the nonreactive granulite-facies rocks that were likely dehydrated during the Arrowsmith orogeny (see Sheet 1). On northern Boothia Peninsula, this event is assumed to extend over the region displayed on Sheet 2 based on similarities in structural trends (T. Frisch, personal communication, 2009). Preliminary data (R.G. Berman, W. J. Davis, J. J. Ryan, and L. Nadeau, unpub. data, 2008) suggest that structural and thermal reworking of the Rae craton extended south on Boothia Peninsula to where two amphibolite-facies samples record ca. 1.92 Ga monazite growth (Sheet 2).

South of the MacDonald fault, the evolution of the low-*P*, ultrahigh-*T* Taltson orogen is well documented. Following continental arc magmatism (east-dipping subduction beneath the western Rae craton) at 1.98 Ga (Thériault, 1992), synorogenic granitic plutons were intruded between 1.96 – 1.92 Ga (Bostock et al., 1991; McNicoll et al., 2000). In order to account for

metamorphic temperatures approaching 900 °C, Berman and Bostock (1997) suggested that an oceanic ridge may have been subducted. It should be noted, however, that De et al. (2000) advocate an intracontinental setting, rather than a continental margin setting, on the basis of geochemical characteristics.

In Saskatchewan, a regional northwest-striking deformational fabric is thought to record a tectonic accretionary event in the Taltson orogen (Ashton et al., 2009). North of the Lake Athabasca, metamorphism thought to be coeval with this northwest-striking fabric (Ashton et al., 2005b), is represented by: a) widespread ca. 1.93 Ga leucogranites derived during crustal melting (Hartlaub et al., 2005), b) the growth of 1.94-1.92 Ga metamorphic monazite in pelitic gneisses west of the Black Bay fault (Sheet 2; R.G. Berman, K. Ashton, W.J. Davis, unpub. data, 2005), and c) 1.93-1.92 Ga lower intercept ages from Archean granitoid rocks west of the Black Bay fault (Ashton et al., 2007). The crustal melt leucogranites occur as far east as the central Beaverlodge domain. The dominantly upper-amphibolite facies conditions assigned to this area are based on the partially melted nature and mineral assemblages of rocks defining the northwest-striking fabric (Ashton et al., 2009).

In distinguishing the effects of the Taltson from the Snowbird orogeny, it is important to note that the metamorphic ages within the core of the Taltson orogen are generally no younger than ca. 1.92 Ga (Ashton et al., 2009). In contrast, the Snowbird orogen is generally characterized by metamorphic ages younger than 1.91 Ga (see below). This distinction in ages strongly suggests that the early deformation (characterized by a shallow, northwest-striking foliation) and medium-*P*, granulite-facies metamorphism dated at 1.92 Ga immediately west of the Snowbird tectonic zone and east of Wholdaia Lake (Sheet 2; Martel et al., 2007) are causally related to the Taltson orogeny. It is on this basis that the high-grade, Taltson aged metamorphism

has been interpolated across northern Saskatchewan (Ashton et al., 2009) and the southern Northwest Territories.

The occurrence of ca. 2.3 Ga K-Ar biotite ages north of Tazin Lake, northwest Saskatchewan (Koster and Baadsgaard, 1970; Banks, 1980) suggests that metamorphic grade there during the Taltson event did not exceed subgreenschist facies. Towards the north, K-Ar biotite ages progressively young down to ca. 1890 Ma (Banks, 1980), suggesting that the metamorphic grade of the Taltson event increases up to at least greenschist facies, but probably not higher since ca. 2.5 – 2.4 Ga K-Ar hornblende ages are preserved (see Sheet 2). To the northeast of the K-Ar data in this region of the southwestern Northwest Territories, there is only one monazite age (R.G. Berman, H. Qui, W.J. Davis, S. Pehrsson, unpub. data, 2006) which is interpreted to indicate Taltson age amphibolite-facies metamorphism of a ca. 2.3 Ga (Arrowsmith orogeny) granulite-facies gneiss. Based on this sample, a large region of the southwestern Northwest Territories, extending up to the MacDonald fault in the southern Thelon tectonic zone, is considered to have experienced Taltson – Thelon age amphibolite-facies metamorphism (Sheet 2).

South of the Athabasca Basin, most of the exposed basement, with the exception of rocks near the Virgin River shear zone, was subjected to low- to moderate-*P*, granulite-facies conditions (Ashton et al., 2009). This thermal event was accompanied by partial melting in most rock types and the formation of a northwest-striking gneissosity (e.g. Scott, 1985). Well-developed northwest-striking magnetic striping in the sub-Phanerozoic portion of this region (Lloyd domain) is thought to be a regional manifestation of that gneissosity (Ashton et al., 2009). Metamorphic zircon of ca. 1.94-1.93 Ga age in ca. 1.98 Ga arc-type plutonic rocks (Stern et al., 2003) is considered coeval with development of the northwest-striking fabric and the granulite-facies assemblages.

The northwest-striking fabrics and presumed ca. 1.94-1.93 Ga assemblages continue northward beneath the unmetamorphosed Athabasca Group and are exposed in the basement core of the Carswell structure. Anatectic granitoid rocks sampled from the basement to the Athabasca Group just south of the Carswell structure also returned 1.93-1.91 Ga U-Pb zircon ion microprobe ages (Brouand et al., 2002). In addition, monazite in pelitic rocks from the basement core of the Carswell structure are ca 1.93 Ga in age (R.G. Berman, K. Ashton, W. J. Davis, unpub. data, 2005). These ages suggest that the metamorphic culmination during the Taltson orogeny extended southeastward from the Northwest Territories and northeastern Alberta to the Virgin River shear zone (Ashton et al., 2009).

Snowbird phase of Hudsonian Orogeny (ca. 1.91 – 1.87 Ga; bright colour series on Sheet 2)

The occurrence of a high-pressure, ca. 1.9 Ga metamorphic belt along much of the length of the Snowbird tectonic zone has been proposed to reflect the ca. 1.9 Ga collision between the Hearne craton and the composite Rae-Chesterfield block (Berman et al., 2007; see also Hoffman, 1988). This orogenic event is considered to represent the earliest phase of the Hudsonian orogeny, involving pre-1.865 Ga (Cumberland – Wathaman Batholiths) microcontinent collisions with the southeast flank Rae craton (Berman et al., 2007) following terminal collisions in the Taltson-Thelon orogen on the west side of the Rae craton. The geochronological and metamorphic data defining this belt form three domains. First, Barrovian-style (kyanite-staurolite-garnet), middle amphibolite-facies metapelitic rocks in the Josephine River belt crystallized during a tectonometamorphic event that is regionally well dated at 1.89 Ga (Sheet 2; Berman et al., 2002; 2007). The Paleoproterozoic age of these sediments, which contain detrital zircon as young as 2.3 Ga (Berman et al., 2007), suggests that they may have formed on the

margin of the composite Rae-Chesterfield block before its interpreted collision with the Hearne craton at ca. 1.91 Ga (Berman et al., 2007). This timing would provide ~20 Ma to thicken and incubate these sediments to the temperature (lower amphibolite-facies) required for ca. 1.89 Ga monazite growth. This domain extends northeast to the western part of exposed bedrock on Southampton Island, where ca. 1.88 Ga monazite inclusions form part of an early fabric within garnet porphyroblasts in samples from the western side of the exposed bedrock (Berman et al., 2011). The inferred eastern limit of the Snowbird orogen on Southampton Island (Sheet 2) lines up with the geophysically defined eastern edge of the Hearne craton, providing a plausible explanation for the apparent absence of ca. 1.88 Ga monazite to the east of this boundary on Southampton Island (Berman et al., 2010c). Further east on southwestern Baffin Island, a thermal event marked by ca. 1.87 Ga monazite growth may be related to the ca. 1.9 Ga accretion of an Archean (Sugluk, Corrigan et al., 2009) crustal block to Meta Incognita microcontinent (Wodicka et al., 2010) prior to the ca. 1.88-1.865 Ga collision of this composite block with the Rae craton (see below and Sheet 3).

The second high-*P* domain is interpolated through a large portion of the Chesterfield block on the basis of several samples near MacQuoid and Angikuni Lakes (Sheet 2). In this region garnet porphyroblasts show post-tectonic, ~10 kbar overgrowths at middle amphibolite to upper amphibolite facies (Berman et al., 2000, 2002b; 2007). The best constraints on timing suggest that this post-tectonic, high-*P* event occurred at 1.88 Ga (Berman et al., 2007). In the vicinity of Yathkyed Lake, a subgreenschist-facies region must have been at higher crustal levels to preserve K-Ar biotite (and hornblende) cooling ages > 2300 Ma (Wanless et al., 1979; Sandeman, 2001). The extent of this low-grade region is only very approximately constrained by available isotopic data (Sheet 2). Therefore the metamorphic patterns shown for ca. 1.9 Ga are

assumed to be the same at ca. 1.8 Ga (see Sheet 3), when there is more control from the distribution of greenschist-facies Paleoproterozoic Hurwitz Group metasedimentary rocks.

The third high-*P* domain, which defines the northern segment of the Snowbird tectonic zone, consists of 12-13 kbar, mafic gabbro-anorthosite that intruded at granulite facies into granitic lower crust at ca. 1.901 Ma (Kramanituur complex; Sanborn-Barrie et al., 2000) at 1.907 Ga (Uvauk complex, Mills et al., 2007), and at 1.905 Ga (Daly Bay complex; Berman et al., 2007). In this segment of the STZ, metamorphism began prior to gabbro-anorthosite emplacement, based on ca. 1.92 – 1.91 Ga monazite in granulite-facies metasediments at Kramanituur complex (Sanborn-Barrie et al., 2001) and 1.917 Ga granulite-facies tonalite at Daly Bay complex (Berman et al., 2007). The region of > 10 kbar lower crust, which contains smaller lenses of high-*P* amphibolite, extends north to the Chesterfield fault zone, a northwest vergent thrust interpreted as responsible for at least partial exhumation of this high-*P* domain (Berman et al., 2007). The timing of this exhumation is constrained by ^{40}Ar - ^{39}Ar hornblende ages between ca. 1.92 – 1.88 Ga (H. A. Sandeman, R. G. Berman, unpub. data, 2001). In the eastern part of this region (with the possible exception of western Southampton Island discussed above), there are no mineralogical or quantitative data to allow assessment of the pressure or timing of crystallization, which is assumed to be ca. 1.9 Ga on Sheet 2.

The third high-*P* domain described above in the northern STZ is similar to ca. 1.91 – 1.90 Ga, 10-12 kbar granulite-facies rocks that occur in the east-vergent, hanging wall of the Chipman shear zone (Martel et al., 2007; Legs Lake shear zone of Mahan et al., 2003), and comprise the central STZ in northwest Saskatchewan (East Athabasca mylonite zone; Williams et al., 1999; Baldwin et al., 2003; Williams and Hanmer, 2005; Flowers et al., 2006; Mahan et al., 2008) and southern Northwest Territories (Snowbird Lake map sheet; Martel et al., 2007). Recent detailed

studies suggest that most of the region has undergone two periods of high-*P* metamorphism and deformation at ca. 2.56 – 2.50 Ga (Baldwin et al., 2003, 2006; Mahan et al., 2006a; Flowers et al., 2008) and ca. 1.9 Ga (Baldwin et al., 2004; Flowers et al., 2006; Mahan et al., 2006a, 2006b) prior to ca. 1.85 – 1.80 Ga partial exhumation along the Chipman shear zone (Mahan et al., 2003; Martel et al., 2007) and ca. 1.8 Ga dextral offset along the Grease River shear zone (Mahan et al., 2005; Dumond et al., 2008). Whereas pressures of 10-12 kbar characterize much of this area in Saskatchewan, 1.904 Ga eclogite (Sheet 2) records conditions up to 1000 °C and 20 kbar (Baldwin et al., 2004). In the eastern part of this region, mafic dykes (Chipman swarm) dated at 1.896 ± 0.001 Ga (Flowers et al., 2006) were emplaced synkinematically with respect to a steep, northeast-striking fabric that regionally records sinistral strike slip displacement and overprints an earlier, shallow-dipping gneissosity (Williams et al., 1995; Mahan et al., 2008). Petrological studies suggest contrasting ca. 1.9 Ga P-T paths, with the eclogite recording prograde burial followed by isothermal decompression (Baldwin et al., 2003, 2004), whereas mafic granulites to the east are interpreted to record isobaric heating prior to cooling and decompression (Mahan et al., 2008). Also enigmatic is the interpretation that rocks in the central STZ of Saskatchewan experienced long-term residence in the deep crust between ca. 2.5 Ga and 1.9 Ga (e.g. Williams and Hanmer, 2005; Flowers et al., 2006; Mahan et al., 2008), whereas the Rae craton in the adjacent region of the Northwest Territories (the area east of Wholdaia Lake; Snowbird Lake map sheet) is characterized by ca. 2.56 Ga basement gneiss which had been exhumed prior to deposition of < 2.07 Ga (youngest detrital zircon) sediments that were buried prior to experiencing lower pressure (7.6 kbar) granulite-facies metamorphism and deformation (northwest-striking, shallow foliation and gneissosity) at 1.92 Ga (Martel et al., 2007).

Based on metamorphic zircon and monazite ages in northern Saskatchewan, the region affected by the 1.91-1.90 Ga “Snowbird” event extends west of the central Snowbird tectonic

zone, broadly coinciding with the structural overprint of regional northeast-trending folds on the earlier (ca. 1.92 Ga) northwest-striking fabric (Ashton et al., 2009). Because of the overlap in this area between the Taltson and Snowbird events, “overprint circles” display the granulite-facies metamorphism in this region (Sheet 2) that is characterized by somewhat lower pressures than those within the central Snowbird tectonic zone, suggesting that a tilted crustal section exposes medium-*P* granulites west of the Grease River shear zone (Ashton et al., 2009). East of the Black Bay fault (see inset on Sheet 2), the overprinting of these two events has produced an as yet unresolved range of 1.93-1.90 Ga chemical monazite ages from pelitic gneisses, even though metamorphic zircon and titanite yield well-constrained 1.91-1.90 Ga ages (Hartlaub, 2004). Because of this lack of geochronological resolution, it is unclear at what time (Taltson or Snowbird) the metamorphic zonation from lower amphibolite-facies to upper amphibolite-facies formed (Ashton et al., 2009). The Black Bay Fault may have been initiated at this time as a northwest-verging thrust fault (Ashton et al., 2009). West of the Black Bay Fault, metamorphic grade is inferred to decrease towards the greenschist facies that is recorded further west in the Waugh Lake group (see inset on Sheet 2).

South of the Athabasca Basin, northwest-striking fabrics (in the Lloyd domain) are overprinted by northeast-trending folds that tighten towards the Virgin River shear zone (Ashton et al., 2009). The earliest northeast-striking mylonitic fabric in the shear zone is thought to be the same age as that folding. Mafic dykes that cut the older gneissosity were deformed by the northeast-trending folds. The presence of locally derived leucosomes axial planar to the northeast-trending folds, hornblende replacing primary pyroxenes in the dykes, and hornblende in the oldest mylonitic rocks suggest that this deformation developed under upper amphibolite-facies conditions (Ashton et al., 2009). The main northeast-striking fabric is older than a suite of

ca. 1.83-1.82 Ga biotite granites (e.g. Stern et al., 2003) that stitch the Virgin River shear zone. Those granites contain only a weak northeast-striking foliation that is thought to result from reactivation of the shear zone (Ashton et al., 2009). Zircon overgrowths have been dated at ca. 1.90 Ga in ca. 1.98 Ga arc-type plutonic rocks (Stern et al., 2003) immediately west of the Virgin River shear zone. Given that the northeast-striking fabric has been established to have formed at 1.90 Ga in the STZ northeast of the Athabasca basin (Flowers et al., 2006), these ca. 1.90 Ga zircon overgrowths probably record coeval development of the northeast-striking structures in this region.

The Virgin River shear zone was subjected to repeated reactivation during its uplift history. The Virgin Schist Group is a lower to middle amphibolite-facies, commonly phyllonitic, supracrustal succession exposed in the core of the Virgin River shear zone. The provenance of these rocks remains unclear as does the age of superimposed metamorphism, although 1.90 Ga is considered a minimum age (Ashton et al., 2009).

Main phase of Hudsonian Orogeny (ca. 1.86 – 1.75 Ga; Sheet 3)

The Trans-Hudson Orogen extends across the western Churchill Province from northern Saskatchewan to Baffin Island. Following decades of focussed research in the Manitoba-Saskatchewan and North Quebec – Baffin Island segments, it is the best characterized of all orogenic events affecting this part of the Canadian shield.

Saskatchewan – Manitoba segment of the Trans-Hudson orogen

In the Saskatchewan – Manitoba segment of the orogen, metamorphism and deformation resulted from arc accretion followed by collision of the Hearne, Sask, and Superior crustal blocks. Metamorphism within the zone of accreted juvenile rocks mainly resulted from their interaction with the Sask craton in the south and Hearne craton in the northwest.

Along the Hearne craton margin, northwest-directed subduction under the continent may have commenced as early as 1.92 Ga (Tran et al., 2003; R. Maxeiner et al., in prep) and been accompanied by arc accretion. Alternatively, subduction with the opposite polarity may have been followed by ca. 1.88 – 1.865 Ga accretion of the La Ronge – Lynn Lake arcs to the southeastern Hearne craton margin (Bickford *et al.* 1990; 1994; Corrigan *et al.* 2005). By 1.86 Ga, continued westward transport of the oceanic plate resulted in development of a major continental arc (Wathaman Batholith) on the eastern margin of the Hearne craton (e.g. Meyer et al., 1992). Evidence for early thermotectonic processes (not shown on Sheets 2 or 3) comes from 1917 Ma volcanogenic rocks of the Porter Bay Complex in the Peter Lake domain (Maxeiner et al., in prep.), which were subjected to a low-grade thermotectonic overprint prior to emplacement of 1913 Ma Porter Bay Complex intrusions (Rayner et al., 2005a). As shown on Sheet 3, the lower sequence of the Campbell River Group and its basement in the Peter Lake domain are interpreted (Ashton et al., 2009) to have been metamorphosed to upper amphibolite facies during

northward subduction and emplacement of the 1.92-1.91 Porter Bay Complex, a proposed continental arc.

Earliest deformation in the Rottenstone domain, Saskatchewan is reportedly older than 1.87 Ga (Bickford et al., 1987; Maxeiner et al., 2004). Bickford et al. (1990) and Meyer et al. (1992) relate this early deformation to arc accretion (i.e. La Ronge arc) along the eastern margin of the Hearne craton via southeast-directed subduction at about 1.87 Ga. Alternatively, early deformation, at least in the La Ronge domain, Saskatchewan, may have been due to intra-oceanic amalgamation processes (Maxeiner et al., 2005) and interaction with the Flin Flon-Glennie Complex (Lucas et al., 1997). The latter was amalgamated with attendant deformation and possible metamorphism from juvenile ocean floor and arc assemblages at 1.88-1.87 Ga (Lucas et al., 1996). Evidence of these earlier metamorphic conditions is scarce due to subsequent overprinting by the main <1.83 Ga metamorphic event (Ashton et al., 2009).

The Flin Flon – Glennie Complex together with sedimentary rocks of the Kisseynew domain, were thrust over the Archean Sask craton (exposed in several tectonic windows) at about 1.83 Ga (Ashton et al., 2005a), resulting in widespread partial melting. Isograds are modified after Plint et al. (1996). Rare relict kyanite observed west of the Saskatchewan – Manitoba border (Ashton and Wheatley, 1986) suggests that these early metamorphic conditions locally involved higher pressures than those inferred from the final stable assemblages (Ashton et al., 2009). The Flin Flon block must have remained at relatively high structural levels during most of the collisional phase of the orogeny as it preserves greenschist- to subgreenschist- facies assemblages.

The resulting episode of renewed tectonic thickening produced widespread upper amphibolite-facies metamorphism in the Rottenstone domain (Ashton et al., 2009), recorded by the emplacement of abundant anatectic granitic sheets that range from ca. 1.84 to 1.81 Ga in age

(MacLachlan et al. 2004; Clark et al., 2005). Metamorphic monazite in pelitic migmatite records an age of ca. 1.82 Ga. In slightly lower-grade rocks of the former Crew Lake Belt of the eastern Rottenstone domain, which exhibit only incipient partial melting (Ashton et al., 2009), metamorphic monazite in psammite ranges from 1.83 to 1.80 Ga (N. Rayner, unpub. data, 2006). Metamorphic titanite from various rock types in the central Rottenstone domain ranges from ca. 1.81 to 1.79 Ga (MacLachlan et al. 2004) and likely records the timing of cooling through the ~600° C closure temperature (Heaman and Parrish, 1991). In the Mudjatik and Wollaston domains of the Hearne craton, this early pulse of high-grade Trans-Hudson metamorphism is recorded by upper amphibolite to granulite-facies mineral assemblages in Wollaston Supergroup rocks (Schledewitz, 1978; Harper et al., 2005; Card et al., 2006).

Ashton et al. (2009) interpret that renewed tectonic thickening attendant to terminal collision caused a thermal pulse responsible for widespread recrystallization of mylonite zones and variable overprinting of the early metamorphic assemblages via terrane accretion (e.g. Tran, 1997). Although the orientation of isograds has been modified by late folding in some areas (e.g. the contact between the northeastern Glennie domain and the Kisseynew domain), elsewhere (e.g. the area northwest of Flin Flon) they crosscut the main regional fabric at high angles (Ashton et al., 2009). Portions of the Flin Flon – Glennie and La Ronge arc complexes that remained at shallow crustal levels during this deformation have been preserved as granite-greenstone belts marked by metamorphic ‘lows’ (Ashton et al., 2009). Most of the Hearne craton, except for the Peter Lake domain, was also reworked at this time under upper amphibolite to granulite-facies conditions (e.g. Orrell et al., 1999; Tran, 2001; Chakungal et al., 2004). This main metamorphic event took place at moderate pressures (i.e. 5-7 GPa) and culminated at about

1.82-1.80 Ga (e.g. Annesley et al., 1992, 1997; Syme et al., 1998; Ashton et al., 1999; Harper et al., 2006; van Breemen et al., 2007).

Titanite ages throughout the orogen (e.g. Ashton et al., 2005a; van Breemen et al., 2007), as well as ^{40}Ar - ^{39}Ar hornblende and mica ages east of the Tabbernor fault (Schneider et al., 2007), document uplift and cooling to about 1.77 Ga, at which time felsic magmas (Nueltin and Jan Lake granite suites) were intruded over much of the region (e.g. Heaman et al., 2003; Bickford et al., 2005). Sporadic 1.76-1.75 Ga titanite and rutile metamorphic ages (Sheet 3), as well as ca. 1.74 – 1.72 Ga ^{40}Ar - ^{39}Ar mica ages west of the Tabbernor fault, may result from high heat flow associated with this pulse of granitic magmatism (Ashton et al., 2009).

North Quebec – Baffin Island segment of the Trans-Hudson orogen

A large part of the metamorphic map for Baffin Island is taken from that of Fraser et al. (1978), updated on the basis of more recent studies in the Ege Bay region (Bethune and Scammell, 2003), central Baffin Island (St-Onge et al., 2005; St-Onge, personal communication), southern Baffin Island (St-Onge et al., 1999; 2006), and southwest Baffin Island (Sanborn-Barrie et al., 2008; St-Onge, personal communication). The following description of the North Quebec – Baffin Island segment of the Trans-Hudson orogen follows that summarized by St-Onge et al. (2002; 2006; see also references within). This segment of the orogen preserves a record or several accretionary and magmatic events prior to collisional reworking of three main tectonic assemblages (from the southern, lowermost assemblage to the northern, uppermost assemblage): (1) lower plate, Archean rocks of the Superior craton with overlying Paleoproterozoic sedimentary and volcanic cover rocks, (2) upper plate, accreted Paleoproterozoic ophiolite (c. 2.0 Ga), fore-arc clastic rocks, and the ca. 1.86 – 1.82 Ga Narsajuaq magmatic arc, and (3) upper plate, accreted rocks representing (a) Meta Incognita microcontinent basement with a <1.93 Ga

north-facing clastic – carbonate shelf succession (Lake Harbour Group), (b) Archean Rae craton basement with 2.2 – 1.92 Ga south-facing clastic and carbonate cover rocks (Piling Group), and (c) 1.865 – 1.848 Ga charnockitic rocks of the Cumberland Batholith.

The earliest event (D_0) involved ca. 1.88 – 1.865 Ga accretion of Meta Incognita microcontinent to the southern Rae craton, which produced a north-vergent, thin-skinned thrust and fold belt (north of the Cumberland Batholith, see Piling Group, Fig. 1) with ca. 1.88 – 1.85 Ga (M_0 ; see Sheet 2 for older ages) cordierite-andalusite assemblages recording P-T conditions of 550 – 600 °C and 3-4 kbar (Allan and Pattison, 2003).

Accretion of Narsajuaq arc (or Sugluk terrane, Corrigan et al., 2009) to the southern margin of Meta Incognita microcontinent produced D_1 deformation, and together with intrusion of the Cumberland Batholith, M_{1a} regional lower granulite-facies (garnet-cordierite-K feldspar) metamorphism. This ca. 1849 – 1835 Ma metamorphic event records conditions of 790 – 845 °C and 7-8 kbar on southern Baffin Island (St-Onge et al., 2007), where subsequent thermal events are recorded at 1833 – 1829 (M_{1b}) and at 1836 – 1825 Ma (M_{1c} at lower structural levels). On southwest Baffin Island, monazite records the main tectonometamorphic event between ca. 1850 and 1835 Ma at metamorphic conditions that vary from upper amphibolite facies (~700 °C – 5 kbar) to lower amphibolite facies (Rayner et al., 2008). In a lower to middle-amphibolite facies supracrustal belt on southwestern Baffin Island, peak P-T conditions of 620 °C - 4 kbar appear to have been attained on an anti-clockwise P-T-t path (Smye et al., 2009) which contrasts with the clockwise P-T-t paths determined for samples further east (St-Onge et al., 2006; Smye et al., 2009). These differences are considered to reflect the pre-deformation emplacement of Narsajuaq arc plutons in the west and syn- to post-deformation intrusion of the Cumberland Batholith in the east (Smye et al., 2009).

Further north in the Ege Bay region, a low-*P* greenschist to amphibolite-facies region dated at ca. 1.85 – 1.82 Ga is juxtaposed against granulite-facies rocks to the north (Jackson and Berman, 2000; Bethune and Scammell, 2003). The south-dipping Isortoq fault zone that separates these areas (Sheet 3) is thought to represent a ca. 1.85 Ga northwest-vergent thrust that was reactivated as a normal fault (Bethune and Scammell, 2003). Further north and northeast, amphibolite- to granulite-facies gneisses (some with thermobarometric data suggesting high pressures, Sheet 3) are thought to have been exhumed during a later stage of southwest-vergent thrusting (Jackson and Berman, 2000).

Collision between the Superior craton and the composite upper plate produced ca. 1820 – 1805 Ma D₂ south-vergent, recumbent folds and thrusts in the upper plate, with M₂ retrograde upper amphibolite-facies metamorphism dated between ca. 1820 – 1808 Ma (St-Onge et al., 2006). In the lower plate, ca. 1830 – 1815 Ma piggy back sequence thrusting produced greenschist- to amphibolite-facies, Barrovian style metamorphism (up to 575 °C and 9.1 kbar) in the Cape Smith belt (St-Onge et al., 2000). Between ca. 1815 to 1785 Ma (Scott and St-Onge, 1995) the Superior craton underwent high-*P* (8 – 10 kbar; 585 – 725 °C; St-Onge and Ijewliw, 1996) hydration reactions at the basement-cover contact. This amphibolite-facies hydration zone has been assumed to extend across the width of the basal décollement at the base of the Cape Smith belt (Sheet 3). The occurrence of young (ca. 1660 – 1590 Ma) K-Ar hornblende cooling ages (Wanless et al., 1974) within two samples of the Superior craton north of the basal décollement (Sheet 3) has been taken to indicate that this portion of the Superior craton experienced at least greenschist-facies thermal conditions. Post-collisional, M₃ metamorphism at ca. 1795 – 1785 Ma involved reequilibration at amphibolite facies and emplacement of leucogranites in both the upper and lower plates.

Central Rae segment of the Trans-Hudson orogen

Metamorphic and geochronological studies in the central Rae craton west of Hudson Bay has demonstrated extensive reworking of this region between ca. 1.86 and 1.77 Ga (Carson et al., 2004; Berman et al., 2005, Berman et al., 2010a). The northwestern part of the Committee Bay belt provides a window into early tectonometamorphism, with metapelitic rocks recording monazite and zircon ages between 1.86 – 1.84 Ga (Carson et al., 2004; Berman et al., 2005; Berman et al., 2010a). Low-*P* (5 kbar maximum) metamorphism as a response to tectonic thickening (dominantly northwest-vergent D_2 folds; Sanborn-Barrie et al., 2011) is indicated by the the absence of plutonism prior to or during metamorphism (Sanborn-Barrie et al., 2011), clockwise P-T paths retrieved from lower to upper amphibolite-facies metapelitic rocks, and garnet and staurolite growth that is synchronous or late with respect to deformation. Given the ~10 to 30 Ma time lag for thermal incubation after onset of thickening (depending on the heat productivity of the rocks and peak temperatures relative to the temperature of monazite growth; Berman et al., 2010b), this metamorphism is considered to reflect ca. 1.87 Ga collision of a Meta Incognita microcontinent with the Rae craton (Berman et al., 2005, 2010a, 2010b), consistent with the 1.88 – 1.865 Ga constraints for the timing of this collision (St-Onge et al., 2002). The eastern end of the Committee Bay belt records D_2 deformation and metamorphism at ca. 1.82 – 1.81 Ga (Berman et al., 2010a; Sanborn-Barrie et al., 2011), which is thought to reflect its relative geographic proximity to the eastern promontory of the Superior craton during its ca. 1.82 Ga collision with the composite Rae - Meta Incognita crustal block (Berman et al., 2010a, 2010b). Further east on southern Melville Peninsula, data are available to evaluate the timing of near-peak metamorphism and deformaton, except for a ca. 1.80 Ga whole rock Rb-Sr age (Henderson, 1983).

On southern Boothia Peninsula north of the Committee Bay belt, the main Paleoproterozoic deformation and metamorphism is recorded by later monazite growth (ca. 1.83 Ga; Berman et al., 2008) than in the western Committee Bay belt, a difference that can be attributed to the deeper structural levels exposed in the Boothia region (~ 7 kbar; Berman et al., 2008) than in the Committee Bay belt (~5 kbar; Berman et al., 2005, 2010a). To the south in the region of Woodburn Lake, the main Paleoproterozoic deformation and lower to middle amphibolite-facies metamorphism is dated at 1.84 – 1.83 Ga (R. G. Berman, W.J. Davis, S. Pehrsson, unpub. data, 2005). On Southampton Island, metamorphic grade ranges from upper amphibolite facies to granulite-facies, with most *in situ* monazite dating (Berman et al., 2011) indicating ca. 1.86 – 1.82 Ga ages. However, some of the granulite-facies metamorphism is very likely older than ca. 1.9 Ga (not shown on Sheet 2), as dated charnockites have yielded disparate ages of ca. 3.0 Ga, 2.67 Ga, and 1.93 Ga (N. Rayner., unpub. data, 2009). Similarly, upper amphibolite to granulite-facies metamorphism of metaplutonic rocks on southern Melville Peninsula (sheet 3) is considered in part Archean and in part Paleoproterozoic in age. The diachroneity in the timing of deformation and metamorphism across the central Rae craton illustrates the domainal pattern of reworking which appears to be controlled by heterogeneity in transmission of tectonic stresses, in exposed crustal levels (e.g. deeper levels in the Boothia mainland region (Berman et al., 2008), in the heat productivity of the thickened rocks which controls the rate at, and grade to which rocks are heated and thermally weakened (e.g. 1.86 Ga monazite growth in the western CBb; Berman et al., 2010b), and in the relative timing of heat input from plutonic rocks (e.g. 1.86 Ga monazite growth on Southampton Island; Berman et al., 2011).

In contrast to the 1.86 – 1.81 Ga tectonometamorphism described above in the Committee Bay belt, the central part of this belt records monazite growth at 1.79 Ga that is associated with

dextral shearing at lower amphibolite facies in the south (Berman et al., 2010a) and with upper amphibolite facies in the north (Carson et al., 2004). Further northwest on southern Boothia Peninsula, ca. 1.79 Ga monazite ages are anomalously young relative to the regions further east (Berman et al., 2008). This younger episode of metamorphic crystallization is thought to reflect slightly deeper crustal levels (and temperatures above that required for monazite growth) within a north to northwest-trending synform produced during cross-folding in the final stages of amalgamation of Laurentia (Berman et al., 2008; 2010a, 2010b; Sanborn-Barrie et al., 2011).

The areal extent of reworking of Rae crust during the Hudsonian orogeny is still quite poorly known, with few constraints outside of the areas discussed above. Large regions with uncertain grade of 1.86 – 1.75 Ga metamorphism occur between Chesterfield Inlet and Wager Bay, and on Boothia and Melville Peninsulas as well as the adjacent area on northwestern Baffin Island. These regions are shown as either undivided amphibolite-facies or greenschist- to amphibolite-facies metamorphic grade, primarily based on the sparse occurrence of K-Ar ages younger than ca. 1750 Ma for hornblende or biotite, respectively (Wanless et al., 1966, 1979; Stevens et al., 1982a, b). The region of greenschist- to amphibolite facies on northwestern Melville Peninsula outlines an area that previously reached granulite facies (Schau et al., 1993), most likely at ca. 2.6 Ga. A K-Ar hornblende age of 2657 Ma (Stevens et al., 1982a) suggests that one area on the northern flank of the Penrhyn Group (just south of Committee Bay) did not exceed greenschist facies.

Northwest Baffin Island is characterized by upper amphibolite-facies regions, but scant geochronological data suggest these are part of the ca. 2.5 – 2.3 Ga Arrowsmith orogen (see Sheet 1), not part of the Trans-Hudson orogen as originally thought (e.g. Fraser et al., 1978). Preservation in this region of ca. 2.31 – 2.39 Ga Rb-Sr isochron ages suggest that Hudsonian

metamorphic grade did not exceed lower amphibolite facies. These interpretations suggest that metamorphic grade decreases northward from the ca. 1.82 Ga Dexterity granulite belt (Jackson and Berman, 2000) which has been juxtaposed against greenschist- to lower amphibolite-facies rocks along the Isortoq fault zone (Bethune and Scammell, 2003). On the northeast side of Baffin Island, upper amphibolite- to granulite-facies areas are considered part of the Northeast Baffin thrust belt (Jackson and Berman, 2000). Within this belt, reconnaissance level thermobarometric results (without geochronological control) indicate high-*P* granulite facies conditions (Jackson and Berman, 2000). Because upper amphibolite-facies surrounding regions do not have any pressure indicators, it is not known whether high-*P* conditions characterize the whole belt.

The north and northwestern limits of reworking related to the Hudsonian orogeny are also poorly constrained. Two geochronological studies (Schultz et al., 2007; Berman et al., 2005) have found no evidence for Hudsonian age zircon or monazite growth in the Queen Maud region, and K-Ar data are lacking there. On southern Boothia peninsula, Hudsonian reworking extends north to a ca. 2.6 Ga granulite belt (Berman et al., 2008; shown as an upper amphibolite-facies belt on Sheet 1;), which is interpreted to have been thrust southwards (Berman et al., 2008), leading to earlier cooling through hornblende closure (ca. 1.8 Ga, N. Joyce and J. Ryan, unpublished data) than the region to the south of this belt (ca. 1.76 Ga, N. Joyce and J. Ryan, unpublished data). Further north, there is an indication from monazite ages in one sample (Sheet 3) of cryptic Hudsonian reworking. On northern Boothia Peninsula, two K-Ar hornblende cooling ages are 1918 ± 12 Ma (Roddick et al., 1992) and 1831 ± 20 Ma (Frisch and Hunt, 1993), whereas K-Ar biotite ages are younger than ca. 1740 Ma (Wanless et al., 1979; Frisch and Hunt, 1993). These data suggest that Hudsonian thermal reworking was at least greenschist facies, and locally exceeded lower amphibolite facies (Sheet 3).

In northwestern Saskatchewan, the Hudsonian event has not been recognized west of the Virgin River shear zone (Orrell et al., 1999). North of Lake Athabasca, the effects of the Hudsonian orogeny diminish west of the Black Bay fault (Ashton et al., 2009). Along strike in the Northwest Territories, the few available data suggest that much of the Rae was at least thermally reworked at amphibolite facies at least as far west as the Howard Lake shear zone. Further west, the Nonacho sediments were metamorphosed to lower subgreenschist facies (Aspler, 1985).

REFERENCES

- Allan, M. M. and Pattison, D. R. M., 2003. Deformation history and metamorphism of a synformal depression, Longstaff Bluff Formation metaturbidite, central Baffin Island, Nunavut; Current Research - Geological Survey of Canada, 2003-C15, 11p.
- Annesley, I.R., Madore, C. and Krogh, T.E., 1992a. U-Pb zircon, titanite and monazite ages from the Wollaston Domain: a summary; Summary of Investigations 1992, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Miscellaneous Report 92-4, p.61-65.
- Annesley, I.R., Madore, C. and Krogh, T.E., 1992b. U-Pb geochronology of some granitoids from the Peter Lake Domain: a summary; Summary of Investigations 1992, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Miscellaneous Report 92-4, p.168-171.
- Annesley, I.R., Madore, C. and Krogh, T.E., 1993. U-Pb zircon, titanite, and monazite geochronology from the Wollaston-Mudjatik domain boundary, Karpinka Lake area, northern Saskatchewan; Geological Association of Canada - Mineralogical Association of Canada, v.18, p. 3.
- Annesley, I.R., Madore, C., Krogh, T.E., Kwok, Y.Y. and Kamo, S.L., 1999. New U-Pb zircon and monazite geochronological results for Archean and Paleoproterozoic basement to the southeastern part of the Athabasca Basin, Saskatchewan; Summary of Investigations 1999, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Paper 99-4.2, p. 90-99.
- Annesley, I.R., Madore, C., Shi, R. and Krogh, T.E., 1997. U-Pb geochronology of thermotectonic events in the Wollaston Lake area, Wollaston Domain: a summary of 1994-1996 results; Summary of investigations 1997, Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Miscellaneous Report 97-4, p. 162-173.
- Ansdell, K. M., 2005. Tectonic evolution of the Manitoba-Saskatchewan segment of the Paleoproterozoic Trans-Hudson Orogen, Canada; Canadian Journal of Earth Sciences, v. 42, p. 741-759.
- Ansdell, K. M., Lucas, S. B., Connors, K. and Stern, R. A., 1995. Kiseynew metasedimentary gneiss belt, Trans-Hudson Orogen (Canada); back-arc origin and collisional inversion; Geology, v. 23, p. 1039-1043.
- Ansdell, K.M. and Stern, R., 1997. The Scimitar Complex: Initial SHRIMP U-Pb and Nd isotope data from granitoid rocks; Summary of Investigations, Saskatchewan Geological Survey, Saskatchewan Energy and Mines Paper 97-4, p. 145-149.
- Ashton, K. E. and Wheatley, K. J., 1986. Preliminary report on the Kiseynew gniesses in the Kiseynew-Wildnest lakes area, Saskatchewan; Current Research of the Geological Survey of Canada, v. 86-1B, p. 305-317.
- Ashton, K., Berman, R. G., Card, C., Maxeiner, R., Harper, C. T. and MacLachlan, K., 2009. Metamorphic map of northern Saskatchewan; Geological Survey of Canada Open File, v. 5443 (1:1000000 scale).
- Ashton, K.E., Card, C., and Modeland, S., 2005b. Geological reconnaissance of the northern Tazin Lake Map Area (NTS 74N), including parts of the Ena, Nolan, Zemplak, and Taltson domains, Rae Province; *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2005-4.2, 24p.

- Ashton, K.E., Card, C.D., Davis, W., and Heaman, L.M., 2007. New U-Pb zircon age dates from the Tazin Lake map area (NTS 74N); *in* Summary of Investigations 2007, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2007-4.2, 8p.
- Ashton, K.E., Card, C.D., Heaman, L.M., and Coolican, J., 1999a. Rae Northeast: analytical update and new field results from the north-central Train Lake domain; *in* Summary of Investigations 1999, Saskatchewan Geological Survey, Saskatchewan Energy Mines, Miscellaneous Report 99-4.2, 3-16.
- Ashton, K.E., Heaman, J.F., Lewry, J.F., Hartlaub, R.P. and Shi, R., 1999b. Age and origin of the Jan Lake Complex: a glimpse at the buried Archean craton of the Trans-Hudson Orogen; *Canadian Journal of Earth Science*, v. 36, p. 185-208.
- Ashton, K.E., Hunt, P.A., and Froese, E., 1992. Age constraints on the evolution of the Flin Flon volcanic belt and Kiseeynew belt, Saskatchewan and Manitoba; *in* Radiogenic Age and Isotopic Studies: Report 5, Geological Survey of Canada, Paper 91-2, p. 55-69.
- Ashton, K.E., Lewry, J.F., Heaman, L.M., Hartlaub, R.P., Stauffer, M.R., and Tran, H.T., 2005a. The Pelican Thrust Zone: basal detachment between the Archean Sask Craton and Paleoproterozoic Flin Flon-Glennie Complex, western Trans-Hudson orogen; *Canadian Journal of Earth Sciences*, v. 42, p. 685-706.
- Aspler, L.B., 1985. Geology of the Nonasho Basin (Early Proterozoic) N.W.T.; Ph.D. Thesis, Carleton University, Ottawa, Ontario, 385p.
- Aspler, L.B., Chiarenzelli, J.R., and McNicoll, V.J., 2002. Paleoproterozoic basement-cover infolding and thick-skinned thrusting in Hearne Domain, Nunavut, Canada; intracratonic response to Trans-Hudson Orogen; *Precambrian Research*, v. 116, p. 331-354.
- Baadsgaard, H. and Godfrey, J.D., 1967. Geochronology of the Canadian Shield in northeastern Alberta 1. Andrew Lake area; *Canadian Journal of Earth Sciences*, v. 4, p. 541-563.
- Baldwin, J. A., Bowring, S. A., Williams, M. L. and Mahan, K., 2006. Eclogites of the Snowbird tectonic zone; petrological and U-Pb geochronological evidence for Paleoproterozoic high-pressure metamorphism in the western Canadian Shield; *Lithos*, v. 88, p. 173-200.
- Baldwin, J.A., Bowring, S.A., and Williams, M.L., 2003. Petrological and geochronological constraints of high pressure, high temperature metamorphism in the Snowbird tectonic zone, Canada; *Journal of Metamorphic Geology*, v. 21, p. 81-98.
- Baldwin, J.A., Bowring, S.A., Williams, M.L., and Williams, I.S., 2004. Eclogites of the Snowbird tectonic zone; petrological and U-Pb geochronological evidence for Paleoproterozoic high-pressure metamorphism in the western Canadian Shield; *Contributions to Mineralogy and Petrology*, v. 147, p. 528-548.
- Banks, C.S., 1980. Geochronology, general geology and structure of Hill Island Lake-Tazin Lake areas; M. Sc. Thesis, University of Alberta, Edmonton, AB, Canada.
- Beaumont-Smith, C.J. and Bohm, C.O., 2002. Structural analysis and geochronological studies in the Lynn Lake greenstone belt and its gold-bearing shear zones (NTS 64C10, 11, 12, 14, 15, 16); Report of Activities 2002, Manitoba Industry, Trade and Mines, Manitoba Geological Survey, p. 159-170.

- Bell, K., 1985. Geochronology of the Carswell area, northern Saskatchewan; *in* Laine R., Alonso, D. and Svab, M. (eds) *The Carswell Structure Uranium Deposits, Saskatchewan*. Geological Association of Canada Paper 29, p. 33-46.
- Berman, R. G. and Bostock, H. H., 1997. Metamorphism in the Northern Taltson Magmatic zone, Northwest Territories; *Canadian Mineralogist*, v. 35, p. 1069-1092.
- Berman, R. G., 1990. Mixing properties of Ca-Mg-Fe-Mn garnets; *American Mineralogist*, v. 75, p. 328-344.
- Berman, R. G., Camacho, A. and Sandeman, H., 2010b. Diachronous Paleoproterozoic deformation and metamorphism in the Committee Bay belt, Rae Province, Nunavut: insights from 40Ar-39Ar cooling ages and thermal modelling; *Journal of Metamorphic Geology*, v. 28, 439–457.
- Berman, R. G., Easton, R. M. and Nadeau, L., 2000a. A new tectonometamorphic map of the Canadian Shield, Introduction; *in*: *Tectonometamorphic studies in the Canadian Shield (Part II)*; *The Canadian Mineralogist*, v. 38, p. 277-285.
- Berman, R.G., Davis, W.J., Aspler, L., Chiarenzelli., 2002b. SHRIMP U/Pb ages of multiple metamorphic events in the Angikuni Lake area, western Churchill Province, Nunavut; *Current Research Geological Survey of Canada*, 2002-F3, p. 1-9.
- Berman, R.G., Davis, W.J., Pehrsson, S., 2007. The collisional Snowbird tectonic zone resurrected: growth of Laurentia during the 1.9 Ga accretionary phase of the Trans-Hudson orogeny; *Geology*, v. 35, p. 911-914.
- Berman, R.G., Davis, W.J., Ryan, J.J., Tella, S., 2002a. In-situ U/Pb SHRIMP geochronology of Barrovian facies-series metasedimentary rocks in the Happy lake and Josephine River supracrustal belts: implications for the Paleoproterozoic architecture of the northern Hearne domain, Nunavut; *Current Research Geological Survey of Canada*, 2002-F4, p. 1-14.
- Berman, R.G., Rayner, N., Sanborn-Barrie, M., Chakungal, J., 2011. New constraints on the tectonothermal history of Southampton Island, Nunavut provided by in situ SHRIMP geochronology and thermobarometry; *Current Research Geological Survey of Canada* (in press).
- Berman, R. G., Ryan, J. J., Tella, S., Sanborn-Barrie, M., Stern, R., Aspler, L., Hanmer, S. and Davis, W., 2000b. The case of multiple metamorphic events in the Western Churchill Province: evidence from linked thermobarometric and in-situ SHRIMP data, and jury deliberations.; *GeoCanada 2000 Conference extended abstracts*, Calgary, Alberta, abstract 836.
- Berman, R.G., Ryan, J.J., Davis, W.J., Nadeau, L., 2008. Preliminary results of linked in-situ SHRIMP dating and thermobarometry of the Boothia mainland area, north-central Rae Province, Nunavut; *Current Research Geological Survey of Canada*, 2008-2, p. 1-13.
- Berman, R. G., Sanborn-Barrie, M., Rayner, N., Carson, C., Sandeman, H. and Skulski, T., 2010a. Petrological and in situ SHRIMP geochronological constraints on tectonometamorphic evolution of the Committee Bay belt, Nunavut; *Precambrian Research*, v. 181, 1-20.
- Berman, R.G., Sanborn-Barrie, M., Stern, R., Carson, C., 2005. Tectonometamorphism at ca. 2.35 and 1.85 Ga in the Rae domain, western Churchill Province, Nunavut, Canada: insights from structural, metamorphic and in situ geochronological analysis of the southwestern Committee Bay belt; *Canadian Mineralogist*, v. 43, p. 409-442.
- Berman, R.G., Sapers, H., Davis, W.J., Rayner, N., Pehrsson, S., Sandeman, H., Sanborn-Barrie, M. and Carr, S., 2010c. The collisional Snowbird tectonic zone revisited: new constraints on the

Chesterfield Inlet segment provided by metamorphic monazite, detrital zircon, and ^{40}Ar - ^{39}Ar geochronology. GeoCanada 2010 Conference extended abstracts, Calgary, Alberta, abstract #670.

- Bethune, K.M. and Scammell, R.J., 1997. Precambrian geology, Koch Island area, District of Franklin, Northwest Territories (part of NTS 37C); Geological Survey of Canada Open File 3391, scale 1:50 000.
- Bethune, K.M. and Scammell, R.J., 2003. Distinguishing between Archean and Paleoproterozoic tectonism, and the evolution of the Isortoq fault zone, Ege Bay area, north-central Baffin Island, Canada; *Canadian Journal of Earth Sciences*, v.40, p. 1111-1135.
- Bickford, M. E., Collerson, K. D. and Lewry, J. F., 1994. Crustal history of the Rae and Hearne provinces, southwestern Canadian Shield, Saskatchewan; constraints from geochronologic and isotopic data; *Precambrian Research*, v. 68, p. 1-21.
- Bickford, M. E., Collerson, K. D., Lewry, J. F., Van, S. W. R. and Chiarenzelli, J. R., 1990. Proterozoic collisional tectonism in the Trans-Hudson Orogen, Saskatchewan; *Geology*, v. 18, p. 14-18.
- Bickford, M. E., Hajnal, Z., Mock, T. D., Steinhart, W. E., Collerson, K. D. and Lewry, J. F., 2005. Origin of the Archean Sask Craton and its extent within the Trans-Hudson Orogen; evidence from Pb and Nd isotopic compositions of basement rocks and post-orogenic intrusions; *Canadian Journal of Earth Sciences*, v. 42, p. 659-684.
- Bickford, M.E., Mock, T.D., Steinhart III, W.E., Collerson, K.D., and Lewry, J.F., 2005. Origin of the Sask Craton and its extent within the Trans-Hudson orogen: evidence from Pb and Nd isotopic compositions of basement rocks and post-orogenic intrusions; *Canadian Journal of Earth Sciences*, v. 42, p. 659-684.
- Bickford, M.E., van Schmus, W.R., Collerson, K.D. and Macdonald, R., 1987. U-Pb zircon geochronology project: new results and interpretation; *in* Summary of Investigations, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 87-4, p. 76-79.
- Bostock, H. H. and van Breemen, O., 1992. Ages of detrital and metamorphic zircons and monazites from a pre-Taltson magmatic zone basin at the western margin of Rae Province; *Canadian Journal of Earth Science*, v. 31, p. 1353-1364.
- Bostock, H. H., van Breemen, O. and Loveridge, W. D., 1991. Further geochronology of plutonic rocks in northern Taltson Magmatic Zone, District of Mackenzie, N.W.T.; *in* Radiometric age and isotopic studies, report 4; Geological Survey of Canada, Current Research, v. 90-2, p. 67-78.
- Bostock, H.H. and Loveridge, W.D., 1988. Geochronology of the Talston Magmatic zone and its eastern cratonic margin, District of Mackenzie; *Radiogenic Age and Isotopic Studies: Report 2*, Geological Survey of Canada, Paper 88-2, p. 59-65.
- Brouand, M., Cuney, M.L. and Deloule, E., 2002. Age and nature of the plutonism of the western part of the Athabasca Basin basement (northern Saskatchewan, Canada); GAC-MAC joint annual meeting, v. 27.
- Card, C., Harper, C.T., Barsi, N., Lesperance, J., and Smith, J., 2006. Investigation of the Wollaston-Mudjatik Transition, Charcoal Lake and Cochrane River (parts of NTS 64L-9, -10, -11, -14, -15,

and -16); *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2006-4.2, 16p.

- Carpenter, R., Duke, N., Sandeman, H., and Stern, R., 2005. Relative and absolute timing of gold mineralization along the Meliadine Trend, Nunavut, Canada; evidence for Paleoproterozoic gold hosted in an Archean greenstone belt; *Economic Geology*, v. 100, p. 567–576.
- Carson, C.J., Berman, R.G., Stern, R.A., Sanborn-Barrie, M., Skulski, T., and Sandeman, H.A., 2004. Age constraints on the Paleoproterozoic tectonometamorphic history of the Committee Bay region, western Churchill Province, Canada: evidence from zircon and in situ monazite SHRIMP geochronology; *Canadian Journal of Earth Sciences*, v. 41, p. 1049-1076.
- Cavell, P.A., Wijbrans, J.R. and Baadsgaard, H., 1992. Archean magmatism in the Kaminak Lake area, District of Keewatin, Northwest Territories: ages of the carbonatite-bearing alkaline complex and some host granitoid rocks; *Canadian Journal of Earth Science*, v. 29, p. 896-908.
- Chakungal, J., Reynolds, P., Jamieson, R. and Corrigan, D., 2004. Slow post-orogenic cooling in a deeply eroded orogen: Reindeer Zone, Trans-Hudson Orogen, Saskatchewan; *Canadian Journal of Earth Sciences*, v. 41, p. 867–880.
- Chiarenzelli, J., Aspler, L., Villeneuve, M. and Lewry, J., 1998. Early Proterozoic evolution of the Saskatchewan Craton and its allochthonous cover, Trans-Hudson Orogen; *Journal of Geology*, v. 106, p. 247-267.
- Clark, G.S., and Schledewitz, D.C.P., 1988. Rubidium - strontium ages of Archean and Proterozoic rocks in the Nejanilini and Great Island domains, Churchill Province, northern Manitoba, Canada; *Canadian Journal of Earth Sciences*, v. 25, p. 246-254.
- Clarke, D. B., Hajnal, Z., Henry, A. S. and Hamilton, M. A., 2005. Composition, age, and origin of granitoid rocks in the Davin Lake area, Rottenstone Domain, Trans-Hudson Orogen, northern Saskatchewan; *Canadian Journal of Earth Sciences*, v. 42, p. 599-633.
- Clarke, D.B., Henry, A.S., and Hamilton, M.A., 2005. Composition, age, and origin of granitoid rocks in the Davin Lake area, Rottenstone Domain, Trans-Hudson Orogen, northern Saskatchewan; *Canadian Journal of Earth Sciences*, v. 42, p. 599-633.
- Corrigan, D., Hajnal, Z., Nemeth, B. and Lucas, S. B., 2005. Tectonic framework of a Paleoproterozoic arc-continent to continent-continent collisional zone, Trans-Hudson Orogen, from geological and seismic reflection studies; *Canadian Journal of Earth Sciences*, v. 42, p. 421-434.
- Corrigan, D., Maxeiner, R.O., and Harper, C.T., 2001. Preliminary U-Pb results from the La Ronge-Lynn Lake Bridge project; Summary of Investigations 2001, Saskatchewan Geological Survey, Saskatchewan Energy and Mines Miscellaneous Report 2001-4.2, p. 111-115.
- Corrigan, D., Pehrsson, S., Wodicka, N. and de Kemp, E., 2009. The Palaeoproterozoic Trans-Hudson Orogen: a prototype of modern accretionary processes; *in* B. J. Murphy, Keppie, J.D. and Hynes, A.J. (eds), *Ancient Orogens and Modern Analogues*, Geological Society, London, p. 459–481.
- Cranstone, D.A., and Turek, A., 1976. Geological and geochronological relationships of the Thompson nickel belt, Manitoba; *Canadian Journal of Earth Sciences*, v. 13, p.1058-1069.

- David, J. Bailes, A.H. and Machado, N., 1996. Evolution of the Snow Lake portion of the Paleoproterozoic Flin Flon and Kiseynew belts, Trans-Hudson Orogen, Manitoba, Canada; *Precambrian Research*, v. 80, p. 107-124.
- Davis, W. J., Hanmer, S., Tella, S., Sandeman, H. A. and Ryan, J. J., 2006. U/Pb geochronology of the MacQuoid supracrustal belt and Cross Bay plutonic complex; key components of the northwestern Hearne Subdomain, western Churchill Province, Nunavut, Canada; *Precambrian Research*, v. 145, p. 53-80.
- Davis, W.J., Hanmer, S., and Sandeman, H.A., 2004. Temporal evolution of the Neoproterozoic central Hearne supracrustal belt; rapid generation of juvenile crust in a suprasubduction zone setting; *Precambrian Research*, v. 134, p. 85-112.
- Davis, W.J., Hanmer, S., Tella, S., Sandeman, H.A., and Ryan, J.J., 2006. U/Pb geochronology of the MacQuoid supracrustal belt and Cross Bay plutonic complex; key components of the northwestern Hearne Subdomain, western Churchill Province, Nunavut, Canada; *Precambrian Research*, v. 145, p. 53-80.
- De, S. K., Chacko, T., Creaser, R. A. and Muehlenbachs, K., 2000. Geochemical and Nd-Pb-O isotope systematics of granites from the Taltson magmatic zone, NE Alberta; implications for early Proterozoic tectonics in western Laurentia; *Precambrian Research*, v. 102, p. 221-249.
- Dumond, G. and Noah McLean, M. L. W., Michael J. Jercinovic, Samuel A. Bowring, 2008. High-resolution dating of granite petrogenesis and deformation in a lower crustal shear zone: Athabasca granulite terrane, western Canadian Shield; *Chemical Geology*, v. 254, p. 175-196.
- Easton, R. M. and Berman, R. G., 2004. Tectonometamorphic map of Ontario, Canada and parts of the United States of America; Geological Survey of Canada Open File, v. 1810, 1:1,500,000 (1 sheet).
- Fedorowich, J.S., Kerrich, R. and Stauffer, M.R., 1995. Geodynamic evolution and thermal history of the central Flin Flon Domain, Trans-Hudson Orogen: constraints from structural development, $^{40}\text{Ar}/^{39}\text{Ar}$, and stable isotope geothermometry; *Tectonics*, v. 14, p. 472-503.
- Flowers, R.M., Bowring, S.A., and Williams, M.L., 2006a. Timescales and significance of high-pressure, high-temperature metamorphism and mafic dike anatexis, Snowbird tectonic zone, Canada; *Contributions to Mineralogy and Petrology*, v. 151, p. 558-581.
- Flowers, R.M., Bowring, S.A., Mahan, K., Williams, M.L. and Williams, I., 2008. Stabilization and reactivation of cratonic lithosphere from the lower crustal record in the western Canadian shield; *Contributions to Mineralogy and Petrology*, v. 156, p. 529-549.
- Flowers, R. M., Mahan, K. H., Bowring, S. A., Williams, M. L., Pringle, M. S. and Hodges, K. V., 2006b. Multistage exhumation and juxtaposition of lower continental crust in the western Canadian Shield; linking high-resolution U/Pb and $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronometry with pressure-temperature-deformation paths; *Tectonics*, v. 25, TC4003, doi:10.1029/2005TC001912.
- Fraser, J. A., Heywood, W. W. and Mazurski, M. A., 1978. Metamorphic map of the Canadian Shield; Geological Survey of Canada, "A" Series Map, v. 1475A (1:3,500,000 scale).
- Frisch, T. and Hunt, P.A., 1988. U-Pb zircon and monazite ages from the Precambrian Shield of Ellesmere and Devon islands, Arctic Archipelago; Radiogenic and Isotopic Studies Report 2, Geological Survey of Canada, Paper 88-2, p. 117-126.

- Frisch, T. and Hunt, P.A., 1993. Reconnaissance U-Pb geochronology of the crystalline core of the Boothia Uplift, District of Franklin, Northwest Territories; Radiogenic Age and Isotopic Studies Report 7; Geological Survey of Canada, Paper 93-2, p. 3-22.
- Gordon, T.M., Hunt, P.A., Bailes, A.H., and Syme, E.C., 1990. U-Pb zircon ages from the Flin Flon and Kiseynew belts, Manitoba: Chronology of crust formation at an Early Proterozoic accretionary margin; *in* Lewry, J.F., and Stauffer, M.R. (eds.) The Early Proterozoic Trans-Hudson Orogen of North America. Geological Association of Canada, Special Paper 37, p. 177-199.
- Harper, C.T., Ebel, C., Yeo, G., Card, C., and Nelson, C., 2005. Wollaston Lake Project: Geology of the Wollaston Supergroup in the Rabbabou Bay – Wellbelove Bay area, northeast Wollaston Lake, Saskatchewan (parts of NTS 64L-06, -07, -10 and -11); *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2005-4.2, 25p.
- Harper, C.T., Rayner, N., and Card, C., 2006. Preliminary U-Pb ages for Archean basement and Proterozoic intrusion from the northern Wollaston Lake area, Saskatchewan; *in* Summary of Investigations 2006, Volume 2, Saskatchewan Geological Survey, Saskatchewan Industry Resources Misc. Rep. 2006-4.2, 9p.
- Harrison, T. M., 1981. Diffusion of ^{40}Ar in hornblende; *Contributions to Mineralogy and Petrology*, v. 78, p. 324-331.
- Harrison, T. M., Duncan, I. & McDougall, I., 1985. Diffusion of ^{40}Ar in biotite; temperature, pressure and compositional effects; *Geochimica et Cosmochimica Acta*, v. 49, p. 2461-2468.
- Hartlaub, R. P., Ashton, K. E., Heaman, L. M. and Chacko, T., 2007. Circa 2.3-Ga Magmatism of the Arrowsmith Orogeny, Uranium City region, Western Churchill Craton, Canada; *Journal of Geology*, v. 115, p. 181-195.
- Hartlaub, R.P., 2004. Archean and Proterozoic evolution of the Beaverlodge Belt, Churchill craton, Canada; Unpubl. Ph.D. thesis, University of Alberta, Edmonton, 189p.
- Hartlaub, R.P., Chacko, T., Heaman, L.M., Creaser, R.A., Ashton, K.E., and Simonetti, T., 2005. Ancient (Meso- to Paleoproterozoic) crust in the Rae Province, Canada: Evidence from Sm-Nd and U-Pb constraints; *Precambrian Research*, v. 141, p. 137-153.
- Hartlaub, R.P., Chacko, T., Heaman, L.M., Creaser, R.A., Ashton, K.E., and Simonetti, T., 2005. Ancient (Meso- to Paleoproterozoic) crust in the Rae Province, Canada: evidence from Sm-Nd and U-Pb constraints; *Precambrian Research*, v. 141, p. 137-153.
- Heaman, L. and Ludden, J.N. Short Course Handbook on Applications of Radiogenic Isotope Systems to Problems in Geology, v. 19, p. 59-102.
- Heaman, L. and Parrish, R., 1991. U-Pb geochronology of accessory minerals; *in*
- Heaman, L.M., Ashton, K.E., Hartlaub, R.P., and Harper, C.T., 1999. Preliminary U-Pb age constraints on the timing of the Clearwater Anorthosite Complex, metamorphism in the Train Lake Domain, and age of the Ourom Lake Group; *in* Summary of Investigations 1999, Volume 1, Saskatchewan Geological Survey, Saskatchewan Energy Mines, Misc. Rep. 99-4.1, p. 137-139.
- Heaman, L.M., Ashton, K.E., Reilly, B.A., Sibbald, T.I.I., Slimmon, W.L., and Thomas, D.J., 1993. 1992-93 U-Pb geochronological investigations in the Trans-Hudson Orogen, Saskatchewan; *in*

- Saskatchewan Geological Survey, Saskatchewan Energy and Mines, Misc. Report 93-4, p. 109-111.
- Heaman, L.M., Kamo, S.L., Delaney, G.D., Harper, C.T., Reilly, B.A., Slimmon, W.L. and Thomas, D.J., 1991. U-Pb geochronological investigation in the Trans-Hudson Orogen, Saskatchewan: preliminary results by the ROM Laboratory in 1990-91; Summary of Investigations 1991, Saskatchewan Geological Survey; Saskatchewan Energy and Mines, Miscellaneous Report 91-4, p. 74-75.
- Heaman, L.M., Maxeiner, R.O. and Slimmon, W.L., 1994. 1993-94 U-Pb geochronological investigations in the Trans-Hudson Orogen, Saskatchewan; Summary of Investigations 1994, Saskatchewan Geological Survey, Saskatchewan Energy and Mines Miscellaneous Report 94-4, p. 96-99.
- Henderson, J., James, D. and Thompson, P., 1999. Geology, Healey Lake - Artillery Lake, Northwest Territories- Nunavut; Geological Survey of Canada Open File, v. 3819, p. (scale 1:250,000).
- Henderson, J.B., and van Breemen, O., 1992. U-Pb zircon ages from an Archean orthogneiss and a Proterozoic metasedimentary gneiss of the Thelon tectonic zone, District of Mackenzie, Northwest Territories; Geological Survey of Canada Paper 91-02, p. 25-33.
- Henderson, J. R., 1983. Structure and metamorphism of the Aphebian Penrhyn Group and its Archean basement complex in the Lyon Inlet area, Melville Peninsula, District of Franklin; Geological Survey of Canada Bulletin, v. 324, p. 50p.
- Hinchey, A.M., Davis, W.J., Ryan, J.J., Nadeau, L., and James, D.T., 2006: Timing of magmatism, granulite-facies metamorphism and deformation in the Archean metaplutonic rocks of the Boothia mainland area within the Rea domain, Northwest Churchill Province, Nunavut; Geological Association of Canada – Mineralogical Association of Canada Program with Abstracts, v. 31, p. 67.
- Hoffman, P. F., 1988. United plates of America, the birth of a craton: Early Proterozoic assembly and growth of Laurentia; Annual Reviews of Earth and Planetary Science, v. 16, p. 543– 603.
- Hunt, P.A. and Roddick, C., 1987. A compilation of K-Ar ages. Radiogenic Age and Isotopic Studies Report 17, Geological Survey of Canada Paper 87-2, p. 143-204.
- Hunt, P.A., and Roddick, J.C., 1988. A compilation of K-Ar ages, Report 18; Radiogenic age and isotopic studies Report 2, Geological Survey of Canada Paper 88-2, p. 127-153.
- Hunt, P.A., and Zwanzig, H.V., 1993. U-Pb ages of intrusion, metamorphism and deformation in the Batty Lake area, Kisseynew Gneiss Belt, Manitoba; Manitoba Energy and Mines, Minerals Division, Report of Activities 1993, p. 34-37.
- Jackson, G. D., 1978. Basement gneisses and the Mary River Group, No. 4 deposit area, northwest Baffin Island, District of Franklin; *in* R.K. Wanless and W.D. Loveridge (eds) Rubidium-strontium Isotopic Age Studies Report 2 (Canadian Shield), Geological Survey of Canada Paper 77-14, p. 18-24.
- Jackson, G. D. and Berman, R. G., 2000. Precambrian Metamorphic and Tectonic Evolution of Northern Baffin Island, Nunavut, Canada; Canadian Mineralogist, v. 38, p. 399-422.

- Jackson, G.D., Hunt, P.A., Loveridge, W.D., and Parrish, R.R., 1990. Reconnaissance geochronology of Baffin Island, N.W.T.; Geological Survey of Canada Paper 89-02, p. 123-148.
- Koster, F. and Baadsgaard, H., 1970. On the geology and geochronology of northwestern Saskatchewan. I. Tazin Lake region; Canadian Journal of Earth Sciences, v. 7, p. 919-930.
- LeCheminant, A.N and Roddick, J.C., 1991. U–Pb zircon evidence for widespread 2.6 Ga felsic magmatism in the central District of Keewatin, N.W.T; Radiogenic age and isotopic studies: report 4. Geological Survey of Canada Paper 90-2, p. 91–99.
- Leech, G.B., Lowdon, J.A., Stockwell, C.H., and Wanless, R.K., 1963. Age determinations and geological studies - Report 4; Geological Survey of Canada, Paper 63-17, 140p.
- Lewry, J. F. and Collerson, K. D., 1990. The Trans-Hudson Orogen: extent, subdivisions and problems; *in* Lewry, J. F. & Stauffer, M. R. (eds) The Early Proterozoic Trans-Hudson Orogen of North America. Geological Association of Canada, Special Paper, v. 37, p. 1-14.
- Loveridge, W. D., Eade, K. E. and Sullivan, R. W., 1988. Geochronological Studies of Precambrian rocks from the southern District of Keewatin; Geological Survey of Canada Paper, v. 88-18, p. 36.
- Loveridge, W. D., Eade, K. E. and Sullivan, R. W., 1988. Geochronological Studies of Precambrian rocks from the southern District of Keewatin; Geological Survey of Canada Paper, v. 88-18, p. 36.
- Lowdon, J. A.; Stockwell, C. H.; Tipper, H. W.; Wanless, R. K., 1963. Age Determinations and Geological Studies; Geological Survey of Canada, Paper 62-17.
- Lucas, S.B., Stern, R.A., Syme, E.C, Reilly, B.A. and Thomas, D.J., 1996. Intraoceanic tectonics and the development of continental crust: 1.92 - 1.84 Ga evolution of the Flin Flon Belt; Geological Society of America Bulletin, v. 108, p. 602-629.
- Lucas, S.B., Stern, R.A., Syme, E.C., Zwanzig, H., Bailes, A.H., Ashton, K.E., Maxeiner, R.O., Ansdell, K.M., Lewry, J.F., Ryan, J.J. and Kraus, J., 1997. Tectonics of the southeastern Reindeer Zone, Trans-Hudson orogen (Manitoba and Saskatchewan); Geological Association of Canada - Mineralogical Association of Canada Program with Abstracts, v. 22, p. 93.
- Machado, N., Krogh, T.E., and Weber, W., 1990. U-Pb geochronology of basement gneisses in the Thompson belt (Manitoba): evidence for pre-Kenoran and Pikwitonei-type crust and Early Proterozoic basement reactivation in the western margin of the Archean Superior Province; Canadian Journal of Earth Sciences, v. 27, p. 794-802.
- Macdonald, R., 1999. Geological Map of Saskatchewan: Precambrian geology. Saskatchewan Geological Survey (1:1,000,000 scale).
- MacLachlan, K., Davis, W. J. and Relf, C., 2005a. Geochronological constraints on Neoproterozoic tectonism; multiple extensional events in the northwestern Hearne Domain, western Churchill Province, Canada; Canadian Journal of Earth Sciences, v. 42, p. 85-109.
- MacLachlan, K., Davis, W. J. and Relf, C., 2005b. Paleoproterozoic reworking of an Archean thrust fault in the Hearne domain, Western Churchill Province: U-Pb geochronological constraints; Canadian Journal of Earth Sciences, v. 42, p. 1313-1330.

- MacLachlan, K., Rayner, N., Dunning, G. and C. Leugner, 2004. New Results and ideas from the Rottenstone Domain Project; *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2004-4.2, 21p.
- MacLachlan, K., Rayner, N., Dunning, G., and Leugner, C., 2004. New results and ideas from the Rottenstone Domain Project; *in* Summary of Investigations 2004, Volume 2, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Misc. Rep. 2004-4.2, 21p.
- Mahan, K. and ML, W., 2005. Reconstruction of a large deep-crustal terrane: Implications for the Snowbird tectonic zone and early growth of Laurentia; *Geology*, v. 33, p. 385-388.
- Mahan, K. H., Goncalves, P., Flowers, R., Williams, M. L. and Hoffman-Setka, D., 2008. The role of heterogeneous strain in the development and preservation of a polymetamorphic record in high-P granulites, western Canadian Shield; *Journal of Metamorphic Geology*, v. 26, p. 669–694.
- Mahan, K. H., Williams, M. L. and Baldwin, J. A., 2003. Contractional uplift of deep crustal rocks along the Legs Lake shear zone, western Churchill Province, Canadian Shield; *Canadian Journal of Earth Sciences*, v. 40, p. 1085–1110.
- Mahan, K.H., Goncalves, P., Williams, M.L., and Jercinovic, M.J., 2006a. Dating metamorphic reactions and fluid flow; application to exhumation of high-P granulites in a crustal-scale shear zone, western Canadian Shield; *Journal of Metamorphic Geology*, v. 24, p. 193-217.
- Mahan, K.H., Williams, M.L., Flowers, R.M., Jercinovic, M.J., Baldwin, J.A., and Bowring, S.A., 2006b. Geochronological constraints on the Legs Lake shear zone with implications for regional exhumation of lower continental crust, western Churchill Province, Canadian Shield; *Contributions to Mineralogy and Petrology*, v. 152, p. 223-242.
- Manitoba Geological Survey, 1979. Geologic Map of Manitoba, Manitoba Department of Mines, Map 79-2 (1:1,000,000 scale).
- Martel, E., van Breemen, O., Berman, R. G. and Pehrsson, S., 2007. Geochronology and tectonometamorphic history of the Snowbird Lake area, NWT: a key window into the significance of the Snowbird tectonic zone; *Precambrian Research*, v. 161, p. 201-230.
- Maxeiner, R. O., Corrigan, D., Harper, C. T., MacDougall, D. G. and Ansdell, K., 2005. Paleoproterozoic arc and ophiolitic rocks on the northwest-margin of the Trans-Hudson Orogen, Saskatchewan, Canada; their contribution to a revised tectonic framework for the orogen; *Precambrian Research*, v. 136, p. 67-106.
- Maxeiner, R.O., Harper, C.T., Corrigan, D., and MacDougall, D.G., 2004. La Ronge–Lynn Lake Bridge Project: Geology of the southern Reindeer Lake area; *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Open File Report 2003-1 (3 maps).
- McDonough, M.R., McNicoll, V.J., Schetselaar, E.M., and Grover, T.W., 2000. Geochronological and kinematic constraints on the crustal shortening and escape in a two-sided oblique-slip collisional and magmatic orogen, Paleoproterozoic Taltson magmatic zone, northeastern Alberta; *Canadian Journal of Earth Science*, v. 37, v. 1549-1573.
- McNicoll, V.J., Theriault, R.J., and McDonough, M.R., 2000. Taltson basement gneisses: U-Pb and Nd isotopic constraints on the basement to the Paleoproterozoic Taltson magmatic zone, northeastern Alberta; *Canadian Journal of Earth Sciences*, v. 37, p. 1575-1596.

- Meyer, M. T., Bickford, M. E. and Lewry, J. F., 1992. The Wathaman Batholith; an early Proterozoic continental arc in the Trans-Hudson orogenic belt, Canada; *Geological Society of America Bulletin*, v. 104, p. 1073-1085.
- Mills, A., Berman, R.G., Davis, W.J., Tella, S., Carr, S.D., Roddick, C., and Hanmer, S., 2007. Thermobarometry and geochronology of the Uvauk Complex: a polymetamorphic Neoproterozoic and Paleoproterozoic segment of the Snowbird tectonic zone, Nunavut, Canada; *Canadian Journal of Earth Sciences*, v. 44, p. 245-266.
- Orrell, S.E., Bickford, M.E., and Lewry, J.F., 1999. Crustal evolution and age of thermotectonic reworking in the western hinterland of the Trans-Hudson Orogen, northern Saskatchewan; *Precambrian Research*, v. 95, p. 187-223.
- Parent, M., Machado, N. and Zwanzig, H. V., 1999. Comportement du système U-Pb dans la monazite et chronologie de la déformation et du métamorphisme des métasédiments du domaine de Kiseynew, orogène trans-hudsonien (Manitoba, Canada); *Canadian Journal of Earth Sciences*, v. 36, p. 1843-1857.
- Parrish, R.R., 1989. U-Pb geochronology of the Cape Smith Belt and Sugluk block, northern Quebec; *Geoscience Canada*, v. 16, p. 126-130.
- Patterson, J.G. and LeCheminant, A.N. 1985. A preliminary geological compilation map of the northeastern Barren Grounds, parts of the Districts of Keewatin and Franklin. Geological Survey of Canada Open File 1138 (scale 1:1,000,000).
- Paul, D; Hanmer, S; Tella, S; Peterson, T D; LeCheminant, A N, 2002. Compilation, bedrock geology of part of the western Churchill Province, Nunavut - Northwest Territories. Geological Survey of Canada, Open File 4236 (scale 1:1,000,000).
- Pehrsson, S., Currie, M., Ashton, K., Berman, R.G., Martel, E., Card, C., and Harper, C., 2011. Bedrock geology compilation of the southern Rae Domain, Western Churchill Province. Geological Survey of Canada, Open File 5744 (scale 1:550,000).
- Plint, H.E., Gordon, T.M., Froese, E. and Berman, R.G., 1996. Metamorphic data from the Flin Flon belt and adjacent Kiseynew belt. Geological Survey of Canada Open File 3160.
- Rayner, N.M., Maxeiner, R.O., and Corrigan, D., 2005a. Progress report on U-Pb geochronology results for the Peter Lake Domain Project; *in* Summary of Investigations 2005, Saskatchewan Geological Survey, Saskatchewan Industry Resources, Miscellaneous Report 2005-4.2, 12p.
- Rayner, N.M., Stern, R.A., and Bickford, M.E., 2005b. Tectonic implications of new SHRIMP and TIMS U-Pb geochronology of rocks from the Sask Craton, Peter Lake Domain, and Hearne margin, Trans-Hudson Orogen, Saskatchewan; *Canadian Journal of Earth Sciences*, v. 42, p. 635-657.
- Rayner, N., St-Onge, M., Berman, R., Sanborn-Barrie, M. and Wodicka, N., 2008. Polyphase tectonometamorphic history in the upper plate of Trans-Hudson orogen (southern Baffin Is.); Goldschmidt Conference, Vancouver, 2008, A780.
- Reilly, B.A., 1993. Bedrock geological mapping, Hatle Lake area (part of NTS 64M-13 and -14); Saskatchewan Energy and Mines Open File Report, 93-2, 21p.
- Roddick, J. C., Quigg, F. B. and Hunt, P. A., 1992. Miscellaneous 40 Ar - 39 Ar Ages and Analytical Procedures; Geological Survey of Canada, Paper, v. 92-02, p. 171-177.

- Roddick, J.C. and van Breemen, O., 1994. U-Pb zircon dating; a comparison of ion microprobe and single grain conventional analyses; Current Research - Geological Survey of Canada, Report: 1994-F, p. 1-9.
- Ryan, J. J., Davis, W.J., Berman, R. G., Sandeman, H.A., Hanmer, S. and Tella, S., 2000. 2.5 Ga granulite-facies activity and post-1.9 low-grade reactivation along the Big lake shear zone, MacQuoid-Gibson lakes area (Nunavut): a fundamental boundary in the Western Churchill Province; Geological Association of Canada - Mineralogical Association of Canada Abstracts with Programs, abstract #924.
- Sanborn-Barrie, M., Carr, S.D., and Theriault, R., 2001. Geochronological constraints on metamorphism, magmatism and exhumation of deep-crustal rocks of the Kramanituar Complex, with implications for the Paleoproterozoic evolution of the Archean western Churchill Province, Canada; Contributions to Mineralogy and Petrology, v. 141, p. 592-612.
- Sanborn-Barrie, M., Davis, W. J., Berman, R. G., Sandeman, H. A. I. and Skulski, T., 2011. Structural Geology of the Committee Bay area: a record of penetrative Paleoproterozoic deformation of an Archean hinterland; Precambrian Research (in prep).
- Sanborn-Barrie, M; St-Onge, M R; Young, M D; James, D T, 2008. Bedrock geology of southwestern Baffin Island, Nunavut: expanding the tectonostratigraphic framework with relevance to mineral resources; Current Research Geological Survey of Canada, v. 2008-6, 16p.
- Sandeman, H.A., 2001. ⁴⁰Ar-³⁹Ar geochronological investigations in the central Hearne domain, western Churchill Province, Nunavut: a progress report; Current Research Geological Survey of Canada, v. 2001-F4, 23p.
- Schau, M., 1993. Geology, northern Melville Peninsula, District of Franklin, Northwest Territories. Geological Survey of Canada Open File 2594 (scale 1:500,000)
- Schau, M., Venance, K., Berman, R. and Brodaric, B., 1993. Metamorphic map, northern Melville Peninsula, District of Franklin, Northwest Territories; Geological Survey of Canada Open File, v. 2594 (scale 1:500,000).
- Schledewitz, D. C. P., 1978. Patterns of regional metamorphism in the Churchill Province of Manitoba (north of 58 degrees); *in* J. A. Fraser & W. W. Heywood (eds) Metamorphism in the Canadian Shield, Geological Survey of Canada, Ottawa, p. 179-189.
- Schneider, D. A., Heizler, M. T., Bickford, M. E., Wortman, G. L., Condie, K. C. and Perilli, S., 2007. Timing constraints of orogeny to cratonization; thermochronology of the Paleoproterozoic Trans-Hudson Orogen, Manitoba and Saskatchewan, Canada; Precambrian Research, v. 153, p. 65-95.
- Schultz, M., Chacko, T., Heaman, L.M., Sandeman, H.A., Simonetti, A., and Creaser, R.A., 2007. The Queen Maud Block: A newly recognized Paleoproterozoic (2.4-2.5 Ga) terrane in northwest Laurentia; Geology, v. 35, p. 707-710.
- Scott, B.P., 1985. Geology of the upper Clearwater River area; Saskatchewan Energy and Mines Open File Report 85-2, 26p.
- Scott, D.J., 1997. Geology, U-Pb, and Pb-Pb geochronology of the Lake Harbour area, southern Baffin Island: implications for the Paleoproterozoic tectonic evolution of northeastern Laurentia; Canadian Journal of Earth Sciences, v. 34, p. 140-155.

- Scott, D.J., 1999. U-Pb geochronology of the eastern Hall Peninsula, southern Baffin Island, Canada: a northern link between the Archean of West Greenland and the Paleoproterozoic Torngat Orogen of northern Labrador; *Precambrian Research*, v. 93, p.5-26.
- Scott, D.J. and Gauthier, G., 1996. Comparison of TIMS (U-Pb. and laser ablation microprobe ICP-MS (Pb) techniques for age determination of detrital zircons from Paleoproterozoic metasedimentary rocks from northeastern Laurentia, Canada, with tectonic implications; *Chemical Geology*, v.131, p. 127-142.
- Scott, D. J. and St-Onge, M. R., 1995. Constraints on Pb closure temperature in titanite based on rocks from the Ungava Orogen, Canada; implications for U-Pb geochronology and P-T-t path determinations; *Geology*, v. 23, p. 1123-1126.
- Scott, D.J., Stern, R.A., St-Onge, M.R. and McMullen, S.M., 2002. U-Pb geochronology of detrital zircons in metasedimentary rocks from southern Baffin Island: implications for the Paleoproterozoic tectonic evolution of Northeastern Laurentia; *Canadian Journal of Earth Sciences*, v. 39, p. 611-623.
- Scott, D.J. and Wodicka, N., 1998. A second report on the U-Pb geochronology of southern Baffin Island, Northwest Territories; in *Radiogenic Age and Isotopic Studies: Report 11*, Geological Survey of Canada, Current Research 1998-F, p. 47-57.
- Shaerer, U. and Deutsch, A., 1990. Isotope systematics and shock-wave metamorphism. II: U-Pb and Rb-Sr in naturally shocked rocks, the Haughton impact structure, Canada; *Geochimica Cosmochimica Acta*, v. 54, p. 3435-3447.
- Skulski, T., Sanborn-Barrie, M. & Sandeman, H., 2003a. *Geology, Walker Lake and Arrowsmith River area, Nunavut*; Geological Survey of Canada Open File 3777 (scale 1:100,000)
- Smye, A. J., St-Onge, M. R. and Waters, D. J., 2009. Contrasting metamorphic pressure-temperature histories within the Trans-Hudson Orogen's hinterland, southwest Baffin Island, Nunavut; *Current Research of the Geological Survey of Canada*, v. 2009-6, p. 21p.
- Spear, F.S., 1993. *Metamorphic phase equilibria and pressure-temperature-time paths*. Mineralogical Society of America, Washington, D.C. 799 p.
- Stern, R.A., and Berman, R.G., 2000. Monazite U-Pb and Th-Pb geochronology by ion microprobe, with an application to in situ dating of an Archean metasedimentary rock; *Chemical Geology*, v. 172, p. 113-130.
- Stern, R.A., Card C.D., Pana, D. and Rayner, N., 2003. SHRIMP U-Pb ages of granitoid basement rocks of the southwestern part of the Athabasca Basin, Saskatchewan and Alberta; in *Radiogenic age and Isotopic Studies, Report 16*, Geological Survey of Canada, Current Research 2003-F, 20p.
- Stevens, R. D., DeLabio, R. N. and Lachance, G. R., 1982a. *Age Determinations and Geological Studies K-Ar Isotopic Ages, Report 15*; Geological Survey of Canada Paper, v. 81-02, 32p.
- Stevens, R. D., Lachance, G. R. and DeLabio, R. N., 1982b. *Age Determinations and Geological Studies K-Ar Isotopic Ages, Report 16*; Geological Survey of Canada, Paper, v. 82-02, p. 56p.
- St-Onge, M. R., Scott, D. J. and Wodicka, N., 1999. *Geological maps and descriptive notes and legend, parts of Meta Incognita Peninsula, Nunavut*; Geological Survey of Canada, "A" Series Map, v. 1979A-1985A (scale 1:100000), p.

- St-Onge, M. R., Scott, D. J. and Wodicka, N., 2002. Review of crustal architecture and evolution in the Ungava Peninsula, Baffin Island area; connection to the Lithoprobe ECSOOT transect; *Canadian Journal of Earth Sciences*, v. 39, p. 589-610.
- St-Onge, M. R., Scott, D. J., Corrigan, D. and Wodicka, N., 2005. Geological maps and descriptive notes and legend, central Baffin Island area, Nunavut; Geological Survey of Canada, "A" Series Map, v. 2077A-2082A (scale 1:100000), p.
- St-Onge, M. R., Searle, M. P. and Wodicka, N., 2006. Trans-Hudson Orogen of North America and Himalaya-Karakoram-Tibetan Orogen of Asia; structural and thermal characteristics of the lower and upper plates; *Tectonics*, v. 25, TC4006, doi:10.1029/2005TC001907, p. 22p.
- St-Onge, M.R., Scott, D.J., Corrigan, D., 2002a. Geology, Dewar Lakes, Nunavut; Geological Survey of Canada Open File 4201.
- St-Onge, M.R., Scott, D.J., Corrigan, D., 2002b. Geology, Straits Bay, Nunavut; Geological Survey of Canada Open File 4200.
- St-Onge, M.R., Scott, D.J., Corrigan, D., 2003a. Geology, Clyde River, Baffin Island, Nunavut; Geological Survey of Canada Open File 4432.
- St-Onge, M.R., Scott, D.J., Corrigan, D., 2003b. Geology, Blanchfield Lake, Baffin Island, Nunavut; Geological Survey of Canada Open File 4433.
- St-Onge, M.R., Scott, D.J., Corrigan, D., 2003c. Geology, Ikpik Bay, Baffin Island, Nunavut; Geological Survey of Canada Open File 4474.
- St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005a. Geology, Ikpik Bay, Baffin Island, Nunavut; Geological Survey of Canada A Series Map 2077A (scale 1,100000).
- St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005b. Geology, Clyde River, Baffin Island, Nunavut; Geological Survey of Canada, A Series Map 2079A (scale 1,100000).
- St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005c. Geology, Straits Bay, Baffin Island, Nunavut; Geological Survey of Canada, A Series Map 2081A (scale 1,100000).
- St-Onge, M.R., Scott, D.J., Corrigan, D., and Wodicka, N., 2005d. Geology, Dewar Lakes, Baffin Island, Nunavut; Geological Survey of Canada, A Series Map 2082A (scale 1,100000).
- St-Onge, M.R., Wodicka, N. and Ijewliw, O., 2007. Polymetamorphic evolution of Trans-Hudson Orogen on Baffin Island, Canada: constraints from Petrology, Structure, and U-Pb dating; *Journal of Petrology*, v. 48, p. 271-302.
- St-Onge, M. R., Wodicka, N. and Lucas, S. B., 2000. Granulite- and amphibolite-facies metamorphism in a convergent-plate margin setting; synthesis of the Quebec-Baffin segment of the Trans-Hudson Orogen; *Canadian Mineralogist*, v. 38, p. 379-398.
- Syme, E.C., Lucas, S.B., Zwanzig, H.V., Bailes, A.H., Ashton, K.E., and Haidl, F.M., 1998. Geology, NATMAP Shield Margin Project area of Flin Flon Belt, Manitoba/Saskatchewan accompanying notes to Geological Survey of Canada Map 1968A; Manitoba Energy and Mines Map A-98-2, Saskatchewan Energy and Mines Map 258A.
- ter Meer, M., 2001. Tectonometamorphic history of the Nowyak Complex, Nunavut, Canada; M. Sc. Thesis, Carleton University, Ottawa, 285p.

- Tella, S., 1993. Geology, Chesterfield Inlet, District of Keewatin, Northwest Territories; Geological Survey of Canada Open File, v. 2756, scale 1:250 000.
- Tella, S., Paul, D., Berman, R.G., Davis, W.J., Peterson, T.D., and Pehrsson, S.J., 2007, Bedrock geology compilation and regional synthesis of the Hearne domain north of 60 °N, western Churchill Province, Nunavut, Canada. Geological Survey of Canada Open File 5441 (scale 1:550,000).
- Thériault, R. J., 1992. Nd isotopic evolution of the Taltson magmatic zone, Northwest Territories, Canada; insights into early Proterozoic accretion along the western margin of the Churchill Province; *Journal of Geology*, v. 100, p. 465-475.
- Thomas, D.J. and Heaman, L.M. 1994. Geologic setting of the Jolu Gold Mine, Saskatchewan: U-Pb constraints on plutonism, deformation, mineralization, and metamorphism; *Economic Geology*, v. 89, p.1017-1029.
- Tran, H. T., Ansdell, K., Bethune, K., Watters, B. and Ashton, K., 2003. Nd isotope and geochemical constraints on the depositional setting of Paleoproterozoic metasedimentary rocks along the margin of the Archean Hearne Craton, Saskatchewan, Canada; *Precambrian Research*, v. 123, p. 1-28.
- Tran, H.T., 1997. Structural relations and thermotectonic history of the Medicine Rapids-Grassy Narrows area, northern Saskatchewan, Canada; M.Sc. thesis, University of Regina, Saskatchewan, 225p.
- Tran, H.T., 2001. Tectonic evolution of the Paleoproterozoic Wollaston Group in the Cree Lake Zone, northern Saskatchewan, Canada; Ph.D. thesis, University of Regina, Regina, Saskatchewan, 458p.
- van Breemen, O. and Bostock, H., 1994. Age of emplacement of Thoa metagabbro, western margin of Rae Province, Northwest Territories: initiation of rifting prior to Talston magmatism? *Radiogenic Age and Isotope Studies, Report 8, Geological Survey of Canada Current Research 1994-F*, p. 61-68.
- van Breemen, O., Harper, C.T., Berman, R.G., and Wodicka, N., 2007. Crustal evolution and Neoproterozoic assembly of the Central and Southern Hearne domain: evidence from U-Pb Geochronology and Sm-Nd isotopes of the Phelps Lake area, northeastern Saskatchewan; *Precambrian Research*, v. 159, p. 33-59.
- van Schmus, W. R., Persons, S. S., Macdonald, R., Sibbald, T. I. I., Patterson, D. F. and Davie, R. F., 1986. Preliminary results from U-Pb zircon geochronology of the Uranium City region, northwest Saskatchewan; *Summary of Investigations by the Saskatchewan Geological Survey*, v. 86-4, p. 108-111.
- Wanless, R.K., Stevens, R.D., Lachance, G.R., and Rimsaite, J.Y.H., 1966. K-Ar isotopic determinations, Report 6; Geological Survey of Canada, Paper 65-17, 101p.
- Wanless, R.K., Stevens, R.D., Lachance, G.R., and Delabio, R.N., 1970. Age Determinations and Geological Studies, K-Ar Isotopic Ages, Report 9, Geological Survey of Canada, Paper 69-2A, 78p.
- Wanless, R.K., Stevens, R.D., Lachance, G.R., and Delabio, R.N., 1972. Report 10, Geological Survey of Canada, Paper 71-2, 96p.

- Wanless, R. K., Stevens, R. D., Lachance, G. R. and Delabio, R. N. D., 1974. Age determinations and geological studies K-Ar isotopic ages, Report 12; Paper - Geological Survey of Canada, 74-2, 63p.
- Wanless, R.K., Stevens, R.D., Lachance, G.R. and Delabio, R.N., 1979. Age determinations and geological studies, K-Ar isotopic ages, Report 14; Geological Survey of Canada, Paper 79-2, p. 1-67.
- Wheeler, J.O., Hoffman, P.F., Card, K.D., Davidson, A., Sanford, B.V., Okulitch, A.V., Roest, W.R., 1996. Geological map of Canada. Geological Survey of Canada, "A" Series Map 1860A (scale 1:5,000,000).
- Williams, M. L. and Hanmer, S., 2005. Structural and metamorphic processes in the lower crust: evidence from the East Athabasca mylonite triangle, Canada, a deep-crustal isobarically cooled terrane; *in* M. Brown and T. Rushmer (eds), *Evolution and Differentiation of the Continental Crust*, Cambridge University Press, Cambridge, p. 232–268.
- Williams, M. L., Brown, M., Hanmer, S., Kopf, C. and Darrach, M., 1995. Syntectonic generation and segregation of tonalitic melts from amphibolite dikes in the lower crust, Striding-Athabasca mylonite zone, northern Saskatchewan; *Journal of Geophysical Research*, v. 100, p. 15717–15734.
- Williams, M. L., Jercinovic, M. J. and Terry, M. P., 1999. Age mapping and dating of monazite on the electron microprobe; deconvoluting multistage tectonic histories; *Geology*, v. 27, p. 1023-1026.
- Williams, M.L. and Jercinovic, M.J., 2002. Microprobe monazite geochronology; putting absolute time into microstructural analysis; *Journal of Structural Geology*, v. 24, p. 1013-1028.
- Wodicka, N. and Scott, D.J., 1997. A preliminary report on the U-Pb geochronology of the Meta Incognita Peninsula, southern Baffin Island, Northwest Territories; in Geological Survey of Canada Current Research 1997-C, p. 167-178.
- Wodicka, N., Whalen, J., St-Onge, M. R. and Corrigan, D., 2010. Meta Incognita microcontinent revisited: insights from U-Pb geochronology and Nd isotopes; GeoCanada 2010 Conference extended abstracts, Calgary, Alberta, abstract #982.