




# GEOCHEMICAL LANDSCAPES: ASPECTS OF VERTICAL AND HORIZONTAL

VA

**Eric C. Grunsky  
and Rick J. McNeil**

**Natural Resources  
Canada- Geological  
Survey of Canada**





# Geochemical Landscape

Earth Sciences Sector

- Geochemical landscape is a geochemical characterization of part of the earth's surface (Polynov, 1960\*), based on the joint influences of climate, relief, geology and vegetation on the chemical processes over a region
- The main application of geochemical landscapes is to support studies in:
  - Agriculture
  - Mineral exploration
  - Human health

\*Quoted in [Nature](#), v. 188,  
Issue 4744, 1960, p. 23



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# Spatial Aspects of Geochemical Landscapes

Earth Sciences Sector

- The scale of the assessment determines the required sample density.
- Search theory sets a working sample density at  $\frac{1}{2}$  the size of the target that is being sought.
- Statistically based sample designs permit defensible results (i.e. analysis of variance).
- The number of sites is also dependent on the number of elements being analyzed if the elements are to be studied simultaneously.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# **Geochemical Landscapes**


## **National Scale**





# Geology of Canada

Earth Sciences Sector



Source: GSC  
Map 1860A  
(1: 5 000 000)

Download map and legend at [http://gsc.nrcan.gc.ca/map/1860a/index\\_e.php](http://gsc.nrcan.gc.ca/map/1860a/index_e.php)



Natural Resources  
Canada

Ressources naturelles  
Canada


Canada





# Surficial Geology of Canada

Earth Sciences Sector



Source: GSC  
Map 1880A  
(1: 5 000 000)

Download map and legend at [http://gsc.nrcan.gc.ca/map/1880a/index\\_e.php](http://gsc.nrcan.gc.ca/map/1880a/index_e.php)



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# Soils of Canada

Earth Sciences Sector



Source: AAFC Map  
(1: 6 500 000)



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada






## Hg (Mercury)

in Stream and Lake Sediment  
and Prairie Soil



Note: Contours based  
on GSC and  
Geoscience BC data



	<u>Median</u>	<u>Number of sites</u>
Lake Sediment	60 ppb	75,359
Stream Sediment	40 ppb	86,838
Prairie Soil	40 ppb	1076

Unpublished,  
based on GSC data





Note: Contours based on GSC, NL provincial and Geoscience BC data

### LOI (Loss-on-Ignition) in Stream and Lake Sediment



	<u>Median</u>	<u>Number of sites</u>
Lake Sediment	33 %	98,960
Stream Sediment	6.9 %	65,468

Unpublished,  
based on GSC data


GSC, 2009

Canada

Canada

Canada





Canada

Canada








# pH in Stream and Lake Water



Note: Contours based  
on GSC and  
Geoscience BC data



	<u>Median</u>	<u>Number of sites</u>
Lake Sediment	6.5	62,204
Stream Sediment	7.4	95,374

GSC, 2009

Unpublished,  
based on GSC data

Canada

Canada

Canada





# Regionality

Earth Sciences Sector

- In a country as large (  $9.2 \times 10^6 \text{ km}^2$ ), geologically diverse (the Precambrian Shield, the Cordilleras and the sedimentary basins) and physically variable (from the Arctic to carolinian and rain forests to grasslands), a single background range has little true meaning or relevance.
- In recognition of this, Environment Canada and Health Canada are moving to regionalize their risk assessments on the basis of ecological units, Ecoprovinces and Ecoregions.



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# EcoProvinces of Canada

Earth Sciences Sector



Source: CanSIS system: [http://geogratis.gc.ca/Ecosystem/1\\_ecosys/ecogifs/ecoprov.gif](http://geogratis.gc.ca/Ecosystem/1_ecosys/ecogifs/ecoprov.gif)



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# EcoRegions of Canada

Earth Sciences Sector



Source:  
CanSIS  
database



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# Example - Map-based Interface

Earth Sciences Sector

Survey count: 595

Refine results using...

Province: All

Year: = All

Organisation: All

Title:

Survey-type: All

Abstract:

Location: Longitude  
Latitude  
or  
NTS:

Submit Search  
Reset

1: 39492305

250K 50K Redraw

- Geoscience Data Repository interface showing location of catalogued geochemical surveys
- On-line at [http://apps1.gdr.nrcan.gc.ca/geochem/main\\_e.phtml](http://apps1.gdr.nrcan.gc.ca/geochem/main_e.phtml) [accessed March 26, 2010]



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# Sites by Ecozone

Earth Sciences Sector



Example

EcoZones

Lambert Conformal Conic  
Lon: 87°57'19" W  
Lat: 63°18'09" N  
Printed at: 2010-02-02



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# **Geochemical Landscapes**

## **Provincial Scale**








# Maritime Geology

Earth Sciences Sector



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# Maritime Soil Types

Earth Sciences Sector



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# Maritime Soil Survey Sites and EcoRegions

Earth Sciences Sector



Source for Ecoregions: Ecological Stratification Working Group, 1996.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# Sampling Strategies





# Sample Design for Soil Surveys

Earth Sciences Sector

- Sample strategy used for the North American Soil Geochemical Landscapes Project in Canada.
  - Generalized Random Tessellation Sample design (GRTS).
- GRTS is a random design based on the following: A *spatial point pattern is a collection of  $n$  points within a region  $D \subset R^2$  (a region  $D$  defined in a 2-dimensional real number space). The number of points is thought of as random and the points are considered to be generated by a stationary isotropic point process in  $R^2$ . This means that there is not preferred origin or orientation in the pattern. (Venables and Ripley, 2004).*
- Initial GRTS design – 1 site / 40km<sup>2</sup>
- Maritime GRTS design – 1 site / 20km<sup>2</sup>



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# Sample Designs

Earth Sciences Sector

- Analysis of variance (AOV) is important in establishing the variance at different levels of design (site spacing) and the repeatability of patterns across an area.
  - Stratified random sample design
  - Balanced nested hierarchical sample design
  - Unbalanced nested hierarchical sample design



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# GRTS Design – Continental Scale

Earth Sciences Sector



Preselected sites  
for the NASGLP

1 site/ 40km<sup>2</sup>



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# North American Soil Geochemistry Landscape Project

Earth Sciences Sector

- The NASGLP was developed from a 2001 meeting between the geological surveys of Mexico, USA and Canada for the purpose of characterizing the soil cover over North America.
- Goals of the NASGLP are:
  - Identify soil geochemical variability across the continent.
  - Identify soil geochemical variability throughout the soil profile.
  - Describe geochemical processes that are observed through the study of the geochemistry, mineralogy
  - Provide essential geochemical information for determining the range of background for a variety of soil materials across the continent.
  - Contribute to the requirements of both environmental and health risk assessments by providing realistic ranges for background.



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# GRTS Design - Maritime Provinces

Earth Sciences Sector



Natural Resources  
Canada



Ressources naturelles  
Canada

Canada




# Example - Other Sample Design

Earth Sciences Sector



Unbalanced Nested  
Stratified Sample Design  
(Garrett, 1983)



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# Sample Strategy - Decisions

Earth Sciences Sector

Target Size

Certainty

Resources



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# **Geochemical Landscapes**

## **Variation in the Vertical Profile**





# Soil Development

Earth Sciences Sector

## Soil Horizon:

A layer of mineral or organic soil material approximately parallel to the land surface that has characteristics altered by processes of soil formation. It differs from adjacent horizons in properties such as color, structure, texture, and consistence and *in chemical, biological or mineralogical composition*

**A-horizon:** forms at or near the surface in the zone of leaching or eluviation of materials (*removal of Fe, Al, clays and organic matter*) in solution or suspension

**B-horizon:** zone of enrichment in organic matter, sesquioxides, or clay; or by the development of soil structure; or by a change of color denoting hydrololysis, reduction, or oxidation

**C-horizon:** unaffected by the pedogenic processes operating in A and B-horizons

*Canadian System of Soil Classification, 3<sup>rd</sup> ed., AAFC*



Natural Resources  
Canada

Ressources naturelles  
Canada



Canada





# Generalized Soil Profile and Processes

Earth Sciences Sector



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# Soil Classification

Earth Sciences Sector

- Soils are classified according to Order, Great Group, and Sub-group.
- Information is available in Chapter 3 of Canadian System of Soil Classification (third edition).  
<http://sis.agr.gc.ca/cansis/taxa/cssc3/chpt3.html>.
- Soil stypes are differentiated on the basis of measured properties of the profile and uses a hierarchical scheme to classify soils from general to specific. The most recent version of the classification system has five categories in its hierarchical structure. From general to specific, the major categories in this system are: orders, great groups, subgroups, families, and series.
- At its most general level, the Canadian System of Soil Classification recognizes 9 different soil orders.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# Soil Orders in Canada

Earth Sciences Sector

The 9 soils orders are as follows:

- (1) Brunisol - is a normally immature soil commonly found under forested ecosystems. The most identifying trait of these soils is the presence of a B horizon that is brownish in color. The soils under the dry pine forests of south-central British Columbia are typically brunisols.
- (2) Chernozem - is a soil common to grassland ecosystems. This soil is dark in color (brown to black) and has an A horizon that is rich in organic matter. Chernozems are common in the Canadian prairies.
- (3) Cryosol - is a high latitudes soil common in the tundra. This soil has a layer of permafrost within one meter of the soil surface.
- (4) Gleysol - is a soil found in an ecosystem that is frequently flooded or permanently waterlogged. Its soil horizons show the chemical signs of oxidation and reduction.
- (5) Luvisol - is another type of soil that develops under forested conditions. This soil, however, has a calcareous parent material which results in a high pH and strong eluviation of clay from the A horizon.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# Soil Orders in Canada

Earth Sciences Sector

- (6) Organic - this soil is mainly composed of organic matter in various stages of decomposition. Organic soils are common in fens and bogs. The profiles of these soils have an obvious absence of mineral soil particles.
- (7) Podzol - is a soil commonly found under coniferous forests. Its main identifying traits are a poorly decomposed organic layer, an eluviated A horizon, and a B horizon with illuviated organic matter, aluminum, and iron. Podzols are common in the forested regions of Southern Ontario.
- (8) Regosol - is any young underdeveloped soil. Immature soils are common in geomorphologically dynamic environments. Many mountain river valleys in British Columbia have floodplains with surface deposits that are less than 3000 years old. The soils in these environments tend to be regosols.
- (9) Solonetzic - is a grassland soil where high levels of evapotranspiration cause the deposition of salts at or near the soil surface. Solonetzic soils are common in the dry regions of the prairies where evapotranspiration greatly exceeds precipitation input. The movement of water to the earth's surface because of capillary action, transpiration, and evaporation causes the deposition of salts when the water evaporates into the atmosphere.



Natural Resources  
Canada


Ressources naturelles  
Canada

Canada



# Common Soil Profiles

Earth Sciences Sector



Natural Resources  
Canada

Ressources naturelles  
Canada


Canada






# Podzol Profile

Earth Sciences Sector



\* *Well developed Ferro-Humic Podzol*



Soil Classification  
Working Group, 1998

LFH: ~5 cm  
Ah: ~4 cm  
Ae: ~10 cm  
Bhf: ~20 cm  
BC: ~20 cm  
C: >1.5 m



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada







# Brunisol Profile

Earth Sciences Sector

*\* Low degree of soil horizon development*



**LFH: ~5 cm**  
**Ah: <2 cm**  
**Bm: ~35 cm**  
**C: >40 cm**

Soil Classification  
Working Group, 1998



Natural Resources  
Canada

Ressources naturelles  
Canada


Canada





# Source Identification

Earth Sciences Sector



Tukey Box Plots for As, Pb in the <2mm fraction of soil from the 0-5 cm and 0-30 cm intervals and from the A,B, and C horizons and in the <0.063 mm fraction of the C horizon. Samples were analyzed by ICP-MS after a 4-acid digestion.



Natural Resources  
Canada

Ressources naturelles  
Canada



Canada



# As in A-horizon (<2 mm), based on CCME threshold levels

Earth Sciences Sector

**Red dots**  
– above limit  
**Green dots**  
– below limit



A-horizon, < 2mm fraction, EPA-3050B digestion



Universal Transverse Mercator - Zone 20 (N)  
Lon: 64°17.525' W  
Lat: 45°45.910' N  
Printed at: 2008-09-12



# As in C-horizon (<2 mm), based on CCME threshold level

Earth Sciences Sector

**Red dots**  
– above limit  
**Green dots**  
– below limit




C-horizon, < 2mm fraction, EPA-3050B digestion

Universal Transverse Mercator - Zone 20 (N)  
Lon: 64°15.364' W  
Lat: 45°45.947' N  
Printed at: 2008-09-12



# As in the Soil Profile By Horizons and Ecoregions

Earth Sciences Sector



As, <2 mm fraction  
EPA 3050B digestion  
ICP-MS analysis

Concentrations vary  
between horizons and  
between ecoregions



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada





# Recommendations

Earth Sciences Sector

- Choose a sampling area suitable for the risk assessment and reaching out to background.
- Choose sample density to optimize process recognition through a minimum number of elements and a maximum number of sample sites.
- Understand the regional geology, soil types, climate and geomorphology.
- Choose suitable soil horizons that reflect the anthropogenic impact and regional and local geochemical variability.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada



# References

Earth Sciences Sector

Ecological Stratification Working Group. 1996. A National Ecological Framework for Canada. Agriculture and Agri-Food Canada, Research Branch, Centre for Land and Biological Resources Research and Environment Canada, State of Environment Directorate, Ottawa/Hull. 125pp. And Map at scale 1:7.5 million.  
<http://www.ec.gc.ca/soer-ree/English/Framework/framework.cfm>. Pdf copy available from [/cansis/publications/ecostrat/intro.html](http://cansis/publications/ecostrat/intro.html)

Fulton, R.J. (comp.). 1995. Surficial Materials of Canada, Geological Survey of Canada, Map 1 880A, scale 1:8,000 000.

Garrett, R.G., 1983: Sampling methodology; in Chapter 4 of Handbook of Exploration Geochemistry, (ed.) R.J. Howarth,, Vol. 2, Statistics and Data Analysis in Geochemical Prospecting. Elsevier, p. 83-110.

Soil Classification Working Group, 1998. The Canadian System of Soil Classification; Agriculture and Agri-Food Canada Publication 1646 (third edition), 187 p.;  
<<http://www.pedosphere.com/resources/cssc3rd/>> [accessed December 16, 2009].

Soil Classification Working Group, 2000: Soils of Canada. Agriculture and Agri-Food Canada, map, scale 1:6 500 000.

Venables, W.N., and Ripley, B.D., 2002, Modern Applied Statistics with S, fourth Edition Springer-Verlag.

Wheeler, J O; Hoffman, P F; Card, K D; Davidson, A; Sanford, B V; Okulitch, A V; Roest, W R , 1996. Geological Map of Canada Map 1860A.Scale 1:5 000 000.



Natural Resources  
Canada

Ressources naturelles  
Canada

Canada