




Figure 1. View westward from ice-free summit of SE part of Angjak Island (sample sites 09-SRB-E108-112; 65°57'N, 62°2'W); 2011-058



Figure 2. View eastward into head of Hood Fjord (65°5'N, 63°5'W); 2011-060

Abstract
 In 2009, as part of the GEM program of the Geological Survey of Canada, Cumberland Peninsula east and southeast of the National Park was mapped. Mapping included several aspects of the regional Proterozoic geology as well as the Quaternary geology. Regional till was sampled for sedimentological and geochemical purposes, and moraines and other deposits were sampled for cosmogenic exposure dating. The eastern part of the peninsula is today an area of intense alpine glaciation and evidently has experienced this style of glaciation throughout the Quaternary. During the last glacial maximum (20–23ka), alpine glaciers advanced to form regional ice divides over the mountains. Ice flow from these divides covered most, possibly all, of the region and supported ice streams along several fronts. This local ice covered with the Laurentide ice Sheet in Cumberland Sound. Despite possibly repeated glaciation, substantial areas show little or no sign of glacial erosion and retain Tertiary surfaces marked with block fields and loess.

Résumé
 En 2009, dans le cadre du programme GEM, la Commission géologique du Canada a cartographié les régions de la péninsule de Cumberland à l'est et au sud-est du parc national. La cartographie inclut plusieurs aspects de la géologie régionale proterozoïque de même que la géologie du Quaternaire. Des échantillons de till régional ont été effectués à des fins d'analyses sédimentologiques et géochimiques ainsi que des moraines et d'autres dépôts à des fins de détermination des âges d'exposition aux rayonnements cosmogoniques. La partie est de la péninsule est aujourd'hui une région de glaciation alpine intense et de toute évidence, ce type de glaciation existait durant le Quaternaire. Durant le dernier maximum glaciaire (20–23ka), les glaciers alpins se sont épanchés pour former des lignes de partage glaciaire régionales. Les glaciers locaux ont recouvert la région et formé des courants glaciaires en bordure de glaciers frontaux. Cette glace locale est restée en coexistence avec l'indépendance lauridienne dans la baie Cumberland. Malgré plusieurs glaciations, des régions importantes ne montrent que très peu ou même aucun signe d'érosion glaciaire et retiennent les surfaces du Tertiaire recouvertes de champs de blocs et de loess.




Cover illustration: Uluksua Glacier, Cumberland Peninsula. Photograph by A. Dyke, 2009-250.
Printed map: Catalogue No. M183-117-2011E ISBN 978-0-660-20833-5
Digital map: Catalogue No. M183-117-2011E-PDF ISBN 978-1-100-18987-9
 © Her Majesty the Queen in Right of Canada 2011 doi:10.46955/28962

CANADIAN GEOSCIENCE MAP 17
 (preliminary version)
SURFICIAL GEOLOGY
HOARE BAY NORTH
 Baffin Island, Nunavut
 1:100 000



ess.nrcan.gc.ca
 Canadian Geoscience Maps
 Cartes géoscientifiques
 du Canada
 Canada



| SURFICIAL DEPOSITS | | PRE-WISCONSINAN | |
|---|--|------------------------|--|
| QUATERNARY | | PRE-WISCONSINAN | |
| HOLOCENE | | PRE-WISCONSINAN | |
| NEOGLACIAL | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: sand and gravel deposited in glacier-dammed lakes | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, deposited behind the ice margin | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, forming terraces, typically at maximum height of glacier-dammed lakes | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, deposited beyond the ice margin | | PRE-WISCONSINAN | |
| TILL: nonsorted glacial debris commonly very bouldery with a silty sand matrix | | PRE-WISCONSINAN | |
| LATERO-FRONTAL MORAINES: 5–100 m high moraine ridges with over-steepened, facing slopes or shallowly buried glacial ice cores and associated ground moraine with minor glaciolacustrine sediments. Distinguished from older moraines by lack of matrix tiller and peat covers. | | PRE-WISCONSINAN | |
| POSTGLACIAL (including Neoglacial) | | PRE-WISCONSINAN | |
| FLUVIAL SEDIMENTS: alluvium; gravel and sand deposited beyond primary influence of Holocene glaciers | | PRE-WISCONSINAN | |
| ALLUVIAL FANS: gravel and sand commonly bouldery, with detrital organic layers and hard soils, 1–20 m thick, formed by steep-gradient streams and debris flows forming channels and levees. | | PRE-WISCONSINAN | |
| ALLUVIAL PLAINS: gravel and sand, 1–10 m thick | | PRE-WISCONSINAN | |
| ALLUVIAL TERRACES: gravel and sand, 1–10 m thick, above limit of modern flooding | | PRE-WISCONSINAN | |
| COLLUVIUM: block and rubble accumulations, 1–50 m thick | | PRE-WISCONSINAN | |
| LANDSLIDE DEPOSITS: rock avalanche debris of coarse blocks, 10 or more metres thick, derived from cliff failure | | PRE-WISCONSINAN | |
| COLLUVIAL FAN DEPOSITS: locally to gravely debris-flow accumulations mixed with silt, 1–50 m thick, traversed by narrow channels and debris levees, typically internal slope steep (20°) at the ends of prominent debris fans; on all surface slopes less than angle of repose but steeper than those of alluvial fans. | | PRE-WISCONSINAN | |
| TALUS: generally active accumulations of blocks and rubble, as much as 50 m thick forming talus cones) sparse at angle of repose below cliffs derived from rock falls. | | PRE-WISCONSINAN | |
| ROCK GLACIER DEBRIS: talus, generally 10–50 m thick, defined by internal flow of buried ice to form tillite glaciers; irregular terraces on talus slopes with transverse ridges on tracks above steep frontal risers; some rock stable and well vegetated, most risers erode, unvegetated, and at angle of repose. | | PRE-WISCONSINAN | |
| MARINE SEDIMENTS: gravel, sand, silt, and minor clay, 1–20 m thick, deposited in beach, lacustrine, and offshore environments during regression of postglacial sea | | PRE-WISCONSINAN | |
| BEACH SEDIMENTS: gravel and sand, commonly bouldery, 1–5 m thick, forming raised beach ridges and washes and the modern, transgressive beach, a barrier beach in places. | | PRE-WISCONSINAN | |
| DELTAIC SEDIMENTS: sand and gravel, typically overlying fine sand and silt, bottomset beds, 5–20 m thick, formed in shallow, lacustrine at marine limit formed at or near the ice margin. | | PRE-WISCONSINAN | |
| EARLY HOLOCENE AND WISCONSINAN | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel, sand, silt, and minor clay, 1–10 m thick, deposited in littoral and deeper water environments in glacier-dammed lakes | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel, sand, silt, and minor clay, 1–2 m thick, deposited behind, at, and in front of the ice margin | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, deposited behind, at, and in front of the ice margin | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, forming fan-shaped deposits | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, forming reactive braided plains | | PRE-WISCONSINAN | |
| GLACIOLACUSTRINE SEDIMENTS: gravel and sand, 1–10 m thick, forming terraced deposits | | PRE-WISCONSINAN | |
| TILL: nonsorted bouldery diamictites, 1–40 m thick, deposited in subglacial and ice marginal environments, tillite composition generally reflecting underlying bedrock | | PRE-WISCONSINAN | |
| LATERO-FRONTAL MORAINES: 5–20 m high ridges and hummocks comprised mainly of till probably overlying older till; glacial ice cores, forming ridges and moraine ridges and less organized, hummocky accumulations formed during ice-marginal regression; moraine crests marked due to colluviation during partial deglaciation of ice cores; matrix somewhat more sandy and less silty than till forming ground moraine, locally contains ice-enclosed stratified drift and debris. | | PRE-WISCONSINAN | |
| TILL WEENER: variably bouldery (10–60% cover; typically 20–40%) diamiction with silty sand matrix, 0.5–2 m thick and discontinuous, insufficiently thick to obscure relief of underlying bedrock. | | PRE-WISCONSINAN | |
| TILL MANTLE: variably bouldery (10–60% cover; typically 20–40%) diamiction with silty sand matrix, 2–10 m thick, sufficiently thick to obscure relief of underlying bedrock. | | PRE-WISCONSINAN | |

CANADIAN GEOSCIENCE MAP 17
SURFICIAL GEOLOGY
HOARE BAY NORTH
 Baffin Island, Nunavut
 1:100 000

Map projection: Universal Transverse Mercator, zone 20, North American Datum 1983
 Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications.
 Elevation is feet above mean sea level.
 Mean magnetic declination 2011, 32°25'W, decreasing 28.4' annually. Readings vary from 32°59'W in the SE corner to 32°40'W in the NW corner of the map.

The Geological Survey of Canada welcomes appreciation or additional information from users. This map conforms to the ISO 9001:2008 standard used by the Scientific and Technical Publishing Services Quality Management System.
 This publication, including digital data, can be downloaded free of charge from GeoGratis (http://geogratis.nrcan.gc.ca). It is also available from the Geological Survey of Canada Bookstore (http://gsc.nrcan.gc.ca/bookstore).

Author: A.S. Dyke
 Geology by A.S. Dyke, 2009
 Geological compilation by A.S. Dyke, 2009
 Cartography by L. Robertson

Initiative of the Geological Survey of Canada, conducted under the auspices of the Multiscale Coastal and Marine (Nunavut) project, as part of Natural Resources Canada's Strategic Energy and Minerals (GEM) program.
 Logistical support provided by the Polar Continental Shelf Project as part of its mandate to promote scientific research in the Canadian North, PCSP 002-09

Recommended citation:
 Dyke, A.S., 2011. Surficial geology, Hoare Bay north, Baffin Island, Nunavut. Geological Survey of Canada, Canadian Geoscience Map 17 (preliminary version), scale = 1:100 000. doi:10.46955/28962

Recommended citation:
 Dyke, A.S., 2011. Surficial geology, Hoare Bay north, Baffin Island, Nunavut. Geological Survey of Canada, Canadian Geoscience Map 17 (preliminary version), scale = 1:100 000. doi:10.46955/28962

CANADIAN GEOSCIENCE MAP 17
 (preliminary version)
SURFICIAL GEOLOGY
HOARE BAY NORTH
 Baffin Island, Nunavut