


Figure 1. Head of Mermel Fiord. The sandy terrace is a wind-eroded, raised glaciomarine delta (sample sites 09-SRB-0371-073) at 36m elevation (66.3°N, 62.3°W), one of a few raised marine deposits along this coast. 2011-54

**Abstract**  
 In 2009, as part of the GEM Program of the Geological Survey of Canada, Cumberland Peninsula east and southeast of the National Park was mapped. Mapping included several aspects of the regional Precambrian geology as well as the Quaternary geology. Regional geology was sampled for sedimentological and geochemical purposes, and moraine and other deposits were sampled for cosmogenic exposure dating. The eastern part of the peninsula is today an area of intense alpine glaciation and evidently has experienced this style of glaciation throughout the Quaternary. During the last glacial maximum (28–20ka), alpine glaciers thickened to form regional ice divides over the mountains. Ice flow from these divides covered most, possibly all, of the region and supplied ice streams along several fiords. This local ice covered with the Laurentide Ice Sheet in Cumberland Sound. Despite possibly repeated glaciation, substantial areas show little or no sign of glacial erosion and retain Tertiary surfaces marked with block fields and tors.

**Résumé**  
 En 2009, dans le cadre du programme GEM, la Commission géologique du Canada a cartographié les régions est et sud-est de la péninsule de Cumberland à l'est et au sud-est du parc national. La cartographie inclut plusieurs aspects de la géologie régionale précambrienne de même que la géologie du quaternaire. Des échantillons de la région ont été effectués à des fins d'analyse sédimentologique et géochimiques ainsi que des moraines et d'autres dépôts à des fins de datation de l'exposition des âges d'exposition au rayonnement cosmogénique. L'ensemble de la péninsule est aujourd'hui une région de glaciation alpine intense et de toute évidence, ce type de glaciation existait durant le Quaternaire. Durant le dernier maximum glaciaire (28–20ka), les glaciers alpins se sont épaissis pour former des lignes de partage glaciaire au-dessus des montagnes. L'écoulement glaciaire provenait de ces lignes de partage glaciaire et a fourni des courants glaciaires en bordure de plusieurs fiords. Cette glace locale entraînait en conséquence avec l'Industrie laurontienne dans la baie Cumberland. Malgré plusieurs glaciations, des régions importantes ne montrent que très peu ou même aucun signe d'érosion glaciaire et conservent les surfaces du Tertiaire recouvertes de champs de blocs et de tors.



**Cover illustration**  
 Uluksuk Glacier, Cumberland Peninsula, Photograph by Art Dyke, 2002-255

**Printed map:**  
 Catalogue No. M183-1/20-2011E  
 ISBN 978-0-662-2089-7

**Digital map:**  
 Catalogue No. M183-1/20-2011E-PDF  
 ISBN 978-1-100-19075-4  
 doi:10.4065/289965


© Her Majesty the Queen in Right of Canada 2011

**CANADIAN GEOSCIENCE MAP 20**  
 (preliminary version)  
**SURFICIAL GEOLOGY**  
**CAPE DYER SOUTH**  
 Baffin Island, Nunavut  
 1:100 000



ess.nrcan.gc.ca

Canadian Geoscience Maps  
 Cartes géoscientifiques du Canada



SURFICIAL DEPOSITS	
<b>QUATERNARY</b>	
<b>HOLOCENE</b>	
<b>NEOGLACIAL</b>	
<b>hGf</b>	Proglacial outwash: gravel and sand, 1–10 m thick, forming fans.
<b>hGfp</b>	Proglacial outwash: gravel and sand 1–10 m thick, forming active terraces.
<b>Tt</b>	Noneroded glacial debris commonly very bouldery with a silty sand matrix.
<b>nTm</b>	Lateral-frontal moraines: 5–10 m high moraine ridges with over-stepped, facing slopes on shallowly buried glacial ice cores and associated ground moraine with more glacioluvial sediments, distinguished from older moraines by lack of mature lichen and plant covers.
<b>POSTGLACIAL (including Neoglaciation)</b>	
<b>AF</b>	Fluvial fans: gravel and sand commonly bouldery with detrital organic layers and buried soils, 1–20 m thick, formed by steep-gradient streams and debris flows forming channels and levees.
<b>Ap</b>	Alluvial plains: gravel and sand, 1–10 m thick.
<b>At</b>	Alluvial terraces: gravel and sand, 1–10 m thick, above limit of modern flooding.
<b>COLLUVIUM: block and rubble accumulations, 1–50 m thick</b>	
<b>Cl</b>	Colluvial fan deposits: blocky to gravelly debris-flow accumulations mixed with some, 1–50 m thick, intersected by narrow channels and debris levees; typically internal scarp slopes (Ca) at the ends of prominent debris hoppers in cliff; surface slopes less than angle of repose but steeper than those of alluvial fans.
<b>Ca</b>	Talus: generally active accumulations of blocks and rubble, as much as 50 m thick forming talus (scree) aprons at angle of repose below cliffs derived from rock falls.
<b>Cr</b>	Rock glacier debris: talus, generally 10–50 m thick, deformed by interstitial flow of buried ice to form talus glaciers; irregular terraces on talus slopes with transverse ridges on leads above leads; frost-free rises; some rises stable and well-vegetated; moist rises unstable, unvegetated, and at angle of repose.
<b>MARINE SEDIMENTS: gravel, sand, silt, and minor clay, 1–20 m thick, deposited in beach, deltaic, and offshore environments during regression of postglacial sea.</b>	
<b>Mr</b>	Beach sediments: gravel and sand, commonly bouldery, 1–5 m thick, forming raised beach ridges and sand dunes and the modern, transgressive beach, a barrier beach in places.
<b>Md</b>	Deltaic sediments: sand and gravel, typically overlying fine sand and silt bottomset beds, 5–20 m thick, forming raised terraces, terraces at marine limit formed at or near the ice margin.
<b>GLACIOLUVIAL SEDIMENTS: gravel and sand, 1–10m thick, deposited behind, at, and in front of the ice margin</b>	
<b>GFf</b>	Proglacial outwash: gravel and sand, 1–10 m thick, forming fan-shaped deposits.
<b>GFp</b>	Proglacial outwash: gravel and sand, 1–10 m thick, forming inactive terraced plains.
<b>GFt</b>	Proglacial outwash: gravel and sand, 1–10 m thick, forming terraced deposits.
<b>TILL: noneroded bouldery diamict, 1–40 m thick, deposited in subglacial and ice marginal environments, lithic composition generally reflecting underlying bedrock</b>	
<b>Tm</b>	Lateral-frontal moraines: 5–40 m high ridges and hummocks comprised mainly of till partially overlying debris-rich glacial ice cores, forming lateral and end moraine ridges and less organized, hummocky accumulations formed during ice-marginal recession; moraine crests muted due to consolidation during partial degradation of ice cores; matrix somewhat more sandy and less silty than till forming ground moraine; locally contains ice-contact stratified drift and outwash.
<b>Tr</b>	Rock-glacierized moraines: ice-cored end or lateral moraines, 5–40 m high, developed from original side of glacial deposition by downslope flow of ice-debris mixture; till mixed in places with some, commonly act as local base level for scree accumulation; hence difficult to distinguish from Cr in places; mainly stable rises.
<b>Tv</b>	Till veneer: vertically bouldery 10–60% cover, typically 20–40% diamictum with silty sand matrix, 0.5–2 m thick and discontinuous; insufficiently thick to obscure relief of underlying bedrock.
<b>Tb</b>	Till blanket: vertically bouldery (10–40% cover) typically 20–40% diamictum with silty sand matrix, 2–10 m thick, sufficiently thick to obscure relief of underlying bedrock.

**PRE-WISCONSINAN**  
**Ww** Residue: felsenmeer, rock rubble, and gravels, 1–2 m thick, mantling bedrock; formed by disintegration of Precambrian bedrock prior to Last Glacial Maximum but including sparse erratics and possibly morphologically degraded till; tonalite felsenmeer composed of metre-scale blocks in diamict matrix with typically 60–80% block cover; finer rubble and gravels on coarse monograptite of Cumberland Batholith; intersected by 1–4 m high, bedrock surfaces shows little or no sign of glacial scouring; horizon retains its pre-Quaternary form, mainly on fill or gently graded, cliff-bordered, upland plateaus, but on moderately steep slopes in places, most surfaces probably covered by cold-based ice during one or more glaciations.

**BEDROCK**

**PRE-QUATERNARY**  
**R1** Gneiss: major outcrops, typically hundreds of metres high, forming serrated faces with multiple debris hoppers; glacially scoured surfaces removed by postglacial scarp retreat prograding basal (or sub-basinal) talus accumulations.  
**R2** Scoured rock: hilly and hummocky surfaces with lake basins and ice moulded eminences resulting from glacial scouring and with patchy veneers of till commonly deposited of matrix material probably covered by warm-based ice during stable intervals of Wisconsin Glaciation, including Last Glacial Maximum.

- Glacier ice margin for 2009 from SPOT5 satellite imagery
- Geological boundary (defined)
- Geological boundary (gridstone)
- Fluted bedrock (direction known)
- Fluted bedrock (direction unknown)
- Drumlin
- Lateral-frontal moraine
- Lateral meltwater channel: bar on upslope side
- Proglacial meltwater channel
- Lacustrine limit
- Cirque
- Arête
- Cliff
- Dyke
- Field observation (point count relative to geological units)
- Strike (ice flow direction known)
- Station, marine shoreline elevation in metres

ess.nrcan.gc.ca

Canadian Geoscience Maps  
 Cartes géoscientifiques du Canada

**Preliminary version** **CANADIAN GEOSCIENCE MAP 20** **Preliminary version**

**SURFICIAL GEOLOGY**  
**CAPE DYER SOUTH**  
 Baffin Island, Nunavut  
 1:100 000

Author: A.S. Dyke  
 Geology by A.S. Dyke, 2009  
 Geological compilation by A.S. Dyke, 2009  
 Cartography by J. Robertson

Initiative of the Geological Survey of Canada, conducted under the auspices of the Multiple Metals Cumberland Peninsula (Nunavut) project, in part of Natural Resources Canada's Geotouring for Energy and Minerals (GEM) program.  
 Logistical support provided by the Polar Continental Shelf Project as part of its mandate to promote scientific research in the Canadian North. PCSP 002-09

Map projection Universal Transverse Mercator: zone 20, North American Datum 1980  
 Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications.  
 Elevations in feet above mean sea level  
 Mean magnetic declination 2011, 33°54'W, decreasing 20' annually. Readings vary from 33°50'W in the SE corner to 34°16'W in the NW corner of the map.

The Geological Survey of Canada welcomes corrections or additional information from users. This map conforms to the ISO 9001:2008 standard used by the Scientific and Technical Publishing Services Quality Management System.  
 This publication, including digital data, can be downloaded free of charge from GeoPub (<http://geopub.nrcan.gc.ca/>). It is also available from the Geological Survey of Canada Bookstore (<http://psc.nrcan.gc.ca/bookstore/>).

Recommended citation:  
 Dyke, A.S., 2011. Surficial geology, Cape Dyer south, Baffin Island, Nunavut, Geological Survey of Canada, Canadian Geoscience Map 20 (preliminary version), scale 1:100 000, doi:10.4065/289965

**CANADIAN GEOSCIENCE MAP 20**  
 (preliminary version)  
**SURFICIAL GEOLOGY**  
**CAPE DYER SOUTH**  
 Baffin Island, Nunavut

**Preliminary version** **CANADIAN GEOSCIENCE MAP 20** **Preliminary version**

**SURFICIAL GEOLOGY**  
**CAPE DYER SOUTH**  
 Baffin Island, Nunavut