



**GEOLOGICAL SURVEY OF CANADA
OPEN FILE 7202**

**Geochemical and isotopic (Nd, O, Pb and Sr) data from
igneous rocks of the Notre Dame subzone and adjacent
tectonostratigraphic zones, western and central
Newfoundland**

J.B. Whalen

2012



Natural Resources
Canada

Ressources naturelles
Canada

Canada



**GEOLOGICAL SURVEY OF CANADA
OPEN FILE 7202**

**Geochemical and isotopic (Nd, O, Pb and Sr) data from
igneous rocks of the Notre Dame subzone and adjacent
tectonostratigraphic zones, western and central
Newfoundland**

J.B. Whalen

2012

©Her Majesty the Queen in Right of Canada 2012

doi:10.4095/291593

This publication can be downloaded free of charge from GEOSCAN
(<http://geoscan.ess.nrcan.gc.ca/>).

Recommended citation

Whalen, J.B., 2012. Geochemical and isotopic (Nd, O, Pb and Sr) data from igneous rocks of the Notre Dame subzone and adjacent tectonostratigraphic zones, western and central Newfoundland; Geological Survey of Canada, Open File 7202. doi:10.4095/291593

Publications in this series have not been edited; they are released as submitted by the author.

Geochemical and isotopic (Nd, O, Pb and Sr) data from igneous rocks of the Notre Dame subzone and adjacent tectonostratigraphic zones, western and central Newfoundland.

Joseph B. Whalen

Geological Survey of Canada, 601 Booth Street, Ottawa, Ontario, K1A 0E8, Canada.

Email: jwhalen@nrcan.gc.ca

The following report presents a compilation of geochemical and isotopic data collected in the course of a numbers of projects starting in 1981 with geological mapping and geochemical sampling of the Silurian Topsail igneous suite (TIS) (Whalen and Currie, 1988; Whalen et al., 1987a,b,c). A Canada-Newfoundland Cooperation Agreement (CNCA) (1990-1994) funded project focused mainly on Ordovician plutonism within the Notre Dame subzone (Fig. 1) (Whalen et al., 1997b) but included isotopic work on the TIS (Whalen et al., 1996). During the Red Indian Line Project, which was funded under the auspices of the 2000 to 2003 Targeted Geoscience Initiative (TGI), more extensive work was carried out within the southern Notre Dame subzone and the northern Dashwoods subzone (Whalen et al., 2006; van Staal et al., 2007, 2008). With funding from the 2005-2010 Targeted Geoscience Initiative 3 Program (TGI3) some further geochemical and isotopic work was carried out in the Buchans-Robert's Arm belt and the Burlington Peninsula, the results of which are included in this report.

The geochemical data presented herein should provide significant assistance in the regional correlation of units, as well as for constraining their tectonic setting and igneous petrogenesis. It complements previously released geochemical databases for central Newfoundland (e.g., Davenport et al., 1996; Jenner, 2002; Rogers, 2004; Zagorevski and Rogers, 2011).

Tectonic and geologic context

This brief overview is mainly summarized from van Staal et al. (1998, 2004, 2009) and van der Velden et al. (2004), and references therein. The Newfoundland Appalachians are interpreted as a relic collision zone between Laurentia, in the west, represented by Grenvillian basement to the Humber Zone, and Gondwanan terranes in the east, represented by Neoproterozoic basement rocks to parts of the Exploits subzone, and the Gander and Avalon zones (see Fig. 1). The western margin of Iapetus (Humber Zone), an Early Paleozoic passive margin sequence deposited on the leading edge of Laurentia, is divided into western external and eastern internal

sectors, based on increasing deformation and metamorphism toward the east. Prior to 470-475 Ma, a narrow oceanic tract (Humber seaway) is thought to have separated the Humber margin from the Notre Dame arc (NDA) (Notre Dame plus Dashwoods subzones in Fig. 1). This continental magmatic arc is thought to have evolved on a ribbon continent, referred to as Dashwoods, which rifted from Laurentia between 550-530 Ma. Closure of the Humber seaway along the Baie Verte Line (BVL) by east-directed subduction beneath Dashwoods was accompanied by ca. 475 Ma obduction of the Tremadoc-early Arenig suprasubduction zone rocks of the Baie Verte oceanic tract onto the Humber margin and 488-480 Ma arc magmatism of the NDA. It was followed by voluminous tonalite plutonism between 466-459 Ma within the NDA, attributed to break-off of the oceanic slab attached to the downgoing Humber margin. Closure of the main Iapetan ocean basin was accomplished during the Caradoc (455-450 Ma) by simultaneous east- and west-directed subduction. These processes juxtaposed the peri-Laurentian NDA and the associated Annieopsquotch accretionary tract (AAT) with the peri-Gondwanan Victoria arc along the Red Indian Line (RIL) (Fig. 1). Accretion of the Victoria arc to composite Laurentia was followed (1) by accretion of the Gander zone along the Dog Bay Line, as a result of west-directed subduction of the Exploits back-arc basin, producing the Silurian Salinic orogeny, and (2) by accretion of the Avalon Zone along the Dover-Hermitage Bay fault, producing the Early Devonian Acadian orogeny.

Early Silurian magmatism and sedimentation west of the RIL, within the Notre Dame subzone, form stitching plutons and overlap basins on the accreted terranes. These events overlap in time with a major pulse between ca. 435-425 Ma of regional deformation plus metamorphism (Salinic orogeny) followed by rapid uplift and cooling within the Humber internal zone. Overall, this activity has been interpreted as reflecting Exploits back-arc closure and subsequent slab break-off.

This report presents a compilation of geochemical and isotopic data collected mainly from Ordovician and Silurian igneous rocks of the Notre Dame and Dashwoods subzones of western Newfoundland (see Fig. 2 for sample distribution). Interpretations based on data contained in this compilation were fundamental to our current understanding of the complex accretionary events that are outlined above.

Geochemical Database

This open file contains 607 whole rock geochemical samples presented as Microsoft Excel[®] worksheet with the shared link between files being sample number. These samples

comprise surface silicate rock samples of mainly plutonic, lesser volcanic and minor epiclastic rocks. The background information is presented as Adobe Acrobat[®], Microsoft Word[®] documents or Microsoft Excel[®] tables, as appropriate, with the data type indicated by the file names. Background information includes analytical procedures and unfiltered data tables, as produced by the laboratories (these tables include repeats, standards, etc.), for samples which were analyzed post-1989. Such lab sourced files are not available for samples that were analyzed pre-1989. These files are grouped in folders named after the project or program during which sampling and analytical work was carried out: Topsail project, Canada-Newfoundland Cooperation Agreement (CNCA) (1990-1994) project, Red Indian Line project and Target Geoscience Initiative (TGI) 3 project. Samples with a WX-prefix were collected, submitted for analysis and interpreted by the author. Samples with HPA-, MRB-, PQB-, VL-, VL-LH-, and VO- prefixes were submitted for analysis and interpreted by the author. Samples with VL01A- and VL01J- were interpreted by the author.

Location Table: The location table contains the sample location data in the UTM, easting and northing (Zone 21; NAD27) format. Locations for samples collected between 1981 and 1994 were compiled from 1:50 000 topographic sheets. Location data for samples collected post-2000 were determined by handheld GPS. In addition to the exact position for each sample, an overall sense of the location of the sample can be deduced from the NTS map sheet name and number. Sample distribution relative to NTS sheets is shown in Figure 2.

Description Table: The description table contains a number of columns that the user can employ to sort or group the data. The column ‘Zone’ is the Newfoundland tectonostratigraphic zone, based on Williams et al. (1988) (see Fig. 1 for key to abbreviations), in which the sample was collected. This is followed by ‘Type’ (volcanic, plutonic or sedimentary) and ‘Subtype’ (mafic (<55 wt. % SiO₂), intermediate (>55 and <65 wt. % SiO₂) or felsic (>65 wt. % SiO₂); this column is blank for sedimentary rocks. The next column gives the name of the person who collected the sample. The following two columns ‘Suite, Complex or Pluton Name’ and ‘Map Unit’ were derived from published geological maps given in the ‘Reference’ column and listed near the end of this document. These maps can be downloaded for free from this NRCan web site (http://apps1.gdr.nrcan.gc.ca/mirage/db_search_e.php). The final column in this table is a free-form rock-type description. Abbreviations employed in these descriptions are as follows:

vfg = very fine grained	amph = amphibole
fg = fine grained	bt = biotite
mg = medium grained	chl = chlorite
cg = coarse grained	cpx = clinopyroxene
vcg = very coarse grained	ep = epidote

eq = equigranular	feld = feldspar
porph = porphyritic	hbl = hornblende
mass = massive	ms = muscovite
fol = foliated	ol = olivine
incl = inclusions	plag = plagioclase
alt = alteration	px = pyroxene
comp = compositional	qtz = quartz
bk = black	kfld = K-feldspar
dk = dark	mag = magnetite
pk = pink	opx = orthopyroxene
wt = white	sl = sillimanite
	st = staurolite

Chemistry Table: The whole rock chemistry data file is presented as a Microsoft Excel[®] worksheet file which contains whole-rock geochemical analytical data. Major elements are recorded as weight percentages of their oxides. Where the oxidation state was determined, iron is presented as FeO and Fe₂O₃, otherwise it is represented as FeO (total). Volatiles are represented as either CO₂ and H₂O or LOI (loss-on-ignition). The minor, trace and rare earth elemental compositions are presented in parts per million. Details of the analytical procedures are presented in the background information folder. Where an element was analysed using multiple methods the value determined by the method that appears most reliable is presented. Elements not determined are given as 0, whereas element analyses that are below the detection limits are presented as a negative value with the number being the detection limit. A large proportion of the geochemical data presented in this report has been previously presented and interpreted in earlier publications given in the reference list including van Staal et al. (2007, 2008), Whalen (1986, 1989), Whalen and Currie (1984, 1988, 1990), Whalen et al. (1987a, 1987b, 1996, 1997a, 1997b, 2006). Also there is overlap in geochemical data presented herein with that contained in the compilation of Rogers (2004).

Isotopes Table: The isotope table contains: whole rock ‘Delta18O’ ($\delta^{18}\text{O}\%$ SMOW); whole rock ‘i87Sr/86Sr’ (initial $^{87}\text{Sr}/^{86}\text{Sr}$); ‘m87Sr/86Sr’ (measured $^{87}\text{Sr}/^{86}\text{Sr}$); ‘i206Pb/204Pb’ (initial $^{206}\text{Pb}/^{204}\text{Pb}$); ‘i207Pb/204Pb’ (initial $^{207}\text{Pb}/^{204}\text{Pb}$); ‘i208Pb/204Pb’ (initial $^{208}\text{Pb}/^{204}\text{Pb}$) (all determined on K-feldspar); whole rock ‘ENd(460)’ (ϵ_{Nd} at 460 Ma); ‘ENd(T)’ (ϵ_{Nd} calculated at age given in column ‘Age (T)’ in Ma (some ages are determined U-Pb zircon ages some interpreted estimates); ‘TDM(Gold)’ (Nd age in Ga based on model of Goldstein et al. (1984)); ‘TDM(DeP)’ (Nd age in Ga based on model of DePaolo (1981)); ‘(143/144)m’ ($^{143}\text{Nd}/^{144}\text{Nd}$ measured); ‘(147/144)m’ ($^{147}\text{Sm}/^{144}\text{Sm}$ measured); ‘(143/144)i’ ($^{143}\text{Nd}/^{144}\text{Nd}$ initial); ‘CHURi’ (CHUR initial); and Nd and Sm in ppm by isotope dilution. Model ages were not calculated

(NC) for samples for which $(^{147}\text{Sm}/^{144}\text{Sm})_{\text{m}}$ was greater than 0.15. To enable easy linkage between tables, the many samples for which no isotopic data is available are included as blank rows. Details of the isotopic analytical techniques and labs where determinations were made are given in the analytical procedures. Almost all the isotopic data presented in this report has been previously interpreted and published in scientific journal papers given in the reference list (van Staal et al., 2007, 2008; Whalen et al., 1987c, 1996, 1997a, 1997b, 2006). As well there is overlap in Nd isotopic data presented herein with that contained in the compilation of Rogers (2004).

Complete Table: This file combines location, description, rock chemistry and isotopes tables.

Discussion of geochemical data

Samples utilized in this study were collected in the course of regional scale mapping in order to identify the geochemical fingerprints of different igneous suites within tectonostratigraphic terranes and to establish geochemical-isotopic characteristics of terranes to aid regional correlations. A discussion of the tectonic significance, distribution and correlation of the volcano-plutonic rocks presented in this database is beyond the scope of this report. For this the reader is referred to various references by the author and van Staal et al. given in the references at the end of this report. Herein, the intent is to highlight the diversity of rock types included in this database and the range of tectonic settings within which they likely were generated.

The geochemical data from igneous rocks was subdivided based on silica content into three groups; (1) mafic ($\text{SiO}_2 < 55$ wt.%); (2) intermediate ($\text{SiO}_2 > 55, < 65$ wt.%); and, (3) felsic ($\text{SiO}_2 > 65$ wt.%). A plot of all these groups on the Q'-ANOR plot of Streckeisen and LeMaitre (1979) (Fig. 2a) illustrates the large diversity of plutonic rock types included in the dataset. As cogenetic igneous suites normally do not cut across boundaries in a K_2O - SiO_2 plot (Fig. 2b), this plot shows that the dataset includes shoshonitic, high-K, medium-K and low-K igneous suites. Intermediate and felsic igneous rocks are plotted on Rb-Y+Nb and Nb-Y granitoid rock tectonomagmatic discrimination plots (Fig. 3a and 3b) which show that the dataset consists mainly of volcanic arc, within-plate and post-collisional granites. Mafic igneous rocks are plotted on various basalt trace element tectonomagmatic discrimination plots (Fig. 4) which suggest the dataset consists of basaltic rocks that formed within diverse arc-related settings. For a discussion

of the isotopic data and its interpretation, the reader is referred to Whalen et al. (1987c, 1996, 1997a,b, and 2006).

Acknowledgements

This is a contribution to the Targeted Geoscience Initiative (TGI) 4 (2010-2015). The following are acknowledged for providing access to their unpublished geochemical data: Johan Lissenberg (Cardiff University, Wales), Sally Pehrsson (Geological Survey of Canada) and Alex Zagorevski (Geological Survey of Canada). Additional acknowledgement is also given to Neil Rogers for his contribution as project leader for the TGI 3 and TGI 4 programs in Newfoundland and for critically reviewing this open file.

References

- Brem, A.G., Lin, S., and van Staal, C.R., 2006, Geology, Harrys River, Newfoundland and Labrador, Geological Survey of Canada, Open File 4921, scale 1:50 000.
- Cabanis, B., and Lecolle, M., 1989, Le diagramme La/10-Y/15-Nb/8; un outil pour la discrimination des series volcaniques et la mise en evidence des processus de melange et/ou de contamination crustale. The La/10-Y/15-Nb/8 diagram; a tool for distinguishing volcanic series and discovering crustal mixing and/or contamination, Comptes Rendus de l'Academie des Sciences, Serie 2, Mecanique, Physique, Chimie, Sciences de l'Univers, Sciences de la Terre, v. 309, no. 20, p. 2023-2029.
- Davenport, P. H., Honarvar, P., Hogan, A., Kilfoil, G., King, D., Nolan, L. W., Ash, J. S., Colman-Sadd, S. P., Hayes, J. P., Liverman, D. G. E., Kerr, A., and Evans, D. T. W., 1996, Newfoundland Department of Mines and Energy, Geological Survey, Open File NFLD/2611 (1996) version 1.0, CD-ROM (1 disk).
- DePaolo, D.J., 1981, Neodymium isotopes in the Colorado Front Range and implications for crust formation and mantle evolution in the Proterozoic, *Nature*, v. 291, p. 193-197.
- Goldstein, S.L., O'Nions, R.K., and Hamilton, P.J., 1984, A Sm-Nd isotopic study of atmospheric dusts and particulates from major river systems, *Earth and planetary Science letters*, v. 70, p. 221-236.
- Hall, L.A.F., and van Staal, C.R., 1999, Geology, Southern End of Long Range Mountains (Dashwoods Subzone), Newfoundland, Geological Survey of Canada, Open File 3727, scale 1:50 000.
- Jenner, G. A., 2002, Assessment report on geochemical exploration for 2001 submission for fee simple grants volume 1 folios 61-62 and for second year supplementary, fourth year supplementary, fifth year, sixth year supplementary, seventh year and ninth year supplementary assessment for licence 4805 on claim 16398, licence 4823 on claims 16431-16432, licence 4867 on claims 16397, 16400-16401, 16424-16426 and 17688, licence 4868 on claim block 6648, and licences 5576M, 5649M, 5668M, 6003M, 7420M, 8295M, 8312M and 8444M on claims in the Buchans area, central Newfoundland, Newfoundland and Labrador Geological Survey, Assessment File 12A/1008, p. 131.
- Le Maitre, R.W., 1989, *A Classification of Igneous Rocks and Glossary of Terms*, Oxford, Blackwell, 193 pp.
- Lissenberg, C.J., Zagorevski, A., McNicoll, V.J., van Staal, C.R., and Whalen, J.B., 2005, Assembly of the Annieopsquotch accretionary tract, Newfoundland Appalachians: Age and geodynamic constraints from syn-kinematic intrusions, *Journal of Geology*, v.113, p.553-570.

- Lissenberg, C.J., Zagorevski, A., Whalen, J.B., and van Staal, C.R., 2005, Geology, Star Lake, Newfoundland (NTS 12A/11), Geological Survey of Canada Open File 1669, 1:50,000.
- Owen, J.V., 1990, Geology, Long Range Inlier, Newfoundland, Geological Survey of Canada, Map 1764A, scale 1:250 000.
- Pearce, J.A., 1996, Sources and settings of granitic rocks, *Episodes* 19, 120-125.
- Pearce, J.A., Harris, N.B.W., and Tindle, A.G., 1984, Trace element discrimination diagrams for the tectonic interpretation of granitic rocks, *Journal of Petrology* 25, 956-983.
- Peccerillo, A., and Taylor, S.R., 1976, Geochemistry of the Eocene calc-alkaline volcanic rocks from Kastamonu area, northern Turkey, *Contributions to Mineralogy and Petrology*, v. 58, p. 63-81.
- Piasecki, M.A.J., Williams, H., and Coleman-Sadd, S.P., 1990, Tectonic relationships along the Meelpaeg, Burgeo and Burlington LITHOPROBE transects in Newfoundland, *In* Current Research, Newfoundland Department of Mines and Energy, Geological Survey Branch, Report 90-1, p. 327-339.
- Rogers, N., 2004, Red Indian Line geochemical database, Geological Survey of Canada Open File 4605.
- Shervais, J. W., 1982, Ti-V plots and the petrogenesis of modern and ophiolitic lavas, *Earth and Planetary Science Letters*, v. 59, no. 1, p. 101-118.
- Skulski, T., Castonguay, S., Moussallam, Y., McNicoll, V., van Staal, C., and Bedard, J., in review, Geology Nippers Harbour and parts of Horse Islands, Cape St. John and Little Bay Island, Newfoundland and Labrador, Geological Survey of Canada, Canadian Geoscience Map 15528, scale 1:50,000.
- Streckeisen, A.L., and LeMaitre, R.W., 1979, Chemical approximation to modal QAPF classification of the igneous rocks, *Neues Jahrbuch für Mineralogie*, v. 136, p. 169-206.
- Szybinski, Z. A., Brem, A. G., van Staal, C. R., Whalen, J. B., McNicoll, V. J., Jenner, G., and Piercey, S., 2006, Geology, Little Grand Lake, Newfoundland (NTS 12A/12), Geological Survey of Canada Open File 1668, scale 1:50,000.
- van Staal, C.R., Lin, S., Hall, L., Valverde, P., and Genkin, 1996, Geology, Rose Blanche, Newfoundland (part of NTS 11-O/11), Geological Survey of Canada, Open File 3165, scale 1:50 000.
- van Staal, C.R., Lissenberg, C.J., Pehrsson, S.J., Zagorevski, A., Valverde-Vaquero, P., Herd, R.K., McNicoll, V., and Whalen, J.B., 2005a, Geology, Puddle Pond, Newfoundland, (12A/05), Geological Survey of Canada Open File 1664, 1:50,000.
- van Staal, C.R., Valverde-Vaquero, P., Zagorevski, A., Pehrsson, S., Boutsma, S., and van Noorden, M.J., 2005b, Geology, King George IV Lake, Newfoundland and Labrador, Geological Survey of Canada, Open File 1665, scale 1:50 000.
- van Staal, C.R., Valverde-Vaquero, P., Zagorevski, A., Rogers, N., Lissenberg, C.J., and McNicoll, V.J., 2005c, Geology, Victoria Lake, Newfoundland and Labrador, Geological Survey of Canada, Open File 1667, scale 1:50 000.
- van Staal, C.R., Whalen, J.B., McNicoll, V.J., Pehrsson, S., Lissenberg, C.J., Zagorevski, A., van Breemen, O., and Jenner, G., 2007, The Notre Dame Arc and the Taconic Orogeny in Newfoundland, *In* Hatcher, R.D., Jr., Carlson, M.P., McBride, J.H., and Martinez Catalan, J.R., eds., 4-D Framework of Continental Crust; Geological Society of America Memoir 200, p. 511-552.
- van Staal, C.R., Whalen, J.B., Valverde-Vaquero, P., Zagorevski, A., and Rogers, N., 2009, Pre-Carboniferous, episodic accretion-related, orogenesis along the Laurentian margin of the northern Appalachians, *In* Murphy, J.B., Keppie, J.D., and Hynes, A.J., eds., Ancient Orogens and Modern Analogues. Geological Society, London, Special Publication; v. 327; p. 271-316.
- Van Staal, C.R. and Zagorevski, A., 2012. Accreted terranes of the Appalachian Orogen in Newfoundland: In the footsteps of Hank Williams, Geological Association of Canada, Guidebook Trip A1, 99 p.

- Whalen, J.B., 1986, Geochemistry of the mafic and volcanic components of the Topsails igneous suite, Western Newfoundland, In Current Research, Part B, p. 121-130, Geological Survey of Canada, Paper 86-1B.
- Whalen, J.B., 1989, The Topsails suite, western Newfoundland: an early Silurian subduction-related magmatic suite? Canadian Journal of Earth Sciences, v. 26, p. 2421-2434.
- Whalen, J.B., and Currie, K.L., 1984, The Topsails igneous terrane, Western Newfoundland: evidence for magma mixing, Contributions to Mineralogy and Petrology, v. 87, p. 319-327.
- Whalen, J.B., and Currie, K.L., 1988, Geology of the Topsails igneous terrane, western Newfoundland, Geological Survey of Canada Map 1680A, scale 1:200 000.
- Whalen, J.B., and Currie, K.L., 1990, The Topsails suite, western Newfoundland: fractionation and magma-mixing in an "orogenic" A-type granite suite, In Stein, H.J. and Hannah, J. L. (Editors), Ore-bearing Granite Systems; Petrogenesis and Mineralizing Processes, Geological Society of America, Special Paper 246, p. 287-299.
- Whalen, J.B., Currie, K.L., and Chappell, B.W., 1987a, A-type granites: geochemical characteristics, discrimination and petrogenesis, Contributions to Mineralogy and Petrology, v. 95, p. 407-419.
- Whalen, J.B., Currie, K.L., and Chappell, B.W., 1987b, A-type granites: descriptive and geochemical data, Geological Survey of Canada Open File 1411.
- Whalen, J.B., Currie, K.L., and van Breeman, O., 1987c, Episodic Ordovician-Silurian plutonism in the Topsails igneous terrane, western Newfoundland, Transactions of the Royal Society of Edinburgh: Earth Sciences, v. 78, p. 17-28.
- Whalen, J.B., Longstaffe, F.J., Jenner, G.A., Robert, F., and Gariépy, C., 1996, Geochemical and isotopic (O, Nd, Pb and Sr) constraints on A-type granite petrogenesis based on the Topsails igneous suite, Newfoundland Appalachians, Journal of Petrology, v. 37, p.1463-1489.
- Whalen, J.B., van Staal, C.R., Longstaffe, F.J., Gariépy, C., and Jenner, G.A., 1997a, Insights into tectonostratigraphic zone identification in southwestern Newfoundland based on isotopic (Nd, O, Pb) and geochemical data, Atlantic Geology, v.33, no.4, p.231-241.
- Whalen, J.B., Jenner, G.A., Longstaffe, F.J., Gariépy, C., and Fryer, B.J., 1997b, Implications of granitoid geochemical and isotopic (Nd, O, Pb) data from the Cambro-Ordovician Notre Dame arc for the evolution of the Central Mobile Belt, Newfoundland Appalachian, In Sinha, A.K., Whalen, J.B., and Hogan, J., eds., Magmatism in the Appalachian Orogen; Geological Society of America Memoir 191, p. 367-395.
- Whalen, J.B., McNicoll, V.J., van Staal, C.R., Lisenberg, C.J., Longstaffe, F.J., Jenner, G.A., and van Breeman, O., 2006, Spatial, temporal and geochemical characteristics of Silurian collision-zone magmatism: an example of a rapidly evolving magmatic system related to slab break-off, Newfoundland Appalachians, Lithos, v.89, p.377-404.
- Williams, H., Colman-Sadd, S.P., and Swinden, H.S., 1988, Tectonic - stratigraphic subdivisions of central Newfoundland, In Current Research, Part B. Geological Survey of Canada, Paper 88-1B, p. 91-98.
- Wood, D.A., 1980, The application of a Th-Hf-Ta diagram to problems of tectonomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of the British Tertiary volcanic province, Earth and Planetary Science Letters, v.50, p.11-30.
- Zagorevski, A., and Rogers, N., 2011, Geochemical Data of the Buchans-Robert's Arm Belt and adjacent units in central Newfoundland and Notre Dame Bay regions, Geological Survey of Canada, Open File 6728, 1CD-ROM.




Figure 1. Tectonostratigraphic zones and subzones of the Newfoundland Appalachians (modified after Williams et al. (1988) and Piasecki et al. (1990)). Abbreviations used for zones in the 'Description Table' are given in the figure legend; RIL = Red Indian Line.




Figure 2. Sample distribution within NTS sheets of western Newfoundland.




Figure 3. Western Newfoundland igneous rocks, grouped by silica content, plotted on: (a) the Q'-ANOR CIPW normative plutonic rock classification diagram of Streckeisen and LeMaitre (1979); and (b) the K₂O versus SiO₂ diagram of Le Maitre (1989) with shoshonitic field from Peccerillo and Taylor (1976). Samples were grouped based on hydrous analyses and plotted anhydrous, thus some samples plot at higher silica content than the grouping parameters.






Figure 4. Western Newfoundland intermediate and felsic igneous rocks plotted on the (a) Rb versus Y+Nb diagram of Pearce (1996); and (b) the Nb versus Y diagram of Pearce et al. (1984). Abbreviations are: syn-COLG and post-COLG, syn- and post-collisional granites; VAG, volcanic arc granites; ORG, ocean ridge granites; WPG, within plate granites.




Figure 5. Western Newfoundland mafic (<55 wt.% SiO₂) igneous rocks plotted on; (a) the Th-Zr-Nb diagram of Wood (1980); (b) the La-Y-Nb diagram of Cabanis and Lecolle (1989); (c) the V-Ti diagram of Shervais (1982). Abbreviations: VAT, volcanic arc tholeiite; N- and E-MORB, normal and enriched mid-ocean ridge basalt; CAB, calc-alkaline basalt; CON, continental tholeiite; alkali con rift, alkaline continental rift basalt; BAB, back-arc basalt.

Contents

readme.rtf – contents and file structure.

TGI4 NL Geochemical-Isotopic Database.pdf – summary document outlining the database.

licence_agreement.rtf – formal end-user licence agreement for digital data.

Primary Data Files

Complete.xls –location, description, chemistry and isotopes of silicate samples

Locations.xls – location of samples included in this open file

Description.xls – descriptive information on the samples included in this open file

Chemistry.xls – whole-rock geochemical data on silicate samples

Isotopes.xls – isotopic data on silicate samples

Background Information

Analytical Procedures

analytical_techniques.pdf

Standards_duplicates.xls

Unprocessed Analytical Data

CNCA lab files

XRAL major elements.xls

MUN XRF on powder pellets.xls

MUN ICPMS traces 1.xls

MUN ICPMS traces 2.xls

Red Indian Line lab files

02037xtr.xls

02064XTR.xls

03002xem.xls

01053.xls

TL0001-0048-SmNd calc.xls

TGI3 lab files

07013XRF.xls

07013MS.xls

0713ET-WHA.xls

O7013XRF_C_complete.xls

CarletonU_Nd_results.xls