

DESCRIPTIVE NOTES

INTRODUCTION
The Surficial Geology Map of NTS 94-004 (Canadian Geoscience Map 108) is the product of collaboration between the Geological Survey of Canada and British Columbia's Ministry of Energy, Mines and Natural Gas as part of the Geo-Mapping for Energy and Minerals Program (GEM-EM) Yukon Basin Project.

APPROACH TO SURFICIAL GEOLOGY MAPPING
The distinctive landscape of NTS 94-004 has led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-004 map area (Hurley and Hicks, 2010; Hurley et al., 2011a,b).

INFERRED GEOLOGICAL HISTORY
The distinctive landscape of NTS 94-004 is largely a product of underlying bedrock and geological structures, with ornamentation by the Late Wisconsinan Laurentide Ice Sheet.

Topographic and drainage patterns were greatly modified during the phase of maximum ice cover (~18–10 ka BP to ~21 k a BP) and ~21 k a BP (ca. 12 calendar ka BP) with thicknesses in excess of 2.5 m observed in major valleys and is suspected that similar duff thicknesses blanketed bedrock (unit B) across the entire map area.


Degradation began sometime after 18 °C ka BP (or ~21 k a BP) and ended before 10 °C ka BP (ca. 12 calendar ka BP) with the resulting active Laurentide Ice Sheet, glacially overthrust and meltwater superglaciated surfaces d' un territoire d'environ 750 km² couvert par le feuillet cartographique ETANE CREEK (NSR/94-004) dans le nord-est de la Colombie-Britannique.

Post-glaciation (10 °C ka BP, or ca. 12 calendar ka BP to present), changes in regional base-level led to episodes of channel incision and aggradation, resulting in the formation of erosional alluvial terraces along most stream and river valleys in the early Holocene.

Canadian Geoscience Map 108 is an output of the Geo-Mapping for Energy and Minerals Yukon Basin Project managed by Carl Ozyer and Amy Lane (GSC/Canada). The assistance of Robert Cooke, Sean Eagles, Vic Dohar, Mike Sigurdson, Scott Tweedy and Martin Legault (NRCan Scientific Publishing Services) was greatly appreciated throughout the map-making process.

Abstract
Canadian Geoscience Map 108 depicts the surficial geology over some 750 km² covered by the Etane Creek map sheet (NTS 94-004) in northeastern British Columbia. The map area encompasses a significant portion of the Tsoo Tabellands (Alberta Plateau) and is incised by creeks drain south and west into Odyin Creek and Muskwa River, and northeast into the Fort Nelson River. Bedrock is mantled by unconsolidated glacial deposits dating to the Late Pleistocene (Late Wisconsinan Glaciation, ~25 ka to ca. 10 ka) and various glacial Holocene (ca. 10 ka to present). Deposits of till, outwash, and minor moraine deposits, non-coalescent Pleistocene upper (Glaciation du Wisconsinien supérieur, de ~25 ka à 10 ka) ainsi que de dépôts glaciaires Holocène (de 10 ka jusqu'à nos jours). Les dépôts de till, de couleur verte sur la carte, sont généralement propices à l'établissement de l'infrastructure. Les dépôts fluvioglaciaires et éoliens, qui reculent un potentiel en orange et en buff, Slopes disturbed by landslides, debris flows, and rock falls appear brown and pink. Glaciolacustrine and organic deposits with sporadically discontinuous permafrost are coloured purple and grey. Alluvial deposits prone to flooding, erosion, and sedimentation appear yellow on the map.

Résumé
La Carte géoscientifique du Canada 108 illustre la géologie sur une zone de 750 km² couverte par le feuillet cartographique ETANE CREEK (NSR/94-004) dans le nord-est de la Colombie-Britannique. La région cartographique comprend une partie importante des hauts plateaux de Tsoo (plateau de l'Alberta) et est entaillée par des ruisseaux qui coulent vers le sud et vers l'ouest pour se jeter dans le ruisseau Odyin et la rivière Muskwa et vers l'est dans le ruisseau Odyin et la rivière Muskwa et le Fort Nelson. Le socle rocheux est couvert de dépôts glaciaires datant de la Glaciation du Wisconsinien supérieur, de ~25 ka à 10 ka) ainsi que de dépôts glaciaires Holocène (de 10 ka jusqu'à nos jours). Les dépôts de till, de couleur verte sur la carte, sont généralement propices à l'établissement de l'infrastructure. Les dépôts fluvioglaciaires et éoliens, qui reculent un potentiel en orange et en buff, Slopes disturbed by landslides, debris flows, and rock falls appear brown and pink. Glaciolacustrine and organic deposits with sporadically discontinuous permafrost are coloured purple and grey. Alluvial deposits prone to flooding, erosion, and sedimentation appear yellow on the map.



National Topographic System reference and index to adjoining published Geological Survey of Canada maps



Cover illustration: Sandstones and conglomerates exposed along an escarpment of the Tsoo Tabellands in northeast British Columbia, view north. Photograph by D.H. Hurley, 2013-081

Natural Resources Canada / Ressources naturelles du Canada

CANADIAN GEOSCIENCE MAP 108 SURFICIAL GEOLOGY ETANE CREEK British Columbia 1:50 000



Canadian Geoscience Maps



Holocene earth materials and landforms Organic Deposits

Owb Peat bogs: fibric to humic organic matter; massive to stratified accumulations; generally greater than 2 m thick; confined to topographic depressions or level areas; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in wetland areas; logs with sporadically discontinuous permafrost and thermokarst depressions potentially unstable if organic material is disturbed or removed.

O Undifferentiated peat bogs and fens: humic to fibric organic matter, massive to stratified accumulations, generally greater than 2 m thick; confined to topographic depressions, level areas or channels; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in wetland areas; logs with sporadically discontinuous permafrost and thermokarst depressions potentially unstable if disturbed or removed during development.

Af Alluvial fan sediments: boulders, gravel, sand and silt; generally massive to planar stratified; well to rapidly drained; greater than 2 m thick; fan morphology with slopes up to 15°; may contain interbedded debris flows and buried organic material; transported and deposited by modern rivers, streams and creeks; subject to periodic flooding; potential source of aggregate.

Al Alluvial terraced sediments: boulders, gravel, sand and silt; generally massive to planar stratified; well to rapidly drained; greater than 2 m thick; may contain interbedded debris flows and buried organic material; underlain by outwash, till or bedrock; transported and deposited by modern rivers, streams and creeks; subject to rare flooding; potential source of aggregate; land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.

Ap Alluvial floodplain sediments: gravel, sand and silt; massive, trough crossbedded, ripple-bedded, planar stratified; well to rapidly drained; greater than 2 m thick; underlain by till or bedrock; transported and deposited by modern rivers, streams and creeks; subject to seasonal flooding; land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.

Ei Loess: silt and sand; generally massive, well-sorted, crossbedded or ripple-bedded; moderately to well drained; discontinuous veneers less than 1 m thick; blankets greater than 2 m thick; underlain by glacial lake sediments, outwash, till and alluvial deposits; derived from deflation, transportation and deposition of fine-grained sediment by wind action; contains little or no ground ice.

Cv Colluvial deposits: Colluvial veneer: clast-supported diamictons and rubble; massive to stratified, poorly-sorted; well to rapidly drained; deposits less than 2 m thick; landscape headscarp range from 300 m to 10.5 km; formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated debris on slopes above 10-15° with greater than 5 m relief prone to mass-wasting; rock falls, topples, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciolacustrine sediments and outwash containing sporadically discontinuous permafrost; where ground ice is found slope failure can occur on surface less than 5°; slope instability could present major problems for construction in some areas.

Cb Colluvial blanket: clast-supported diamictons and rubble; massive to stratified, poorly-sorted, well to rapidly drained; deposits greater than 2 m thick; landscape headscarp range from 300 m to 10.5 km; formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated debris on slopes above 10-15° with greater than 5 m relief prone to mass-wasting; rock falls, topples, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciolacustrine sediments and outwash containing sporadically discontinuous permafrost; where ground ice is found slope failure can occur on surface less than 5°; slope instability could present major problems for construction in some areas.

GLb Late Pleistocene earth materials and landforms Glaciolacustrine deposits: Glaciolacustrine blanket: silt and clay with subordinate sand, gravel and diamictone; massive or rhythmically interbedded; slump structures and dropstones locally present; poor to moderately drained, generally greater than 2 m thick; kettle lakes and irregular topography underlain by bedrock, tills and outwash; transported by and deposited from sediment-laden meltwater, subsaqueous gravelly flows and thermal melting of ice in proglacial lakes; where sporadically discontinuous permafrost is, or was present, glaciolacustrine sediments may be subject to thermokarst processes; slopes less than 5° are potentially unstable and prone to landslides and debris flows.

GfH Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamictone; generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick; irregular hummocks and kettled topography; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by rivers and streams flowing from, or in contact with, glacial ice; potential source of groundwater and gravel aggregate when material is gravel rich.

Tm Moraine ridges: sand, silt and clay-rich diamictons, massive, matrix-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 m high; major moraines up to 12 km in length and 10 m high; ridge shape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional fill moraines; major ridges feature are large recessional end moraines and ice-thrust ridges, generally suitable for infrastructure placement.

Ts Streamlined till: silt and clay-rich diamictons, massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 5 km in length; generally less than 50 m wide and 20 m high; formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement.

Tv Till veneer: sand, silt and clay-rich diamictons, massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; less than 2 m thick draping bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation beneath active, warm-based ice and in situ melting from stagnant cold-based ice; generally suitable for infrastructure placement.

R Pre-Quaternary earth materials and landforms Bedrock: Undifferentiated bedrock: conglomerates, sandstone, siltstone, shale and limestone; exposed in escarpments between 300 m and 80 km in length; slopes above 10-15° with greater than 5 m relief prone to rock falls, topples, rock slides and debris flows; Paleozoic; unconformably overlain by Mesozoic sedimentary rocks; limestone and classic sedimentary rocks are a potential source of crushed granular aggregate.

- Geological boundary (Confidence: approximate)
Bedrock scarp
Major moraine ridge (unspecified)
Other moraine ridge (unspecified)
Drumlin ridge
Major meltwater channel scarp
Minor meltwater channel central axis (unspecified, sense: known)
Terrace scarp (environment: glacioluvial)
Terrace scarp (environment: fluvial)
Terrace scarp (environment: glaciolacustrine)
Station location (ground observation)

Recommended citation: Hurley, D.H., Hicks, A.S., Chow, W., and Mirmohammadi, M. 2013. Surficial geology, Etane Creek, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 108 (preliminary), scale 1:50 000. doi:10.4095/292000

Preliminary publications in this series have not been scientifically edited.

Authors: D.H. Hurley, A.S. Hicks, W. Chow, and M. Mirmohammadi
Geomatics by D.H. Hurley, W. Chow, and M. Mirmohammadi
Cartography by W. Chow
Initiative of the Geological Survey of Canada, conducted under the auspices of the Yukon Basin Project as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program

SURFICIAL GEOLOGY ETANE CREEK British Columbia 1:50 000



The Geological Survey of Canada welcomes corrections or additional information from users. This publication is available for free download through GEOCAN (http://geocan.nrc.ca/gc/can/).

Map projection Universal Transverse Mercator, zone 10, North American Datum 1983. Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations in feet above mean sea level. Magnetic declination 2013, 20'28"E, decreasing 21" annually.

Recommended citation: Hurley, D.H., Hicks, A.S., Chow, W., and Mirmohammadi, M. 2013. Surficial geology, Etane Creek, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 108 (preliminary), scale 1:50 000. doi:10.4095/292000

CANADIAN GEOSCIENCE MAP 108 SURFICIAL GEOLOGY ETANE CREEK British Columbia