

INTRODUCTION

The Surficial Geology Map of NTS 94-O16 (Canadian Geoscience Map 118) is the product of collaboration between the Geological Survey of Canada and the British Columbia Ministry of Energy, Mines and Natural Gas as part of the Geoscience for Energy and Minerals Program (GEM) Energy Yukon Basin Project.

Map preparation and field-based observations have led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-O16 map area (Hurley and Hickin, 2010; Hurley et al., 2011a,b).

The surficial geology map area is largely a product of underlying bedrock and tectonic structures, with ornamentation by the Late Pleistocene Laurentide Ice Sheet. The map area encompasses a portion of the Foothills Province and the northernmost part of the Canadian Shield.

Map preparation and field-based observations have led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-O16 map area (Hurley and Hickin, 2010; Hurley et al., 2011a,b).

Map preparation and field-based observations have led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-O16 map area (Hurley and Hickin, 2010; Hurley et al., 2011a,b).

ACKNOWLEDGMENTS

Canadian Geoscience Map 118 is an outcrop of the Geo-Mapping for Energy and Minerals Yukon Basin Project managed by Carl Cyren and Larry Lane (GS/Cyren).

Abstract

Canadian Geoscience Map 118 depicts the surficial geology over some 750 km² covered by the Stanislas Creek map area in northwestern British Columbia. The map area is drained by the Fortune and Stanislas creeks and their tributaries which flow north into the northwest-draining Pełlet River. Bedrock is mantled by unconsolidated earth materials dating to the late Pleistocene-Late Wisconsinian Glaciation, > 25 ka to ca. 10 ka and non-glacial Holocene (< 10 ka to present).


Deposits of till green on the map, are generally suitable for placement of infrastructure. Glaciofluvial deposits with mineral, aggregate, and groundwater potential are coloured orange. Slopes disturbed by landslides, debris flows, and rock falls represent high-risk areas. Glaciolacustrine and organic deposits with sporadically discontinuous permafrost are coloured purple and grey. Alluvial deposits prone to flooding, erosion, and sedimentation appear yellow on the map.

Catalogue No. M183-1118-2013E-PDF ISBN 978-1-103-21798-5 doi:10.4095/29285



CANADIAN GEOSCIENCE MAP 118 SURFICIAL GEOLOGY STANISLAS CREEK

British Columbia 1:50 000



Holocene earth materials and landforms

- Owb: Peat bogs: fibric to humic organic matter, massive to stratified accumulations; generally greater than 2 m thick, confined to topographic depressions or level areas... Owf: Fens: fibric organic matter, massive to stratified, generally greater than 2 m thick, confined to topographic depressions, level areas or channels... O: Undifferentiated peat bogs and fens: humic to fibric organic matter, massive to stratified accumulations, generally greater than 2 m thick... Af: Alluvial fan sediments: boulders, gravel, sand and silt; generally massive to planar stratified, well to rapidly drained... Ai: Alluvial terrace deposits: boulders, gravel, sand and silt; generally massive to planar stratified, well to rapidly drained... Ap: Alluvial floodplain deposits: gravel, sand and silt; massive, trough cross-bedded, rippled-bedded, planar stratified... Cv: Colluvial veneer: clast-supported diamictites and rubble; massive to stratified, poorly-sorted... Gb: Glaciolacustrine deposit: massive or rhythmically interbedded; slump structures and irregularly locally present... GFb: Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamict; generally massive to stratified... GFc: Esker ridges: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamictite... GFf: Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamictite... Th: Till blanket: sand, silt and clay-rich diamictites; massive, matrix-supported and compact... Tm: Moraine ridges: sand, silt and clay-rich diamictites; massive, matrix-supported... Ts: Streamlined till: silt and clay-rich diamictites; massive, matrix-supported and compact...

Late Pleistocene earth materials and landforms

- Gfb: Glaciolacustrine deposit: massive or rhythmically interbedded; slump structures and irregularly locally present... GFb: Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamict; generally massive to stratified... GFc: Esker ridges: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamictite... GFf: Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamictite... Th: Till blanket: sand, silt and clay-rich diamictites; massive, matrix-supported and compact... Tm: Moraine ridges: sand, silt and clay-rich diamictites; massive, matrix-supported... Ts: Streamlined till: silt and clay-rich diamictites; massive, matrix-supported and compact...

- Geological boundary (Confidence: approximate) Major moraine ridge (end, interlobate, or unspecified) Other moraine ridge (DeGeer, minor lateral, recessional, rogen, washboard, other transverse or unspecified) Esker ridge (sense: unknown or unspecified) Drumlin ridge Major meltwater channel scarp Minor meltwater channel central axis (marginal, overflow, subglacial or unspecified; sense: known) Terrace scarp (environment: glaciofluvial) Terrace scarp (environment: fluvial) Station location (ground observation or stratigraphic section)

Recommended citation Hurley, D.H., Hickin, A.S., Chow, W., and Mirzaei, M., 2013. Surficial geology, Stanislas Creek, British Columbia: Geological Survey of Canada, Canadian Geoscience Map 118 (preliminary), scale 1:50 000. doi:10.4095/29285

Preliminary publications in this series have not been scientifically edited.