CANADIAN GEOSCIENCE MAP 121 Holocene earth materials and landforms DESCRIPTIVE NOTES Peat bogs: fibric to humic organic matter; massive to stratified accumulations; INTRODUCTION generally greater than 2 m thick; confined to topographic depressions or level This Surficial Geology Map of NTS 94-O/13 (Canadian Geoscience Map 121) is the product of collaboration between the Geological Survey of Canada and the British Columbia Ministry of Energy, Mines and Natural Gas as part of the Geo-mapping for Energy and Minerals Program areas; underlain by poorly drained till, glaciolacustrine and other GEM-Energy Yukon Basins Project. The accompanying geodatabase includes field observation points and field photos, landform features as unconsolidated sediments: formed by decomposition of plant material in lines, and surficial geology unit polygons. The map and geodatabase are essential baseline geoscience information for a range of potential wetland areas; bogs with sporadically discontinuous permafrost and end-users including resource explorationists, geotechnical engineers, land-use managers, terrestrial ecologists, archaeologists thermokarst depressions potentially unstable if organic material is disturbed or geoscientists and communities in northern British Columbia. By providing new insight into the distribution and origins of surficial earth materials, CGM 121 will help to reduce the economic costs and risks associated with the sustainable development of energy and mineral resources in NTS 94-0/13. Environmental impact assessments for new access roads, work camps, well pads, pipeline and power **Fens**: fibric organic matter; massive to stratified; generally greater than 2 m transmission line corridors, water storage and waste management systems and other infrastructure will benefit from the geoscience nformation presented here. By identifying areas prone to geological hazards (e.g., landslides, permafrost, flooding), CGM 121 will also help to thick; confined to topographic depressions, level areas and meltwater $protect \, natural \, resources, in frastructure \, and \, communities \, vulnerable \, to \, climate \, change \, in \, Canada's \, north.$ channels; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in APPROACH TO SURFICIAL GEOLOGY MAPPING wetland areas; fens are prone to flooding following drainage damming by Terrain mapping and field-based benchmarking studies have led to a better understanding of the regional distribution of surficial deposits permafrost, landslides and other geomorphic processes in the NTS 94-O/13 map area (Huntley and Hickin, 2010; Huntley et al., 2011a-b). Surficial earth materials and landforms were interpreted using a combination of stereo-pair air photos (BCB97010, 15BCB97015 Undifferentiated peat bogs and fens: humic to fibric organic matter; massive 15BCB97029, 15BCB97075 and 15BCB97088 series), LANDSAT 7 satellite imagery (http://glovis.usgs.gov/ [URL 2011]) and Shuttle Radar to stratified accumulations; generally greater than 2 m thick; confined to Topography Mission digital elevation models (http://dds.cr.usgs.gov/srtm/ [URL 2011]). The base map was generated from CANVEC shape topographic depressions, level areas or channels; underlain by poorly drained files (http://geogratis.cgdi.gc.ca/geogratis/ [URL 2011]). Surficial geology polygons and landform line symbols were digitized using till, glaciolacustrine and other unconsolidated sediments; formed by commercially available computer software packages (Global Mapper, ArcMap and ArcGIS) and compared to published maps, reports and decomposition of plant material in wetland areas; may contain sporadically archived digital data (e.g., Stott and Taylor, 1968; Bednarski, 2003a-d; Clement et al., 2004; Bednarski, 2005a-b; Trommelen and Levson, discontinuous permafrost and thermokarst depressions; potentially unstable if 2008; Demchuk, 2010). The geodatabase accompanying this map conforms to the Science Language for the Data Management component disturbed or removed during development. of the GEM Geological Map Flow process (cf. Huntley and Sidwell, 2010; Huntley et al., 2011a; Deblonde et al., 2012). Fieldwork was undertaken in 2009 and 2010 to ground truth surficial geology polygons interpreted from air photos and satellite imagery, and to Alluvial fan sediments: boulders, gravel, sand and silt; generally massive to gather characteristics that could not be determined through remote predictive mapping. Earth materials were defined on the basis of facies and landform associations, texture, sorting, colour, sedimentary structures, degree of consolidation, and stratigraphic contact relationships field stations and remote observations from helicopters. The distribution of glacial and non-glacial landforms is depicted on the surficial geology map. Map units in the Legend are presented chronostratigraphically and include organic deposits, alluvial, colluvial, eolian, glaciolacustrine and glaciofluvial sediments, tills and areas of bedrock. INFERRED GEOLOGICAL HISTORY The distinctive landscape of NTS 94-O/13 is largely a product of underlying bedrock and geological structures, with ornamentation by the Late massive to planar stratified; well to rapidly drained; greater than 2 m thick; may Wisconsinan Laurentide Ice Sheet. In the northeast, the plateau is underlain by conglomerate, sandstone, carbonaceous shale and coal contain interbedded debris flows and buried organic material; underlain by (Upper Cretaceous Wapiti Formation). Over much map area, shale, siltstone and sandstone (Upper Cretaceous Kotaneelee Formation) lie outwash, till or bedrock; transported and deposited by modern rivers, streams beneath surficial units. Undifferentiated clastic bedrock (Lower Cretaceous Fort St. John Group) is exposed along hill crests in the northwest Topography and drainage patterns were greatly modified during the phase of maximum ice cover (>1814 C ka BP or >21.4 calendar ka BP). and wildlife resources. Unconsolidated sediment thicknesses in excess of 2-5 m are observed in major valleys and it is suspected that similar drift thicknesses blanket bedrock (R) across the map area. Silt- and clay-rich Laurentide tills have low clast contents (<20%) of proximally derived Cretaceous siliciclastic sedimentary rocks and distal igneous and metamorphic clasts from the Canadian Shield, hundreds of kilometres to the northeast Drumlin ridges up to several hundred metres in length suggest clay-rich tills (unit Ts) were deposited beneath active, rapidly flowing warm-based glacial ice (Huntley and Hickin, 2010; Huntley et al., 2011b). Drumlinized till (unit Ts) is most pronounced south of the Petitot River and west of the Maxhamish Escarpment where ice flowed up-hill, and thick accumulations of till were deposited over soft bedrock and unconsolidated advance-phase sediments. Many lake basins were excavated by erosion and ice-thrusting as Laurentide ice and subglacial meltwater scoured and deformed older glacial deposits and weak bedrock. Deglaciation began sometime after 18 ¹⁴C ka BP (or >21.4 calendar ka BP) and ended before 10 ¹⁴C ka BP (ca. 12 calendar ka BP), with the Colluvial deposits retreating active Laurentide Ice Sheet, stagnant ice masses in lowlands, glaciofluvial outwash and landslide debris blocking and reordering Colluvial veneer: clast-supported diamictons and rubble; massive to stratified, regional drainage. The mapped distribution of moraine ridges (unit Tm) implies that ice margins receded to the east across the map sheet poorly-sorted; well to rapidly drained; deposits less than 2 m thick; landslide (Huntley and Hickin, 2010). Some large end moraines are deformed and streamlined suggesting that receding lobes remained active during retreat and occasionally rapidly advanced. Minor moraine ridges drape drumlins in cross-cutting patterns and are interpreted as crevasse headscarps range from 300 m to 10.5 km; formed by the weathering and illings and squeeze moraines deposited shortly after drumlinization ended, or as ice retreated from the map area (Huntley et al., 2011b). down-slope movement of earth materials by gravitational processes; bedrock Hummocky till (unit Th) found with short segments of subareal-subglacial meltwater channels and eskers indicate that bodies of stagnant and unconsolidated debris on slopes above 10-15° with greater than 5 m relief glacier ice remained in lowland areas west of the Maxhamish Escarpment (Huntley et al., 2011a-b). As ice retreated from the map area, prone to mass-wasting; rock falls, topples, rock slides and debris flows occur proglacial lakes formed on the Liard Plateau and were linked by spillways that drained meltwater northward into the Mackenzie River basin. In where shale, sandstone and carbonate strata is exposed close to the surface: the map area, glaciolacustrine deposits (unit GLb), glaciofluvial terraces (unit GFt), and meltwater channels incised into till and bedrock indicate that glacial lake levels fell stepwise through deglaciation, with stable elevations at approximately 420 m, 380 m and <300 m. Fineretrogressive rotational debris slides, debris flows and slumps occur in grained glacial earth materials have been re-worked by eolian activity and discontinuous loess covers glacial lake and till deposits in some glaciolacustrine sediments and outwash containing sporadically discontinuous Post-glaciation (10 [™]C ka BP, or ca. 12 calendar ka BP to present), changes in regional base-level led to episodes of channel incision and aggradation, resulting in the formation of erosional alluvial terraces along most stream and river valleys. In the early Holocene, pulses of fluvial terrace building followed initial valley incision by the Liard and other major rivers. Most streams and rivers have alluvial fans (unit Af) and Late Pleistocene earth materials and landforms terraces (unit Åt) <5 m above active floodplains (unit Ap) consisting of gravel overlain by silt and sand. Poorly drained clay-rich till on the plateaux and glaciolacustrine sediments in lowland areas are covered by extensive postglacial peat deposits (unit Owb), fens (unit Owf) and Glaciolacustrine deposits undifferentiated wetlands (unit O). Discontinuous permafrost is sporadically encountered in glaciolacustrine and some peat deposits. Charcoal, observed in dug pits on alluvial terraces, suggest forest fires may have contributed to periods of landslide activity on slopes and local fluvial aggradation. Landslides and colluviated deposits (unit Cv) are common where bedrock outcrops form escarpments, and where shale or ine-grained glacial deposits are exposed along steep cutbanks. Stream networks and wetlands draining plateau watersheds are disrupted by peaver activity and, to a lesser extent, by roads and infrastructure where they cross streams, rivers and organic deposits (Huntley and Hickin, 2010; Huntley and Hickin, 2011a-b). outwash; transported by and deposited from sediment-laden meltwater, subaqueous gravity flows and thermal melting of ice in proglacial lakes, where sporadically discontinuous permafrost is, or was present, glaciolacustrine Canadian Geoscience Map 121 is an output of the Geo-Mapping for Energy and Minerals Yukon Basins Project managed by Carl Ozyer and Larry Lane (GSC-Calgary). The assistance of Robert Cocking, Sean Eagles, Vic Dohar, Mike Sigouin, Scott Tweedy and Martin Legault sediments may be subject to thermokarst processes; slopes less than 5° are potentially unstable and prone to landslides and debris flows. (NRCAN Scientific Publishing Services) was greatly appreciated throughout the map-making process. A critical review of CGM 121 was Glaciofluvial deposits provided by Ron DiLabio (GSC-Ottawa). Canadian Geoscience Map 121 depicts the surficial La Carte géoscientifique du Canada 121 illustre la geology over some 790 km² covered by the Sandy géologie des matériaux superficiels d'un territoire Creek map sheet (NTS 94-O/13) in northeastern British d'environ 790 km² couvert par le feuillet cartographique Columbia. The map area lies at the western limit of the de Sandy Creek (SNRC 94-O/13), dans le nord-est de Liard Plateau and is incised by the Liard River and its la Colombie-Britannique. La région cartographique se tributaries. Bedrock is mantled by unconsolidated earth situe à la limite occidentale du plateau de Liard et est materials dating to the Late Pleistocene (Late entaillée par la rivière Liard et ses affluents. Le socle Wisconsinan Glaciation, > 25 ka to ca. 10 ka) and nonrocheux est couvert de matériaux terrestres non consolidés remontant au Pléistocène supérieur glacial Holocene (ca. 10 ka to present). Deposits of till, green on the map, are generally suitable for placement (Glaciation du Wisconsinien supérieur, de > 25 ka à of infrastructure. Glaciofluvial deposits with mineral. env. 10 ka) ainsi que de matériaux non glaciaires de aggregate, and groundwater potential are coloured l'Holocène (d'env. 10 ka jusqu'à nos jours). Les dépôts orange. Slopes disturbed by landslides, debris flows, de till, de couleur verte sur la carte, sont généralement and rock falls appear brown and pink. Glaciolacustrine and organic deposits with sporadically discontinuous dépôts fluvioglaciaires, qui recèlent un potentiel en Hummocky till: sand and silt-rich diamictons; massive to stratified, matrixpermafrost are coloured purple and grey. Alluvial minéraux, en agrégats et en eau souterraine, sont and clast-supported; clast contents less than 20% and contain sub-rounded figurés par la couleur orange. Les versants dérangés deposits prone to flooding, erosion, and sedimentation par des glissements de terrain, des coulés de débris et appear yellow on the map. des chutes de blocs sont représentés en brun et en rose. Les dépôts glaciolacustres et organiques, qui renferment sporadiquement du pergélisol discontinu, sont représentés en violet et en gris. Les dépôts alluviaux sujets aux inondations, à l'érosion et à la sédimentation apparaissent en jaune sur la carte. infrastructure placement. CGM 121 | CGM 12 CGM 122 | CGM 125 National Topographic System reference and index to adjoining published Geological Survey of Canada maps **Cover illustration** Catalogue No. M183-1/121-2013E-PDF View northwest of moraine ridges, drumlins, ISBN 978-1-100-21792-5 infrastructure placement. neltwater channels and organic deposits over the doi:10.4095/292398 iard Plateau, northeast British Columbia. ---- Geological boundary (Confidence: approximate) hotograph by D.H. Huntley. 2013-099 © Her Majesty the Queen in Right of Canada 2013 Bedrock scarp other transverse or unspecified) Natural Resources Ressources naturelles du Canada ----- Landslide escarpment (Status: inactive or unspecified) **CANADIAN GEOSCIENCE MAP 121** Major meltwater channel scarp **SURFICIAL GEOLOGY** Terrace scarp (environment: glaciofluvial) TTTTTTTTT Terrace scarp (environment: fluvial) **SANDY CREEK** Terrace scarp (environment: glaciolacustrine) British Columbia 1:50 000

REFERENCES

Bednarski, J.M., 2003a. Betalamea Lake, Northwest Territories - Yukon Territory - British Columbia (NTS 95B/4); Geological Survey of Canada, Open File 4502, scale 1:50 000. Bednarski, J.M., 2003b. Surficial geology of Fort Liard, Northwest Territories - British Columbia. Geological Survey of Canada, Open File 1760, scale 1:50 000.

Bednarski, J.M., 2003c. Surficial geology of Lake Bovie, Northwest Territories - British Columbia. Geological Survey of Canada, Open File 1761, scale 1:50 000. Bednarski, J.M., 2003d. Surficial geology of Celibeta Lake, Northwest Territories - British Columbia. Geological Survey of Canada, Open File 1754, scale 1:50,000.

Bednarski, J.M., 2005a. Surficial Geology of Etsine Creek, British Columbia, Geological Survey of Canada, Open

Bednarski, J.M., 2005b. Surficial Geology of Gote Creek, British Columbia, Geological Survey of Canada, Open Clement, C., Kowall, R. Huntley, D. and Dalziel, R., 2004. Ecosystem units of the Sahtaneh area; Slocan Forest roducts (Fort Nelson) Report, 39 pages and appendices.

Deblonde, C., Plouffe, A., Boisvert, E., Buller, G., Davenport, P., Everett, D., Huntley, D., Inglis, E., Kerr, D., Moore, A., Paradis, S.J., Parent, M., Smith, R., St-Onge, D., and Weatherston, A., 2012.

Trommelen, M. and Levson, V.M., 2008. Quaternary stratigraphy of the Prophet River, norheastern British

Columbia; Canadian Journal of Earth Sciences, Vol. 45, pages 565-575.

Science Language for an Integrated Geological Survey of Canada Data Model for Surficial Maps Version 1.1 Results of Geological Survey of Canada Surficial Legend Review Committee; Geological Survey of Canada, Demchuk, T., 2010. Surficial geology of the Komie Creek area (NTS 094P/05). British Columbia Ministry of

planar stratified; well to rapidly drained; greater than 2 m thick; fan morphology Energy, Mines and Petroleum Resources, Open File 2010-08; Geological Survey of Canada Open File 6568, with slopes up to 15o; may contain interbedded debris flows and buried scale 1:50 000. organic material; transported and deposited by modern rivers, streams and Huntley, D.H. and Hickin, A.S., 2010. Surficial deposits, landforms, glacial history and potential for granular aggregate and frac sand: Maxhamish Lake Map Area (NTS 94-O), British Columbia. Geological Survey of creeks; subject to periodic flooding; potential source of aggregate. Alluvial terraced sediments: boulders, gravel, sand and silt; generally Canada, Open File 6430, 17 pages. Huntley, D., Hickin, A. and Chow, W., 2011a. Surficial geology, geomorphology, granular resource evaluation and

geohazard assessment for the Maxhamish Lake map area (NTS 94-O), northeastern British Columbia; Geological Survey of Canada, Open File 6883, 20 pages. and creeks; subject to rare flooding; potential source of aggregate; land use Huntley, D.H., Hickin, A.S. and Ferri, F., 2011b. Provisional surficial geology, glacial history and paleogeographic activities may adversely affect stream courses and conditions, and impact fish reconstructions of the Toad River (NTS 94-N) and Maxhamish Lake map areas (NTS 94-O), British Columbia. Geoscience Reports 2011, BC Ministry of Energy, pages 37-55. Alluvial floodplain sediments: gravel, sand and silt; massive, trough $Huntley, D.H.\ and\ Sidwell,\ C.F.,\ 2010.\ Application\ of\ the\ GEM\ surficial\ geology\ data\ model\ to\ resource\ evaluation$ cross-bedded, rippled-bedded, planar stratified; well to rapidly drained; greater

and geohazard assessment for the Maxhamish Lake map area (NTS 94-O), British Columbia. Geological than 2 m thick; underlain by till or bedrock; transported and deposited by Survey of Canada, Open File 6553, 22 pages. modern rivers, streams and creeks; subject to seasonal flooding; land use activities may adversely affect stream courses and conditions, and impact fish Stott, D.F. and Taylor, G.C., 1968. Geology of Maxhamish Lake. Geological Survey of Canada, Map 2-1968,

Late Pleistocene to Holocene earth materials and landforms

permafrost; where ground ice is found slope failure can occur on surfaces less than 5°; slope instability could present major problems for construction in Glaciolacustrine blanket: silt and clay with subordinate sand, gravel and diamicton; massive or rhythmically interbedded; slump structures and ropstones locally present; poor to moderately drained; generally greater than 2 m thick; kettle lakes and irregular topography underlain by bedrock, tills and

Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamicton; generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick; terrace scarps range from 100 m to 8 km in length; in contact with, and overlying other till units, outwash and glaciolacustrine sediments; deposited by meltwater confined to proglacial channels and spillways; potential source of groundwater and granular aggregate when material is gravel rich.

Till blanket: sand, silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glacigenic deformation of sediment beneath active, warm-based ice and in situ melting from stagnant cold-based ice; stable terrain, generally suitable for infrastructure placement.

granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; drapes till and other glacial deposits; deposited by in situ melting from stagnant cold-based ice and modified by meltwater: evidence for ice collapse includes slump structures, kettle lakes and irregular topography; potential source of aggregate when material is gravel rich; generally suitable for infrastructure placement. Moraine ridges: sand, silt and clay-rich diamictons; massive,

matrix-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 m high; major moraines up to 12.5 km in length and 10 m high; ridges drape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional push moraines; major ridges features are large recessional end moraines and ice-thrust ridges; generally suitable for

Streamlined till: silt and clay-rich diamictons: massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield: moderately well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 9 km in length; generally less than 50 m wide and 20 m high; formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glacigenic deformation of sediment beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement.

Till veneer: sand, silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; less than 2 m thick draping bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glacigenic deformation beneath active, warm-based ice and in situ melting from stagnant cold-based ice; generally suitable for

• • • • • • Major moraine ridge (end, interlobate, or unspecified) Other moraine ridge (DeGeer, minor lateral, recessional, rogen, washboard,

Preliminary

Station location (ground observation or stratigraphic section)

Recommended citation Huntley, D.H., Hickin, A.S., Chow, W., and Mirmohammadi, M., 2013. Surficial geology, Sandy Creek, British Columbia; Geological Survey of Canada, Canadian Geoscience Map 121 (preliminary), scale 1:50 000. doi:10.4095/292398

Preliminary Canadian M. Mirmohammadi

Authors: D.H. Huntley, A.S. Hickin, W. Chow, and Geology by D.H. Huntley and A.S. Hickin (2009–2010) Geological compilation by D.H. Huntley (2009–2011)

Preliminary Geomatics by D.H. Huntley, W. Chow, and Cartography by W. Chow Initiative of the Geological Survey of Canada, conducted under the auspices of the Yukon Basin

Project as part of Natural Resources Canada's Geo

mapping for Energy and Minerals (GEM) program

SURFICIAL GEOLOGY **SANDY CREEK British Columbia**

CANADIAN GEOSCIENCE MAP 121

Map projection Universal Transverse Mercator, zone 10. North American Datum 1983 Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations in feet above mean sea level Magnetic declination 2013, 20°45′E

decreasing 22' annually.

Preliminary

The Geological Survey of Canada welcomes corrections or additional information from users. his publication is available for free download through GEOSCAN (http://geoscan.ess.nrcan.gc.ca/).

Preliminary

70 4**71**000m. E.

Preliminary

reliminary publications li this series have not been scientifically edited.

CANADIAN GEOSCIENCE MAP 121 SURFICIAL GEOLOGY SANDY CREEK British Columbia

Canada

Geoscience Maps