

DESCRIPTIVE NOTES

INTRODUCTION
This Surficial Geology Map of NTS 94-O19 (Canadian Geoscience Map 123) is the product of collaboration between the Geological Survey of Canada and the British Columbia Ministry of Energy, Mines and Natural Gas as part of the Geoscience for Energy and Minerals Program (GEM-Energy Yukon Basin Project).

APPROACH TO SURFICIAL GEOLOGY MAPPING
This map and field-based benchmarks have led to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-O19 map area (Huntley and Hickin, 2010; Huntley et al., 2011a,b).

FIELDWORK WAS UNDERTAKEN IN 2009 AND 2010 TO GROUND TRUTH SURFICIAL GEOLOGY POLYGONS INTERPRETED FROM AIR PHOTOS AND SATELLITE IMAGERY, AND TO GATHER ADDITIONAL DATA THAT COULD NOT BE OBTAINED THROUGH REMOTE SENSING.

INFERRED GEOLOGICAL HISTORY
The distinctive landscape of NTS 94-O19 largely a product of underlying bedrock and geological structures, with ornamentation by the Upper Wisconsinan Laurentide Ice Sheet.

TOPOGRAPHY AND DRAINAGE PATTERNS WERE GREATLY MODIFIED DURING THE PHASE OF MAXIMUM ICE COVER (>18 °C ka BP or >21 A calendar ka BP). LANDSLIDES AND OTHER GEOMORPHIC PROCESSES IN THE NTS 94-O19 MAP AREA (HUNTLEY AND HICKIN, 2010; HUNTLEY ET AL., 2011A,B).

DEGLACIATION BEGAN SOMETIME AFTER 18 °C ka BP (>21 A calendar ka BP) AND ENDED BEFORE 10 °C ka BP (ca. 12 calendar ka BP), WITH THE RETREATING ACTIVE LAURENTIDE ICE SHEET, STAGNANT ICE MASSES IN TOWARDS, GLACIOFLUVIAL OUTWASH AND INSTABLE DEBRIS BLOCKING AND REORDERING TERRAIN.

POST-GLACIATION (10 °C ka BP, OR EARLIER, 12 CALENDAR KA BP) TO PRESENT, CHANGES IN REGIONAL BASE-LEVEL LED TO EPISODES OF CHANNEL INCISION AND AGGRAVATION, RESULTING IN THE FORMATION OF OROSSAL ALLUVIAL TERRACES ALONG MOST STRAIGHT AND NEARLY VERTICAL, PULSES OF FLUVIAL TERROR BUILDING FOLLOWED BY RELATIVE VELOCITY INCREASE.

ACKNOWLEDGMENTS
Canadian Geoscience Map 123 is an output of the Geo-Mapping for Energy and Minerals Yukon Basin Project managed by Carl Ozyer and Amy Law (GSC-Calgary).

Abstract
Canadian Geoscience Map 123 depicts the surficial geology over some 700 km² covered by the Trail Lake map sheet (NTS 94-O19), in northeastern British Columbia. Here, the Eshbo Plateau lies within the Pettit River watershed and is drained by east-flowing Gole Creek, and northward by Dilly and Yesashelle creeks.

Résumé
La Carte géoscientifique du Canada 123 illustre la géologie des matériaux superficiels d'un territoire d'environ 700 km² couvert par le feuillet cartographique de Trail Lake (SMRC 94-109), dans le nord-est de la Colombie-Britannique.

Table with 2 columns: CGM 119, CGM 118, CGM 126, CGM 123, CGM 127, CGM 124. Includes National Topographic System reference and index to adjoining published Geological Survey of Canada maps.



Cover Illustration
Rotational landslide triggered by the Kwigiana River incising the Eshbo Plateau in northeast British Columbia, view west. Photograph by D.H. Huntley, 2013-101.

Natural Resources Canada Ressources naturelles du Canada

CANADIAN GEOSCIENCE MAP 123 SURFICIAL GEOLOGY TRAIL LAKE BRITISH COLUMBIA 1:50 000



Canadian Geoscience Maps



Holocene earth materials and landforms Organic Deposits

- Owb Peat bogs: fibric to humic organic matter; massive to stratified accumulations; generally greater than 2 m thick, confined to topographic depressions or level areas, underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in wetland areas; bogs with sporadically discontinuous permafrost and thermokarst depressions potentially unstable if organic material is disturbed or removed.
Owf Fens: fibric organic matter; massive to stratified; generally greater than 2 m thick, confined to topographic depressions, level areas and meltwater channels; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in wetland areas; fens are prone to flooding following drainage damming by beaver activity.
O Undifferentiated peat bogs and fens: humic to fibric organic matter; massive to stratified accumulations; generally greater than 2 m thick, confined to topographic depressions, level areas and meltwater channels; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in wetland areas; may contain sporadically discontinuous permafrost and thermokarst depressions; potentially unstable if disturbed or removed during development.
At Alluvial terraced deposits: boulders, gravel, sand and silt; generally massive to planar stratified; well to rapidly drained; greater than 2 m thick; may contain interbedded debris flows and buried organic material; underlain by outwash, till or bedrock; transported and deposited by modern rivers, streams and creeks; subject to rare flooding; potential source of aggregate; land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.
Ap Alluvial floodplain sediments: gravel, sand and silt; massive, trough cross-bedded, topsoiled, planar stratified; well to rapidly drained; greater than 2 m thick; underlain by till or bedrock; transported and deposited by modern rivers, streams and creeks; subject to seasonal flooding; land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.
Cv Colluvial deposits
Cv1 Colluvial veneer: clast-supported diamictons and rubble; massive to stratified, poorly-sorted; well to rapidly drained; deposits less than 2 m thick; landslide headscarp ranges from 300 m to 10 to 15 km; formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated till on slopes above 10-15° with greater than 5 m relief prone to mass-wasting; rock falls, topgles, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciolacustrine sediments and outwash containing sporadically discontinuous permafrost; where ground ice is found slope failure can occur on surfaces less than 5°; slope instability could present major problems for construction in some areas.
Cb Colluvial blanket: clast-supported diamictons and rubble; massive to stratified, poorly-sorted; well to rapidly drained; deposits greater than 2 m thick; landslide headscarp ranges from 200 m to 10 to 15 km; formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated till on slopes above 10-15° with greater than 5 m relief prone to mass-wasting; rock falls, topgles, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciolacustrine sediments and outwash containing sporadically discontinuous permafrost; where ground ice is found slope failure can occur on surfaces less than 5°; slope instability could present major problems for construction in some areas.
OLB Glaciolacustrine blanket: silt and clay with subordinate sand; gravel and diamicton; massive or rhythmically interbedded; slump structures and drapenets locally present; poor to moderately drained; generally greater than 2 m thick; kettle, irregular topography underlain by bedrock, till and outwash; transported by and deposited from sediment laden meltwater; subaqueous gravity flows and thermal melting of ice in proglacial lakes; were sporadically discontinuous permafrost; as is present, glaciolacustrine sediments may be subject to thermokarst processes; slopes less than 5° are potentially unstable and prone to landslides and debris flows.
GFH Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; irregular hummocks and kettled topography; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by rivers and streams flowing from, or in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.
GFR Esker terraces: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; range from 100 m to 8 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by subglacial meltwater in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.
GFI Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; terrace scarp ranges from 100 m to 8 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by meltwater confined to proglacial channels and spillways; potential source of groundwater and granular aggregate when material is gravel rich.
Tb Till deposits
Tb1 Till blanket: sand, silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath active, warm-based ice and in situ melting from stagnant cold-based ice; stable terrain, generally suitable for infrastructure placement.
Th Hummocky till: sand and silt-rich diamictons; massive to stratified, matrix- and clast-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; drapes till and other glacial deposits; deposited by in situ melting from stagnant cold-based ice and modified by meltwater; evidence for ice collapse includes slump structures, kettle lakes and irregular topography; potential source of aggregate when material is gravel rich; generally suitable for infrastructure placement.
Tm Moraine ridges: sand, silt and clay-rich diamictons; massive, matrix-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 m high, major moraines up to 12.5 km in length and 10 m high; ridges drape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional push moraines; major ridges features are large recessional end moraines and ice-thrust ridges; generally suitable for infrastructure placement.
Ts Streamlined till: silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 8 km in length; generally less than 50 m wide and 20 m high, formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement.

Late Pleistocene earth materials and landforms Glaciolacustrine deposits

- GFH Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; irregular hummocks and kettled topography; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by rivers and streams flowing from, or in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.
GFR Esker terraces: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; terrace scarp ranges from 100 m to 8 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by subglacial meltwater in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.
GFI Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified; some slump structures; moderately to well-drained; greater than 2 m thick; terrace scarp ranges from 100 m to 8 km in length; in contact with, and overlying till units, outwash and glaciolacustrine sediments; deposited by meltwater confined to proglacial channels and spillways; potential source of groundwater and granular aggregate when material is gravel rich.
Tb Till deposits
Tb1 Till blanket: sand, silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath active, warm-based ice and in situ melting from stagnant cold-based ice; stable terrain, generally suitable for infrastructure placement.
Th Hummocky till: sand and silt-rich diamictons; massive to stratified, matrix- and clast-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; drapes till and other glacial deposits; deposited by in situ melting from stagnant cold-based ice and modified by meltwater; evidence for ice collapse includes slump structures, kettle lakes and irregular topography; potential source of aggregate when material is gravel rich; generally suitable for infrastructure placement.
Tm Moraine ridges: sand, silt and clay-rich diamictons; massive, matrix-supported; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 m high, major moraines up to 12.5 km in length and 10 m high; ridges drape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional push moraines; major ridges features are large recessional end moraines and ice-thrust ridges; generally suitable for infrastructure placement.
Ts Streamlined till: silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 8 km in length; generally less than 50 m wide and 20 m high, formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glaciogenic deformation of sediment beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement.

- Geological boundary (Confidence: approximate)
Major moraine ridge (unspecified)
Other moraine ridge (unspecified)
Esker ridge (sense: unknown or unspecified)
Drumlin ridge
Major meltwater channel scarp
Minor meltwater channel central axis (unspecified; sense: known)
Station location (ground observation or stratigraphic section)

Recommended citation
Huntley, D.H., Hickin, A.S., Chow, W., and Mirmohammadi, M., 2013. Surficial geology, Trail Lake, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 123 (preliminary), scale 1:50 000. doi:10.4095/292400

CANADIAN GEOSCIENCE MAP 123 SURFICIAL GEOLOGY TRAIL LAKE BRITISH COLUMBIA 1:50 000

Authors: D.H. Huntley, A.S. Hickin, W. Chow, and M. Mirmohammadi. Cartography by W. Chow. Initiative of the Geological Survey of Canada, conducted under the auspices of the Yukon Basin Project as part of Natural Resources Canada's Geomapping for Energy and Minerals (GEM) program. Map projection Universal Transverse Mercator, zone 10, North American Datum 1983. Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations in feet above mean sea level. Magnetic declination 2013, 20°19'E, decreasing 22' annually. The Geological Survey of Canada welcomes corrections or additional information from users. This publication is available for free download through GEOSCAN (http://geoscan.nrcan.gc.ca/).

Preliminary publications in this series have not been scientifically edited.